

Transformer-based AI approach to unravel long-term, time-dependent prognostic complexity in patients with advanced NSCLC and PD-L1 $\geq 50\%$: insights from the pembrolizumab 5-year global registry

Alessio Cortellini ^{1,2,3}, Valentina Santo,^{1,4} Leonardo Brunetti,^{1,3} Edoardo Garbo,⁴ David J Pinato ^{3,5}, Giulia La Cava,¹ Jarushka Naidoo,^{6,7,8} Artur Katz,⁹ Monica Loza,¹⁰ Joel W Neal,¹⁰ Carlo Genova,¹¹ Scott Gettinger,¹² So Yeon Kim ¹², Ritujith Jayakrishnan,¹² Talal El Zarif,¹² Marco Russano,¹ Federica Pecci,⁴ Alessandro Di Federico,⁴ Joao V Alessi ¹³, Michele Montrone ¹⁴, Dwight H Owen ^{15,16}, Sara Ramella,^{2,17} Diego Signorelli,¹⁸ Mary Jo Fidler,¹⁹ Mingjia Li ¹⁵, Andrea Camerini,²⁰ Balazs Halmos ²¹, Bruno Vincenzi,^{1,2} Giulio Metro,²² Francesco Passiglia,²³ Sai Yendamuri ²⁴, Annalisa Guida,²⁵ Michele Ghidini,²⁶ Antonio D'Alessio ³, Giuseppe L Banna,²⁷ Claudia A M Fulgenzi,³ Salvatore Grisanti,²⁸ Francesco Grossi,²⁹ Armida D'Incecco,³⁰ Eleni Josephides,³¹ Mieke Van Hemelrijck,³² Alessandro Russo,³³ Alain Gelibter,³⁴ Gianpaolo Spinelli,³⁵ Monica Verrico,³⁶ Bartłomiej Tomasik,³⁷ Raffaele Giusti ³⁸, Kirsty Balachandran,²³ Emilio Bria ^{29,30}, Martin Sebastian,³¹ Maximilian Rost,³¹ Martin Forster,³⁹ Uma Mukherjee,³⁹ Lorenza Landi,³³ Francesca Mazzoni,³⁴ Avinash Aujayeb,³⁵ Manuel Dupont,³⁶ Alessandra Curioni-Fontecedro,³⁶ Rita Chiari,⁴⁰ Vincenzo Sforza,⁴¹ Marcello Tiseo ^{42,43}, Alex Friedlaender,⁴⁴ Alfredo Addeo,⁴⁵ Federica Zoratto,⁴⁶ Michele De Tursi,⁴⁷ Luca Cantini,⁴⁸ Elisa Roca,⁴⁹ Giannis Mountzios ⁵⁰, Danilo Rocco,⁵¹ Luigi Della Gravara,⁵¹ Sukumar Kalvapudi,²⁴ Alessandro Inno,⁵² Paolo Bironzo,²³ Rafael Di Marco Barros,³¹ David O'Reilly,⁶ Orla Fitzpatrick,⁶ Eleni Karapanagiotou,³¹ Isabelle Monnet,⁵³ Javier Baena,⁵⁴ Marianna Macerelli,⁵⁵ Aida Piedra,⁵⁶ Francesco Agustoni,⁵⁷ Diego Luigi Cortinovis ^{58,59}, Giuseppe Tonini,^{1,2} Gabriele Minuti,³³ Chiara Bennati,⁶⁰ Laura Mezquita ^{61,62}, Teresa Gorria,⁶³ Alberto Servetto,⁶⁴ Teresa Beninato,⁶⁵ Giuseppe Lo Russo,⁶⁵ Arsela Prelaj ^{65,66}, Andrea De Giglio ⁶⁷, Jacobo Rogado,⁶⁸ Laura Moliner,⁶⁹ Ernest Nadal ⁶⁹, Federica Biello,⁵ Frank Aboubakar Nana,⁷⁰ Anne-Marie Dingemans,⁷¹ Joachim G J V Aerts ⁷¹, Roberto Ferrara,^{72,73} Taher Abu Hejleh,^{74,75} Kazuki Takada,⁷⁶ Abdul Rafeh Naqash,⁷⁷ Marina Chiara Garassino ⁷⁸, Solange Peters,⁷⁹ Heather A Wakelee,¹⁰ Amin H Nassar ¹², Biagio Ricciuti ⁴, Paolo Soda,^{80,81} Camillo Maria Caruso,⁸⁰ Valerio Guarrasi⁸⁰

To cite: Cortellini A, Santo V, Brunetti L, et al. Transformer-based AI approach to unravel long-term, time-dependent prognostic complexity in patients with advanced NSCLC and PD-L1 $\geq 50\%$: insights from the pembrolizumab 5-year global registry. *Journal for ImmunoTherapy of Cancer* 2025;13:e012423. doi:10.1136/jitc-2025-012423

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/jitc-2025-012423>).

CMC and VG contributed equally.

CMC and VG are joint senior authors.

Accepted 31 July 2025



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

For numbered affiliations see end of article.

Correspondence to

Dr Alessio Cortellini;
alessiocortellini@gmail.com

ABSTRACT

Background With nearly one-third of patients with advanced non-small cell lung cancer (NSCLC) and PD-L1 Tumor Proportion Score $\geq 50\%$ surviving beyond 5 years following first-line pembrolizumab, long-term outcomes challenge traditional paradigms of cancer prognostication. The emergence of non-cancer-related factors and time-

dependent trends underscores the need for advanced analytical frameworks to unravel their complex interplay.

Methods We analyzed the Pembro-real 5Y registry, a global real-world dataset of 1050 patients treated across 61 institutions in 14 countries with a long-term follow-up and a large panel of baseline variables. Two complementary approaches were employed: ridge

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Immune checkpoint inhibitors have redefined the prognosis of patients with advanced non-small cell lung cancer (NSCLC), particularly those with PD-L1 expression $\geq 50\%$, with nearly one-third surviving beyond 5 years when treated with pembrolizumab. However, conventional models struggle to capture the long-term, time-dependent influence of host and non-cancer-related factors, and there is a need for innovative analytical frameworks that address data complexity and missingness in real-world settings.

WHAT THIS STUDY ADDS

⇒ This study applies a novel integrative framework combining ridge regression and a transformer-based artificial intelligence (AI) model (not another imputation method) to analyze a large, global, real-world cohort of patients with advanced NSCLC treated with first-line pembrolizumab. It reveals that the prognostic relevance of clinical features evolves over time: while Eastern Cooperative Oncology Group Performance Status, steroid use, and metastatic burden dominate early mortality, long-term outcomes are more strongly influenced by systemic health indicators such as dyslipidemia, body mass index, and absence of cardiovascular disease. The AI model successfully handles missing data without imputation and uncovers dynamic, non-linear patterns not detected by traditional methods.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ This study supports a shift toward holistic, time-aware patient management strategies in immunotherapy-treated NSCLC. It emphasizes the need to integrate dynamic risk assessment tools into clinical practice and highlights the growing importance of systemic health and comorbidities in survivorship care. The findings also encourage future research to validate AI-based models and adopt explainable, real-world compatible methodologies in prognostication.

regression, chosen for its ability to address multicollinearity while retaining interpretability, and not another imputation method (NAIM), a transformer-based artificial intelligence model designed to handle missing data without imputation. Endpoints included risk of death at 6, 12, 24, 60 months and 5-year survival.

Results The ridge regression model achieved a c-statistic of 0.66 (95% CI: 0.59 to 0.72) for the risk of death and an area under the curve (AUC) of 0.72 (95% CI: 0.65 to 0.78) for 5-year survival, identifying Eastern Cooperative Oncology Group Performance Status (ECOG-PS) ≥ 2 , increasing age, and metastatic burden as primary risk factors. However, wide CIs for some predictors highlighted statistical instability. NAIM demonstrated robust handling of missing data, with a c-index of 62.98 ± 2.11 for risk of death and an AUC of 60.52 ± 3.71 for 5-year survival. The comprehensive SHapley Additive exPlanations analysis revealed dynamic, time-dependent patterns, with early mortality dominated by acute factors (eg, ECOG-PS, steroids) and long-term outcomes increasingly influenced by systemic health markers (eg, absence of hypertension, increasing body mass index). Unexpected insights included the protective role of dyslipidemia (but not statins) and the nuanced impact of smoking status, reflecting evolving disease dynamics and host-tumor interplay.

Conclusions Our integrative framework illuminates the complexity of long-term outcomes in patients with NSCLC treated with pembrolizumab, uncovering dynamic, non-linear prognostic trends. This analysis provides insights into patient trajectories, emphasizing the need for holistic, long-term management strategies.

INTRODUCTION

The treatment landscape of non-small cell lung cancer (NSCLC) has undergone a profound transformation with the advent of immune checkpoint inhibitors (ICIs) and targeted therapies. These advances have established unprecedented long-term survival outcomes, particularly in patients with oncogene-addicted tumors and those with high PD-L1 expression treated with ICIs.^{1,2} In the context of ICIs and chemo-immunotherapy, survival rates of $\approx 20\%$ at 5 years have become a consistent reality in advanced NSCLC populations,^{3–5} with single-agent PD-1 inhibition achieving 5-year survival rates of 30% in patients with PD-L1 expression $\geq 50\%$.^{6,7}

This shift is leading the oncology community to a re-evaluation of traditional paradigms in the prognostication and management of patients with NSCLC. Historically, the expected survival of advanced-stage patients was limited to less than 1 year,⁸ with baseline prognostic factors such as performance status (PS) and metastatic burden exerting a proportional and consistent impact on outcomes throughout the disease trajectory.⁹ However, as survival horizons expand, non-cancer-related, host-specific factors are emerging as critical determinants of long-term outcomes.

The interaction between host factors and tumor biology is particularly pronounced in the context of ICIs, which modulate the immune-tumor interplay. Factors linked to the metabolic status,^{10–13} body composition,^{14–17} and systemic inflammation^{18,19} are not only relevant to immune function but also influence treatment efficacy and patient trajectories. As such, the evolving complexity of patient outcomes demands novel analytical methodologies capable of capturing dynamic, non-linear, and time-sensitive trends.

The Pembro-real 5Y is a comprehensive real-world, global, dataset derived from 61 institutions including patients with stage IV NSCLC with PD-L1 Tumor Proportion Score (TPS) of $\geq 50\%$ treated with first-line pembrolizumab monotherapy outside of clinical trials.²⁰ It captures detailed clinical, demographic, and treatment-related data, with a particular emphasis on long-term outcomes, including overall survival (OS) at 5 years.

In this study, we propose an integrative approach to the Pembro 5Y dataset that combines conventional statistical models with advanced artificial intelligence (AI)-based methodologies.

This dual framework aims to address the growing granularity of available clinical data by combining the traditional need for interpretable metrics with the modern imperative for dynamic risk stratification, leveraging conventional and AI-driven methodologies to uncover non-linear prognostic patterns that extend beyond traditional oncologic determinants.

METHODS

Study design

The objective of this study is to uncover novel insights into the risk of death and the determinants of 5-year survival

in patients with advanced stage NSCLC with PD-L1 TPS $\geq 50\%$ treated with first-line pembrolizumab, leveraging conventional and AI-driven methodologies. With this approach, we aim to delineate both expected and unexpected prognostic factors, offering a comprehensive understanding of patient trajectories and informing personalized strategies in this rapidly evolving field.

The Pembro-real 5Y is a large real-world global dataset that includes 1050 patients with stage IV NSCLC and PD-L1 TPS $\geq 50\%$, treated with first-line pembrolizumab at 61 institutions across 14 countries and has already been reported in detail.²⁰ The primary eligibility criteria included receiving first-line pembrolizumab monotherapy outside of clinical trials, having a PD-L1 TPS of $\geq 50\%$, and starting treatment by May 31, 2018. To ensure at least 5 years of follow-up for long-term responders, the minimum data cut-off for patients still alive was set at May 1, 2023.

The 5-year survival rate, defined as the crude rate of patients alive at the 5-year mark from the treatment initiation date, and OS, defined as the time from treatment initiation to death or loss to follow-up, were selected as the clinical endpoints of interest.

To address the complexity of the dataset, characterized by long-term follow-up, extensive baseline information with inherent collinearity, and significant missing data, we adopted a dual analytical approach: conventional statistical modeling using ridge regression and a transformer-based AI model. These methods were applied to the same panel of baseline variables (provided in the online supplemental methods) including clinicopathologic oncological characteristics, demographics, concomitant medications, and comorbidities. Ridge regression was chosen because it allows us to address the challenges of multicollinearity and high-dimensional datasets without eliminating potentially important variables and maintaining interpretability. To complement the conventional analysis, we employed NAIM (not another imputation method), a transformer-based AI model specifically designed for resilience to missing data. This model was originally developed and tested in the context of OS prediction in NSCLC patients and has demonstrated superior performance over conventional imputation-based approaches in prior studies.^{21–23} While external validation in independent cohorts is still lacking, its architecture offers promising features for real-world application, particularly in datasets characterized by missingness and high-dimensional clinical information. Unlike traditional methods requiring complete datasets or imputations, NAIM uses a masked self-attention mechanism to make predictions solely from available data without imputing missing values. Additionally, explainable AI techniques, specifically the SHAP (SHapley Additive exPlanations) method,²⁴ were applied to quantify each feature's contribution to model predictions. SHAP values, rooted in game theory and economics, provide an interpretable measure of each variable's effect on survival or death. This nuanced approach captures time-dependent and non-linear interactions, offering insights

unavailable through conventional regression models. By combining these methodologies, we balanced the interpretability and clinical applicability of conventional statistics with the flexibility and depth of AI, aiming to extract meaningful patterns and provide actionable insights into the predictors of long-term outcomes in NSCLC patients.

To further analyze the importance of the included variables in determining the risk of death over time, we used a graphical representation of the cumulative sum of the absolute contributions of the features across four pre-established time points: 6 months, 12 months, 24 months and 60 months. This approach aggregates the absolute SHAP values, allowing for the identification of feature importance independent of specific time points. Visualizing these cumulative sums allows us to assess which features consistently play a dominant role in influencing predictions throughout the observed period. This representation provides a comprehensive view of feature relevance, aiding in the interpretation of the model's behavior and offering insights into the underlying clinical factors that drive survival outcomes.

Statistical analysis

We used descriptive statistics to report baseline clinicopathologic features of interest, assessing differential distribution through the χ^2 test. The 5-year survival rate was reported as a crude rate with 95% CIs. Median follow-up was estimated using the reverse Kaplan-Meier method, while the median OS was computed using the Kaplan-Meier method and compared with the log-rank test.

In the ridge regression analysis, given the substantial heterogeneity in missingness across the baseline variables (ranging from 1.4% to 89.6%), we opted for a pragmatic and reproducible strategy in the modeling phase. Specifically, categorical variables with residual missingness were retained with “unknown/not-tested” encoded as an explicit factor level. This allowed us to preserve the full analytic sample and avoid the introduction of uncertain or artificial imputations, particularly in variables where the proportion of missing data exceeded commonly accepted thresholds for reliable multiple imputation. This approach aligns with the real-world nature of the dataset, where data incompleteness may reflect systemic differences in clinical documentation, patient complexity, or diagnostic work-up and may carry independent prognostic information. Consequently, HRs reported for the “unknown/not-tested” levels in the regression models represent the impact of such data absence, rather than true biological strata. The extent of missingness for each baseline variable is detailed in table 1.

For the risk of death, a Cox proportional hazards model with ridge regularization was applied to manage multicollinearity and penalize large coefficients. Cross-validation was used to determine the optimal penalty parameter (λ), minimizing the partial likelihood deviance. Variables with non-zero coefficients were subsequently refitted into a standard Cox model to estimate HRs and 95% CIs. Model performance was evaluated using the concordance

Table 1 Patients' characteristics of the overall cohort

	Overall study population No 1050 (%)		Overall study population No 1050 (%)
Age, (years)		Baseline corticosteroids	
Median (range)	69 (31–92)	None	845 (80.5)
		<10mg pred	114 (10.8)
		≥10mg pred	91 (8.7)
Sex		Baseline proton pump inhibitors	
Female	422 (40.2)	No	598 (57.0)
Male	628 (59.8)	Yes	400 (38.1)
		Unknown	52 (5.0)
Ethnicity		Baseline antibiotics	
White	870 (82.9)	No	943 (89.8)
Black/African-American	31 (3.0)	Yes	107 (10.2)
Asian	29 (2.7)		
Hispanic	9 (0.8)		
Others	26 (2.5)		
Unknown	85 (8.1)		
WHO BMI category		Baseline statins	
Obese	162 (15.4)	No	681 (64.9)
Overweight	286 (27.2)	Yes	313 (29.8)
Normal-weight	418 (39.8)	Unknown	56 (5.3)
Under-weight	41 (3.9)		
Unknown	143 (13.7)		
ECOG-PS		Baseline metformin	
0–1	859 (81.8)	No	887 (84.5)
≥2	161 (15.3)	Yes	124 (11.8)
Unknown	30 (2.9)	Unknown	39 (3.7)
Histology		Other glucose-lowering medications	
Squamous	238 (22.8)	No	787 (75.0)
Adenocarcinoma	768 (73.5)	Yes	64 (6.1)
Others/NOS	39 (3.7)	Unknown	199 (19.0)
Smoking status		Hypertension	
Current smokers	280 (26.7)	No	469 (44.7)
Former smokers	660 (62.9)	Yes	523 (49.8)
Never smokers	95 (9.0)	Unknown	58 (5.5)
Unknown	15 (1.4)		
PD-L1 TPS		Myocardial infarction	
≥90%	559 (53.2)	No	858 (81.7)
50%–89%	281 (26.8)	Yes	128 (12.2)
Not specified	210 (20.0)	Unknown	64 (6.1)
CNS metastases		Other cardiovascular conditions	
No	841 (80.1)	No	715 (68.1)
Yes	209 (19.9)	Yes	271 (25.8)
		Unknown	64 (6.1)
Bone metastases		Type 2 diabetes	
No	686 (65.3)	No	809 (77.0)

Continued

Table 1 Continued

	Overall study population		Overall study population	
	No 1050 (%)		No 1050 (%)	
Yes	364 (34.7)	Yes	177 (16.9)	
		Unknown	64 (6.1)	
Liver metastases		Pulmonary disease		
No	885 (84.3)	No	711 (67.7)	
Yes	165 (15.7)	Yes	275 (26.2)	
		Unknown	64 (6.1)	
Lung metastases		Dyslipidemia		
No	396 (37.7)	No	666 (63.4)	
Yes	654 (62.3)	Yes	314 (29.9)	
		Unknown	70 (6.7)	
Pleural metastases		Autoimmune diseases		
No	787 (75.0)	No	933 (88.9)	
Yes	263 (25.0)	Yes	53 (5.0)	
		Unknown	64 (6.1)	
Adrenal glands metastases		Other comorbidities		
No	839 (79.9)	No	563 (53.6)	
Yes	211 (20.1)	Yes	374 (35.6)	
		Unknown	113 (10.8)	
Other metastatic sites		–	–	
No	900 (85.7)	–	–	
Yes	150 (14.3)			
Number of metastatic sites		–	–	
≤3	946 (90.1)	–	–	
>3	104 (9.9)			
EGFR mutation status		–	–	
No	890 (84.8)	–	–	
Yes	19 (1.8)			
Not-tested	141 (13.4)			
ALK translocation status		–	–	
No	883 (84.1)	–	–	
Yes	6 (0.6)			
Not-tested	161 (15.3)			
ROS-1 translocation status		–	–	
No	710 (67.6)	–	–	
Yes	9 (0.9)			
Not-tested	331 (31.5)			
KRAS mutation status		–	–	
No	367 (35.0)	–	–	
Yes	224 (21.3)			
Not-tested	459 (43.7)			
BRAF mutation status		–	–	
No	491 (46.8)	–	–	
Yes	26 (2.5)			
Not-tested	533 (50.7)			

Continued

Table 1 Continued

	Overall study population		Overall study population
	No 1050 (%)		No 1050 (%)
Tumor mutational burden	–		–
Non-high	92 (8.8)	–	–
High	17 (1.6)		
Not-tested	941 (89.6)		

Missing values were reported as “unknown”, “not-specified” or “not-reported” as appropriate. ALK, anaplastic lymphoma kinase; BMI, body mass index; CNS, central nervous system; ECOG-PS, Eastern Cooperative Oncology Group Performance Status; EGFR, epidermal growth factor receptor; KRAS, Kirsten rat sarcoma virus; NOS, not otherwise specified; PD-L1, programmed death-ligand 1; pred, prednisone; ROS-1, proto-oncogene tyrosine-protein kinase ROS; TPS, Tumor Proportion Score.

index (C-index). For the 5-year survival analysis, logistic ridge regression was employed on the same imputed datasets and covariates, with the binary outcome indicating survival at 5 years. Cross-validation identified the optimal λ , and selected variables were refitted into a logistic regression model to estimate ORs and 95% CIs. Model discrimination was assessed via the area under the receiver operating characteristic curve (AUC) and c-statistics. These analyses were performed using the R-studio software, R Core Team (2021). R: A language and environment for statistical computing implemented with the *glmnet*, *survival* packages, ensuring robust handling of multicollinearity and providing interpretable, clinically relevant metrics like ORs and HRs.

As previously described, the NAIM method is a transformer-based AI model that does not require imputation. Instead, it leverages a masked self-attention mechanism that allows the model to learn directly from partially observed data, using only the available features without artificially reconstructing missing values. Numerical features were normalized to (0,1), and categorical variables were encoded numerically. The first block of the model, called feature embedding, encoded these inputs as embeddings, which involves assigning trainable vectors to each possible clinical characteristic. At this stage, missing data were already addressed by assigning it a specific, non-trainable vector. Following this, the core of the model used masked self-attention to make predictions based exclusively on the available patient data.

The analyses were performed using a fivefold cross-validation strategy, stratified on the target variable to maintain the same class proportions in the training and testing sets. The model, which is implemented in Python using the PyTorch library, was evaluated using various metrics, including AUC, accuracy, F1-score, Matthews correlation coefficient (MCC), and G-mean, each of which captures a different aspect of predictive performance. Furthermore, in addition to the results obtained on the test samples, we report the training performance to assess the correlation between the target variable and the various features considered.

SHAP values were then used to provide an interpretable measure of each variable's contribution in defining clinical outcome and were visualized through summary plots. Features were ordered by their absolute contribution, with high values (red) and low values (blue) positioned to indicate their influence on outcomes. For instance, red dots on the right side of the plot imply a positive association with the outcome, while blue dots on the left suggest negative association. Missing values were represented as gray dots. [Figure 1](#) provides a visual overview of the methodological framework and the interpretation of the results.

RESULTS

Cohort characteristics

The Pembro-real 5Y cohort has been extensively described elsewhere²⁰; overall 1063 consecutive patients were entered in the registry (online supplemental table 1 reports the number of patients entered by each participating institution) with a final eligible population of 1050 patients after the exclusion of 13 patients lost to follow-up.

Although the overall cohort has been previously described, a comprehensive overview of the clinicopathologic characteristics of the study cohort, including the absolute number and percentage of missing values for each baseline variable, is reported in [table 1](#).

At the median follow-up period of 70.3 months (95% CI: 69.0 to 70.9) and a total of 805 death events, 282 patients were alive at the 5-year landmark, resulting in a 5-year survival rate of 26.9% (95% CI: 23.8% to 30.2%) while the median OS for the study population was 21.8 months (95% CI: 19.1 to 25.7, 805 events).

Ridge regression analysis

The ridge regression analysis for the risk of death was conducted using an optimal penalty parameter (λ) determined via cross-validation, which was 0.631. The multivariate model is reported in [table 2](#) and demonstrated moderate discriminatory ability with a c-statistic of 0.66 (95% CI: 0.59 to 0.72). Significant predictors of

Analytical framework




Figure 1 Visual summary of the methodological pipeline used to analyze long-term outcomes of patients with advanced NSCLC (PD-L1 $\geq 50\%$) treated with first-line pembrolizumab in the Pembro-Real 5Y Registry. Two parallel approaches were adopted: Conventional modeling using penalized ridge regression for both 5-year survival (logistic) and overall survival (Cox), producing interpretable ORs and HRs with associated CIs, and model performance metrics (eg, C-statistic, AUC). NAIM (AI model) employing a transformer-based deep learning architecture to model time-dependent survival dynamics using SHAP values for feature interpretability and multiple performance metrics (AUC, F1-score, Matthews correlation coefficient [MCC], etc). Outputs were compared side-by-side. Shared predictors (eg, ECOG-PS, corticosteroid use) reinforced robustness, while NAIM identified additional time-sensitive or non-linear predictors (eg, BMI, dyslipidemia). This integrative approach supports both baseline prognostication and longitudinal survivorship strategies. AUC, area under the curve; BMI, body mass index; CV, cardiovascular; ECOG-PS, Eastern Cooperative Oncology Group Performance Status; NAIM, Not Another Imputation Method; NSCLC, non-small cell lung cancer; SHAP, SHapley Additive exPlanations; TMB, Tumor Mutational Burden.

increased mortality risk included Eastern Cooperative Oncology Group Performance Status (ECOG-PS) ≥ 2 (HR 2.06, 95% CI: 1.67 to 2.53), the use of both low dose (HR 1.64, 95% CI: 1.29 to 2.08) and high dose steroids (HR 1.37; 95% CI: 1.04 to 1.81), the presence of liver (HR 1.35, 95% CI: 1.07 to 1.69), bone (HR 1.34, 95% CI: 1.13

to 1.58), and pleural metastases (HR 1.21, 95% CI: 1.01 to 1.45), the presence of other cardiovascular comorbidities (HR 1.22, 95% CI: 1.01 to 1.46), being never smoker (HR 1.35, 95% CI: 1.01 to 1.81), the exposure to proton pump inhibitors (PPI) at baseline (HR 1.20, 95% CI: 1.02 to 1.40) and increasing age (HR 1.01 per

Table 2 Ridge regression multivariable analyses for the risk of death (overall survival—using Cox proportional hazard model) and the 5-year survival state (using logistic regression)

Variable (data missingness)	Overall survival (risk of death)	5year survival (probability of being alive)
	HR (95% CI)	OR (95% CI)
Age (continuous)	1.01 (1.01 to 1.02)	0.97 (0.96 to 0.99)
Sex		
Male vs female	0.97 (0.83 to 1.14)	1.01 (0.73 to 1.39)
Ethnicity (8.1%)		
White	1	1
Black/African American	0.66 (0.41 to 1.06)	1.68 (0.69 to 3.93)
Asian	1.10 (0.72 to 1.67)	0.42 (0.11 to 1.21)
Hispanic	0.29 (0.11 to 1.77)	4.44 (0.94 to 23.66)
Others	0.96 (0.61 to 1.52)	0.42 (0.11 to 1.20)
Unknown	1.03 (0.75 to 1.39)	0.75 (0.39 to 1.39)
WHO BMI (13.7%)		
Normal weight	1	1
Underweight	1.31 (0.87 to 1.96)	0.89 (0.35 to 2.05)
Overweight	1.07 (0.89 to 1.29)	0.94 (0.63 to 1.38)
Obese	0.96 (0.76 to 1.20)	1.09 (0.69 to 1.72)
Unknown	0.98 (0.76 to 1.27)	1.25 (0.73 to 2.12)
ECOG-PS (2.9%)		
0–1	1	1
≥2	2.06 (1.67 to 2.52)	0.39 (0.23 to 0.65)
Unknown	1.31 (0.79 to 2.17)	0.70 (0.21 to 2.09)
Histology		
Squamous	1	1
Adenocarcinoma	1.06 (0.88 to 1.29)	1.18 (0.68 to 1.83)
Others/NOS	1.03 (0.66 to 1.62)	1.62 (0.74 to 3.77)
Smoking status (1.4%)		
Current smoker	1	1
Former smoker	1.04 (0.87 to 1.24)	0.81 (0.57 to 1.15)
Never smoker	1.35 (1.01 to 1.81)	0.57 (0.30 to 1.08)
Unknown	1.44 (0.77 to 2.69)	0.09 (0.01 to 0.60)
PD-L1 tumor proportion score (20.0%)		
50–89	1	1
≥90	0.95 (0.79 to 1.13)	1.08 (0.76 to 1.56)
Not reported	0.92 (0.74 to 1.13)	1.16 (0.24 to 1.76)
CNS metastases		
Yes vs No	1.04 (0.85 to 1.28)	0.90 (0.42 to 1.37)
Liver metastases		
Yes vs No	1.35 (1.07 to 1.69)	0.63 (0.47 to 1.06)
Bone metastases		
Yes vs No	1.32 (1.12 to 1.56)	0.67 (0.58 to 0.95)
Lung metastases		
Yes vs No	1.04 (0.88 to 1.23)	1.04 (0.94 to 1.48)
Pleural metastases		

Continued

Table 2 Continued

Variable (data missingness)	Overall survival (risk of death)	5year survival (probability of being alive)
	HR (95% CI)	OR (95% CI)
Yes vs No	1.22 (1.02 to 1.45)	0.63 (0.61 to 0.93)
Adrenal glands metastases		
Yes vs No	1.11 (0.91 to 1.35)	0.95 (0.73 to 1.42)
Other metastatic sites		
Yes vs No	0.94 (0.75 to 1.18)	1.49 (0.11 to 2.24)
Number of metastatic sites		
>3 vs ≤3	1.12 (0.80 to 1.55)	0.68 (NA-1.47)
Corticosteroids at baseline		
No	1	1
<10mg prednisolone or eq.	1.64 (1.28 to 2.06)	0.70 (0.41 to 1.18)
≥10mg prednisolone or eq.	1.37 (1.04 to 1.81)	0.76 (0.46 to 1.36)
Baseline proton pump inhibitors (5.0%)		
No	1	1
Yes	1.20 (1.03 to 1.41)	0.85 (0.41 to 1.19)
Unknown	1.25 (0.71 to 2.21)	1.22 (0.64 to 3.49)
Baseline antibiotics		
Yes vs No	1.10 (0.86 to 1.41)	0.81 (0.37 to 1.39)
Baseline statins (5.3%)		
No	1	1
Yes	1.01 (0.80 to 1.25)	1.01 (0.30 to 1.58)
Unknown	0.99 (0.60 to 1.64)	0.83 (0.34 to 2.19)
Baseline metformin (3.7%)		
No	1	1
Yes	1.05 (0.78 to 1.41)	0.65 (0.58 to 1.22)
Unknown	0.70 (0.44 to 1.07)	1.35 (0.40 to 3.12)
Other glucose lowering medications (19.0%)		
No	1	1
Yes	1.27 (0.88 to 1.84)	0.73 (0.31 to 1.66)
Unknown	1.05 (0.83 to 1.34)	0.91 (0.55 to 1.47)
EGFR mutational status (13.4%)		
Wild type	1	1
Mutant	1.14 (0.64 to 2.04)	0.91 (0.26 to 2.87)
Not tested	1.42 (0.95 to 2.14)	0.64 (0.02 to 1.56)
ALK translocation status (15.3%)		
Wild type	1	1
Translocated	1.41 (0.55 to 3.58)	0.54 (0.37 to 3.99)
Not tested	1.08 (0.71 to 1.64)	0.91 (0.31 to 2.16)
ROS-1 translocation status (31.5%)		
Wild type	1	1
Translocated	0.68 (0.31 to 1.48)	1.65 (0.50 to 7.28)
Not tested	0.94 (0.73 to 1.18)	0.84 (0.38 to 1.40)
KRAS mutational status (47.3%)		
Wild type	1	1

Continued

Table 2 Continued

Variable (data missingness)	Overall survival (risk of death)	5year survival (probability of being alive)
	HR (95% CI)	OR (95% CI)
Mutant	0.83 (0.67 to 1.04)	1.34 (0.88 to 2.04)
Not tested	1.03 (0.78 to 1.33)	0.96 (0.56 to 1.65)
BRAF mutational status (50.7%)		
Wild type	1	1
Mutant	0.99 (0.61 to 1.57)	0.81 (0.26 to 2.19)
Not tested	1.15 (0.89 to 1.48)	0.88 (0.52 to 1.46)
Tumor mutational burden (89.6%)		
Non-high	1	1
High	0.60 (0.29 to 1.24)	2.99 (0.92 to 9.93)
Not tested	1.04 (0.76 to 1.40)	1.08 (0.62 to 1.97)
Hypertension (5.5%)		
No	1	1
Yes	1.01 (0.85 to 1.19)	0.78 (0.54 to 1.12)
Unknown	1.47 (0.41 to 5.36)	0.91 (0.08 to 12.17)
Myocardial infarction (6.1%)		
No	1	1
Yes	0.93 (0.73 to 1.18)	0.88 (0.52 to 1.46)
Unknown	1.32 (0.05 to 35.74)	0.81 (0 to NA)
Other cardiovascular conditions (6.1%)		
No	1	1
Yes	1.22 (1.01 to 1.46)	0.74 (0.49 to 1.10)
Unknown	1.14 (0.32 to 4.00)	1.05 (0.10 to 12.10)
Type 2 diabetes (6.1%)		
No	1	1
Yes	0.80 (0.59 to 1.09)	1.28 (0.70 to 2.32)
Unknown	0.29 (0.04 to 2.07)	1690577.00 (0.00 to NA)
Pulmonary disease (6.1%)		
No	1	1
Yes	0.87 (0.72 to 1.04)	1.01 (0.11 to 1.45)
Unknown	0.67 (0.22 to 2.01)	1.40 (0.91 to 19.44)
Dyslipidemia (6.7%)		
No	1	1
Yes	0.84 (0.67 to 1.04)	1.42 (0.11 to 2.24)
Unknown	0.78 (0.35 to 1.72)	0.70 (0.78 to 3.13)
Autoimmune diseases (6.1%)		
No	1	1
Yes	0.71 (0.50 to 1.01)	1.63 (0.84 to 3.11)
Unknown	6.04 (0.49 to 73.35)	0.01 (NA)
Other comorbidities (10.8%)		
No	1	1
Yes	0.92 (0.79 to 1.09)	1.07 (0.77 to 1.49)
Unknown	0.61 (0.41 to 0.88)	2.63 (1.25 to 5.53)

Continued

Table 2 Continued

Variable (data missingness)	Overall survival (risk of death)	5year survival (probability of being alive)
	HR (95% CI)	OR (95% CI)
Missing values were reported as “unknown”, “not-specified” or “not-reported” as appropriate. ALK, anaplastic lymphoma kinase; BMI, body mass index; CNS, central nervous system; ECOG-PS, Eastern Cooperative Oncology Group Performance Status; EGFR, epidermal growth factor receptor; KRAS, Kirsten rat sarcoma virus; NOS, not otherwise specified; PD-L1, programmed death-ligand 1.		

unit increase; 95% CI: 1.00 to 1.02). Conversely, Hispanic ethnicity (HR 0.28, 95% CI: 0.11 to 0.74) was identified as a protective factor while the absence of information on other comorbidities was associated with decreased risk of death (HR 0.60, 95% CI: 0.41 to 0.88). Variables with wide CIs were noted, reflecting statistical instability due to sparse events. Online supplemental figure 1 summarizes the ridge regression coefficients, ranking ECOG PS ≥ 2 as the most influential variable (coefficient: 0.39), followed by missing information on the smoking status (0.24) and low-dose steroid use (0.23), while Hispanic ethnicity (−0.47) and TMB-high (−0.24) were associated with the lowest coefficients.

The ridge regression analysis for the 5-year survival state used an optimal lambda of 0.203, as determined by cross-validation. The model achieved an AUC of 0.72 (95% CI: 0.65 to 0.78), indicating acceptable discriminatory performance. Statistically significant predictors associated with decreased probability of being alive at 5 years included ECOG PS ≥ 2 (OR 0.39; 95% CI: 0.22 to 0.65), increasing age (OR 0.97 per year; 95% CI: 0.96 to 0.99), bone (OR 0.67, 95% CI: 0.46 to 0.95) and pleural metastases (OR 0.62, 95% CI: 0.42 to 0.91). In addition, missing information on the smoking status was associated with a reduced probability of 5-year survival (OR 0.09, 95% CI: 0.01 to 0.60). Conversely, missing information on other comorbidities was associated with an increased probability of being alive at 5 years (OR 2.63, 95% CI: 1.25 to 5.53) (table 2). Ridge regression coefficients are reported as online supplemental figure 2; missing information on the smoking status was associated with the lowest coefficient (−0.67), followed by ECOG PS ≥ 2 (−0.39) while Hispanic ethnicity (0.73) and TMB-high (0.51) were associated with the highest coefficients.

NAIM analysis

The NAIM model's performance for predicting the risk of death was evaluated across four distinct time points (6, 12, 24, and 60 months) and through cumulative feature importance analysis. The NAIM model for risk of death, using all features, demonstrated a c-index of 62.98 ± 2.11 on the training set and 61.24 ± 2.15 on the all-feature evaluation set, indicating moderate predictive performance. While the training c-index suggests some degree of overfitting, the consistency between training and evaluation

NAIM model for the prediction of the risk of death across the 4 pre-defined time-points




Figure 2 Histogram plot summarizing the cumulative SHAP values from the NAIM analysis for the risk of death. The length of each bar represents the SHAP value, indicating the relative importance of each variable within the model. Features were ordered by their absolute contribution. The c-index (% $\pm$ SD) was 79.76 ± 2.44 for the training set and 62.98 ± 2.11 for the overall model. Variable's definition and categorization details are reported in online supplemental methods. ALK, anaplastic lymphoma kinase; BMI, body mass index; CNS, central nervous system; ECOG-PS, Eastern Cooperative Oncology Group Performance Status; EGFR, epidermal growth factor receptor; GLM, glucose- lowering medications; NAIM, Not Another Imputation Method; NOS, not otherwise specified; PPI, proton pump inhibitors; pred, Prednisone; SHAP, SHapley Additive exPlanations; TMB, tumor mutational burden; TPS, Tumor Proportion Score.

results reinforces the robustness of the model's performance for this task.

For the cumulative importance (figure 2), the variables with the highest impact across all time points were ECOG-PS, steroid use, and the presence of bone metastases. These variables were consistently the strongest predictors of mortality, reflecting their clinical relevance in determining short-term and long-term risk. Further analysis of time-specific SHAP values (online supplemental figures 3–6) revealed notable temporal variations in the importance and directionality of individual variables. It must be noted that the decreasing magnitude of SHAP values over time may indicate reduced model confidence or predictive ability at intermediate time points (12 and 24 months). At 6 months, ECOG-PS ≥ 2 emerged as the strongest predictor of mortality, with high SHAP values underscoring its acute prognostic significance. Steroid use and bone metastases followed as key contributors, reflecting their impact on short-term outcomes. However, as time progressed to 12 and 24 months, the magnitude of SHAP values for these predictors decreased markedly,

highlighting a diminishing influence on mortality risk over time. This decline suggests that short-term prognostic factors, such as ECOG-PS and steroids, lose relevance as other host-related or systemic variables gain prominence. Notably, variables such as PPI use and smoking status/pulmonary disease displayed dynamic trends over time. At 6 months, these factors had a more pronounced influence on mortality risk but showed variable contributions at later time points, reflecting evolving patterns of disease progression and competing risks. Meanwhile, BMI demonstrated increasing importance over time, transitioning from a marginal role at earlier time points to becoming a significant protective factor by 60 months.

The 5-year survival state analysis (figure 3) provided insights into predictors of long-term survival. The NAIM model for predicting the 5-year survival state, using all features, demonstrated an AUC of 60.52 ± 3.71 on the evaluation set and 78.53 ± 3.25 on the training set, indicating a notable degree of overfitting. Additional metrics for the evaluation set included an accuracy of 53.65 ± 2.97 , an F1-score of 45.43 ± 23.10 , an MCC of 10.72 ± 8.35 , and

NAIM model for the prediction of the 5-year survival




Figure 3 Paired dot plot and histogram and plot summarizing the SHAP values from the NAIM analysis for the 5-year survival. Features were ordered by their absolute contribution, with high values (red) and low values (blue) positioned to indicate their influence on outcomes. For instance, red dots on the right side of the plot imply a positive association with the probability of being alive at 5 years, while blue dots on the right side imply a negative association. Missing values were represented as gray dots. The length of each bar represents the SHAP value, indicating the relative importance of each variable within the model. The metrics for the model (% $\pm$ SD) were as follows: the training set achieved an AUC of 78.53 ± 3.25 , an accuracy of 74.41 ± 2.56 , an F1-score of 71.38 ± 2.82 , an MCC of 47.34 ± 3.41 , and a G-Mean of 64.89 ± 3.17 . In the evaluation set, the model demonstrated an AUC of 60.52 ± 3.71 , an accuracy of 53.65 ± 2.97 , an F1-score of 45.43 ± 23.10 , an MCC of 10.72 ± 8.35 , and a G-Mean of 36.71 ± 7.61 . Variables definition and categorization details are reported in online supplemental methods. ALK, anaplastic lymphoma kinase; AUC, area under the curve; BMI, body mass index; CNS, central nervous system; ECOG-PS, Eastern Cooperative Oncology Group Performance Status; EGFR, epidermal growth factor receptor; GLM, glucose-lowering medications; NAIM, not another imputation method; NOS, not otherwise specified; PPI, proton pump inhibitors; pred, Prednisone; Matthews correlation coefficient: MCC; SHAP, SHapley Additive exPlanations; TMB, tumor mutational burden; TPS, Tumor Proportion Score.

a G-mean of 36.71 ± 7.61 . The SHAP analysis identified several key predictors of 5-year survival. The absence of hypertension emerged as the most impactful variable, demonstrating a strong positive association with long-term survival. Similarly, the absence of other cardiovascular diseases and the presence of dyslipidemia were also strongly associated with an increased likelihood of being alive at 5 years. The absence of bone metastases further emerged as a significant predictor of long-term survival. In contrast, variables such as ECOG-PS ≥ 2 and steroid use were negatively associated with 5-year survival but were not prioritized among the most influential variables. Interestingly, BMI displayed a strong association with 5-year survival, with increasing BMI being linked to a higher likelihood of long-term survival, despite its mixed distribution in the SHAP plots.

DISCUSSION

Our study is unique in its application of both conventional and advanced AI-based methodologies to a large, granular dataset with long-term follow-up, capable of capturing and describing unconventional and non-proportional

baseline variable impacts over time. With one in three patients surviving beyond 5 years in the setting of NSCLC with PD-L1 TPS $\geq 50\%$,^{5,7} a comprehensive approach to care becomes essential, underscoring the growing importance of systemic health in the survivorship phase.

Previous studies leveraging AI and machine learning in advanced NSCLC have predominantly focused on static baseline factors and short-term outcomes.^{25,26} In contrast, our approach stands out by being the first to examine long-term outcomes and their temporal evolution in patients treated with ICIs.

The AI-based NAIM model and ridge regression provide complementary approaches for analyzing clinical outcome; each method offers unique strengths and limitations, reflecting their distinct capabilities to address the complexity of real-world data: long-term follow-up, extensive baseline granularity, significant missing data, and variable collinearity.

While ridge regression provides a static, interpretable estimate of the relative prognostic weight of each variable, the NAIM model allows for a dynamic view of how the relevance of individual features evolves over time.

By combining these methods, we aimed to gain deeper insights into both static and dynamic predictors of patient outcomes, uncovering temporal trends and unexpected patterns that inform personalized management in the evolving oncology landscape. For instance, the prognostic relevance of ECOG-PS and corticosteroid use is well-established in the immunotherapy setting, the emergence of dyslipidemia and the absence of cardiovascular disease as features associated with long-term benefit is a novel observation. These findings may reflect underlying systemic resilience and deserve further validation across diverse therapeutic contexts.

Ridge regression provides interpretable metrics essential for clinical translation. ECOG-PS $\geq$ 2, increasing age, metastatic burden, exposure to steroids, and PPI were associated with the risk of death and inversely associated with the 5-year survival state. These findings align with the known impact of PS, age, and metastatic sites on prognosis.⁹ However, the wide CIs of some variables reflect statistical instability due to low event rates or multicollinearity, undermining the robustness of these estimates. In addition, the observation that missing variables (other comorbidities “unknown” and smoking status “unknown”) and under-represented subgroups, such as Hispanic ethnicity, have a statistically significant impact on outcomes warrants careful interpretation as these estimates may reflect pre-existing correlations rather than true causal effects. Although prior studies have suggested potential biological or socioeconomic factors contributing to improved outcomes in Hispanic populations,^{27 28} these findings require further investigation. Moreover, while the ridge regression coefficients allow for ranking variables by magnitude, it is critical to note that variables with large coefficients but no significant impact may lead to misinterpretations of their true value, as their statistical insignificance could result from factors such as multicollinearity, limited event rates, or noise in the dataset rather than a genuine prognostic association.

The NAIM model provides insights into how prognostic factors influence outcomes not just at baseline, but at different time points during follow-up. This ability to capture temporal dynamics helps identify variables whose importance may increase, decrease, or shift over time, offering a more flexible and realistic understanding of patient trajectories under immunotherapy. The use of SHAP values offers a broader perspective by capturing the relative importance and directionality of variables across multiple predictions, unveiling non-proportional effects. However, the decreasing magnitude of SHAP values over time may indicate reduced model confidence or predictive ability at intermediate time points (12 and 24 months). This trend suggests that while the model effectively captures early predictors of mortality, its ability to sustain strong predictive insights diminishes as the follow-up period extends.

NAIM confirmed several classical prognostic variables while revealing their time-dependent trends, such as ECOG-PS and steroid exposure. ECOG-PS $\geq$ 2 emerged as

the strongest determinant of mortality risk within the first 6 months but diminished over time. By contrast, patients with ECOG-PS 0–1 exhibited increased mortality at 12, 24, and 60 months, reflecting the early dominance of poor PS in acute settings. A similar trend was observed for steroids, strongly associated with early mortality but reversing direction over time. By 5 years, non-steroid exposure was associated with survival, suggesting a diminishing prognostic role as disease trajectories stabilize. Similarly, the presence of bone metastases was confirmed as one of the strongest determinants, following a time-dependent pattern as well. Initially strongly associated with early mortality, its relevance diminished over time, while the absence of bone metastases remained a significant determinant of the 5-year survival state. Interestingly, NAIM revealed that bone metastases were the most significant metastatic site influencing outcomes, while other sites such as liver or CNS metastases, or the overall burden of disease, played a less prominent role.

NAIM also revealed unexpected patterns that differed from conventional expectations and underscored the unique capacity of the model to detect complex relationships. Before delving into technical details, it is worth noting that NAIM identified predictors whose prognostic importance evolved over time or emerged unexpectedly, such as cardiovascular health, metabolic factors, and smoking history. These findings suggest that some features gain relevance only at later stages of survivorship or through interactions not captured in traditional modeling frameworks. Unlike ridge regression, which retains increasing age as a significant prognostic factor, the NAIM analysis does not identify it as a relevant predictor over time. This discrepancy may stem from NAIM’s ability to account for collinearity, possibly allowing comorbidities to emerge as stronger determinants of long-term outcomes. Hypertension and the absence of other cardiovascular diseases gained disproportionate importance at 60 months, reflecting the cumulative impact of cardiovascular health on survivorship. Never-smokers and those without pulmonary disease were associated with increased mortality risk at 6 months, a finding linked to the historical association of smoking with good outcomes in immunotherapy due to inflammatory tumor microenvironments and higher TMB levels.^{29 30} By 12 and 24 months, these risks reversed, likely reflecting competing risks and evolving dynamics. BMI and dyslipidemia transitioned from marginal roles to significant factors over time, partially aligning with the “obesity paradox” in immune-oncology but suggesting a credible impact in the long term.^{12–15 31 32}

Taken together, this integrative modeling approach enhances clinical interpretability by combining the transparency of traditional regression with the flexibility of deep learning, allowing for a more nuanced understanding of patient trajectories under immunotherapy. Concordant predictors across both models (eg, ECOG-PS, corticosteroid use, and tumor burden) can guide baseline risk stratification and treatment discussions, offering clinicians interpretable, stable

markers for early decision-making. In parallel, time-varying signals captured only by NAIM (eg, BMI, dyslipidemia, or cardiovascular comorbidities) may inform longitudinal patient monitoring, survivorship planning, or tailored follow-up strategies as prognostic relevance shifts over time. Together, these complementary insights foster a layered interpretation of risk, supporting both immediate clinical action and long-term care strategies in a real-world setting.

This study has several limitations. First, it is based on a retrospective, observational dataset, which may carry inherent confounding and selection biases. However, this design reflects the real-world clinical setting in which pembrolizumab is routinely administered to patients with advanced NSCLC and allowed the inclusion of a large, multinational population across 61 centers. To mitigate these biases, we employed rigorous statistical methods, including penalized regression, multivariable adjustment, and an AI-based framework designed to handle incomplete and non-linear data structures. Second, this study did not include a comparator arm treated with chemotherapy or targeted agents. This choice was deliberate: our aim was not to assess the relative efficacy of pembrolizumab but to characterize the time-varying impact of baseline clinical features within a homogeneous, real-world immunotherapy cohort. Including patients treated with alternative therapies would have introduced substantial clinical and molecular heterogeneity (eg, PD-L1-negative tumors, oncogene-addicted subtypes), thus reducing the internal validity of the predictive models. This rationale guided our cohort definition and is a key aspect of our study design.

Missing data remain a pervasive challenge in real-world datasets, arising from various causes such as human error, non-response, or systematic loss. The NAIM model showed moderate predictive performance, with metrics such as c-index of 61.2, an AUC of 60.5–78.5 and low F1-score and MCC values, reflecting a degree of overfitting and limited generalizability. These findings underscore the need for further validation and potential refinement before clinical application. Future studies should validate these findings in external datasets with more complete and diverse representation.

Despite the mentioned limitations, the nuanced, time-dependent insights revealed by the AI-based NAIM model complement the static interpretability of ridge regression, together providing a robust framework for understanding the interplay of oncologic, systemic, and temporal factors. Over the course of a 5-year follow-up, the risk of death in patients with NSCLC treated with immunotherapy evolves to increasingly include non-cancer-related factors. This shift underscores the growing importance of systemic health and holistic care in the survivorship phase. With nearly one-third of patients surviving beyond 5 years in this setting, modern oncology must adopt comprehensive management strategies that address

comorbidities, support metabolic and nutritional health, and incorporate time-sensitive interventions.

Author affiliations

- ¹Operative Research Unit of Medical Oncology, Fondazione Policlinico Universitario Campus Bio-Medico, Rome, Italy
- ²Department of Medicine and Surgery, Università Campus Bio-Medico di Roma, Rome, Italy
- ³Department of Surgery and Cancer, Hammersmith Hospital Campus, Imperial College London, London, UK
- ⁴Lowe Center for Thoracic Oncology, Dana-Farber Cancer Institute, Boston, MA, USA
- ⁵Translational Medicine Department, University of Eastern Piedmont, Novara, Italy
- ⁶Department of Oncology, Beaumont Hospital, Beaumont RCSI Cancer Centre, Dublin, Ireland
- ⁷RCSI University of Health Sciences, Dublin, Ireland
- ⁸Department of Oncology, Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins University, Baltimore, MD, USA
- ⁹Oncology Center, Hospital Sirio-Libanês, São Paulo, SP, Brazil
- ¹⁰Department of Medicine, Division of Oncology, Stanford University, Stanford Cancer Institute, Stanford University, Stanford, CA, USA
- ¹¹Academic Oncology Unit, IRCCS Ospedale Policlinico San Martino, Genova, Italy
- ¹²Section of Medical Oncology, Department of Internal Medicine, Yale Cancer Center, Yale School of Medicine, New Haven, CT, USA
- ¹³Lowe Center for Thoracic Oncology, Dana Farber Cancer Institute, Boston, Massachusetts, USA
- ¹⁴Medical Thoracic Oncology Unit, IRCCS Istituto Tumori "Giovanni Paolo II", Bari, Italy
- ¹⁵Division of Medical Oncology, Department of Internal Medicine, The Ohio State University Comprehensive Cancer Center, Columbus, OH, USA
- ¹⁶Pelotonia Institute for Immuno-Oncology, OSUCCC – James, The Ohio State University, Columbus, OH, USA
- ¹⁷Operative Research Unit of Radiation Oncology, Fondazione Policlinico Universitario Campus Bio-Medico, Rome, Italy
- ¹⁸Niguarda Cancer Center, ASST Grande Ospedale Metropolitano Niguarda, Milano, Italy
- ¹⁹Division of Hematology/Oncology/Stem cell transplant, Rush University Medical Center, Chicago, IL, USA
- ²⁰Medical Oncology, Versilia Hospital, Azienda USL Toscana Nord Ovest, Lido di Camaiore, Italy
- ²¹Department of Oncology, Montefiore Medical Center, Albert Einstein College of Medicine, Bronx, NY, USA
- ²²Medical Oncology, Santa Maria della Misericordia Hospital, Azienda Ospedaliero-Universitaria di Perugia, Perugia, Italy
- ²³Department of Oncology, University of Turin, San Luigi Gonzaga Hospital, Turin, Italy
- ²⁴Department of Thoracic Surgery, Roswell Park Comprehensive Cancer Center, Buffalo, NY, USA
- ²⁵SC Oncologia Medica e Traslationale, Azienda Ospedaliera Santa Maria di Terni, Terni, Italy
- ²⁶Medical Oncology, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy
- ²⁷Department of Oncology, Portsmouth Hospitals University NHS Trust, Portsmouth, Portsmouth, UK
- ²⁸Department of Oncology and National Centre for HIV Malignancy, Chelsea and Westminster Hospital, London, UK
- ²⁹Università Cattolica del Sacro Cuore, Fondazione Policlinico Universitario Agostino Gemelli IRCCS, Rome, Italy
- ³⁰Ospedale Isola Tiberina - Gemelli Isola, Rome, Italy
- ³¹Department of Medicine II, Hematology/Oncology, University Hospital Frankfurt, Frankfurt, Germany
- ³²Translational Oncology and Urology Research, School of Cancer and Pharmaceutical Sciences, King's College London, London, UK
- ³³Clinical Trials Center: Phase 1 and Precision Medicine, IRCCS National Cancer Institute Regina Elena, Rome, Italy
- ³⁴Department of Oncology, Careggi University Hospital, Florence, Italy
- ³⁵Department of Respiratory Medicine, Northumbria Healthcare NHS Foundation Trust, Cramlington, UK
- ³⁶Clinic of Oncology, Cantonal Hospital Fribourg, Faculty of Science and Medicine, University of Fribourg, Fribourg, Switzerland

- ³⁷Department of Oncology and Radiotherapy, Faculty of Medicine, Medical University of Gdańsk, Gdańsk, Poland
- ³⁸Medical Oncology Unit, Sant'Andrea Hospital, Roma, Italy
- ³⁹The UCL Cancer Institute, University College London Hospitals NHS Trust, London, UK
- ⁴⁰UOC Oncologia, AST Pesaro Urbino, Pesaro, Italy
- ⁴¹Thoracic Medical Oncology, Istituto Nazionale Tumori, IRCCS "Fondazione G. Pascale", Naples, Italy
- ⁴²Department of Medicine and Surgery, University of Parma, Parma, Italy
- ⁴³Medical Oncology Unit, University Hospital of Parma, Parma, Italy
- ⁴⁴Clinique Générale Beaulieu, Geneva, Switzerland
- ⁴⁵Oncology Service, University Hospital of Geneva, Geneva, Switzerland
- ⁴⁶Medical Oncology, Ospedale Santa Maria Goretti, Latina, Lazio, Italy
- ⁴⁷Department of Innovative Technologies in Medicine & Dentistry, University G. D'Annunzio, Chieti-Pescara, Chieti, Italy
- ⁴⁸Fortrea Inc, Durham, North Carolina, USA
- ⁴⁹Oncologia Toracica - Lung Unit, Ospedale P. Pederzoli, Peschiera del Garda, Italy
- ⁵⁰4th Department of Medical Oncology and Clinical Trials Unit, Henry Dunant Hospital Center, Athens, Greece
- ⁵¹Department of Pulmonary Oncology, AORN dei Colli Monaldi, Naples, Italy
- ⁵²Medical Oncology Unit, IRCCS Ospedale Sacro Cuore Don Calabria, Negrar di Valpolicella, Italy
- ⁵³Service de Pneumologie, Centre Hospitalier Intercommunal, Creteil, France
- ⁵⁴Hospital Universitario 12 de Octubre, Madrid, Spain
- ⁵⁵Department of Oncology, Azienda Sanitaria Universitaria Friuli Centrale (ASUFC), Udine, Italy
- ⁵⁶Department of Medical Oncology, Hospital de la Santa Creu i Sant Pau, Barcelona, Spain
- ⁵⁷Medical Oncology Unit, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy
- ⁵⁸Division of Medical Oncology, Fondazione IRCCS San Gerardo dei Tintori di Monza, Monza, Italy
- ⁵⁹Department of Clinical Medicine, University of Milano-Bicocca, Milano, Italy
- ⁶⁰Oncology Unit, Santa Maria delle Croci Hospital Ravenna, AUSL della Romagna, Ravenna, Italy
- ⁶¹Medical Oncology Department, Hospital Clinic de Barcelona, Barcelona, Spain
- ⁶²Translational Genomics and Targeted Therapies in Solid Tumors, IDIBAPS, Barcelona, Spain
- ⁶³Department of Medical Oncology, Hospital Clinic, Barcelona, Spain
- ⁶⁴Department of Clinical Medicine and Surgery, University of Naples Federico II, Naples, Italy
- ⁶⁵Medical Oncology Department, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy
- ⁶⁶Department of Electronics, Information, and Bioengineering, Polytechnic University of Milan, Milan, Italy
- ⁶⁷Department of Medical and Surgical Sciences (DIMEC), University of Bologna, Bologna, Italy
- ⁶⁸Medical Oncology Department, Hospital Universitario Infanta Leonor, Madrid, Spain
- ⁶⁹Department of Medical Oncology, Catalan Institute of Oncology (ICO) L'Hospitalet, Barcelona, Spain
- ⁷⁰Division of Pneumology, Cliniques Universitaires St-Luc, UCLouvain, Brussels, Belgium
- ⁷¹Department of Pulmonary Medicine, Erasmus MC Cancer Institute, University Medical Center, Rotterdam, Netherlands
- ⁷²Università Vita-Salute San Raffaele, Milano, Italy
- ⁷³Department of Medical Oncology, IRCCS Ospedale San Raffaele, Milan, Italy
- ⁷⁴Department of Internal Medicine, Medical Oncology, King Hussein Cancer Center, Amman, Jordan
- ⁷⁵Department of Internal Medicine, Hematology, Oncology and Blood and Marrow Transplantation, University of Iowa, Iowa City, IA, USA
- ⁷⁶Department of Surgery, Saiseikai Fukuoka General Hospital, Fukuoka, Japan
- ⁷⁷Medical Oncology/TSET Phase 1 Program, The University of Oklahoma Stephenson Cancer Center, Oklahoma City, Oklahoma, USA
- ⁷⁸Thoracic Oncology Program, Section of Hematology/Oncology, Department of Medicine, University of Chicago, Chicago, Illinois, USA
- ⁷⁹Centre Hospitalier Universitaire Vaudois, Lausanne University, Lausanne, Switzerland
- ⁸⁰Research Unit of Computer Systems and Bioinformatics, Department of Engineering, Università Campus Bio-Medico di Roma, Rome, Italy

⁸¹Department of Diagnostics and Intervention, Radiation Physics, Biomedical Engineering, Umeå University, Umeå, Sweden

Social media Alessio Cortellini, X @ACortelliniMD

Acknowledgements ACo acknowledges the support of the International Association for the Study of Lung Cancer (IASLC) young investigators grant 2024. EB acknowledges the support by the Associazione Italiana per la Ricerca sul Cancro (AIRC, Investigator Grant No. IG20583), by Institutional funds of Università Cattolica del Sacro Cuore (UCSC-project D1), and funds of Ministero della Salute (Ricerca Corrente 2024). CMC is a Ph.D. student enrolled in the National Ph.D. in Artificial Intelligence, XXXVII cycle, course on Health and Life Sciences, organized by Università Campus Bio-Medico di Roma. PS, CMC, and VG acknowledge the funding from: PRIN 2022 MUR 20228MZFAA-AIDA (CUP C53D23003620008); PNRR MUR project PE0000013-FAIR. PS acknowledges the support by the National Academic Infrastructure for Supercomputing in Sweden (NAISS) and the Swedish National Infrastructure for Computing (SNIC) at Alvis @ C3SE, partially funded by the Swedish Research Council through grant agreements no. 2022-06725 and no. 2018-05973.

Contributors All authors contributed to the publication according to the ICMJE guidelines for the authorship. All authors read and approved the submitted version of the manuscript (and any substantially modified version that involves the author's contribution to the study). Each author has agreed both to be personally accountable for the author's own contributions and to ensure that questions related to the accuracy or integrity of any part of the work, even ones in which the author was not personally involved, are appropriately investigated, resolved, and the resolution documented in the literature. Study concept and design: ACo, BR, VG, CMC. Acquisition of data: ACo, VSa, GLC, DR, LB, EG, DJP, JN, AK, MLo, JWN, CG, SGe, SYK, RJ, TEZ, MRu, FPe, ADF, JVA, MMo, DHO, DS, MJF, MLI, Aca, SR, BH, BV, GMe, FPa, SY, AGu, MG, AD'Alessio, GLB, CAMF, SGr, FG, AD'Incecco, EJ, MVH, AR, AGe, GS, MV, BT, RG, KB, EB, MS, MRo, MF, UM, LL, FM, AAu, MD, AC-F, RC, Vsf, MT, AF, AAd, FZ, MDT, LC, ER, Gmo, LDG, SK, AI, PB, RDMB, DO'R, OF, EK, IM, JB, MMA, API, FA, DLC, GT, GMi, CB, LMe, TG, AS, TB, GLR, AP, ADG, JR, LMo, EN, FB, FAN, A-MD, JGJVA, RF, TAH, KT, ARN, MCG, SP, HW, AHN, BR, PS, CMC and VG. Analysis and interpretation of data: ACo, BR, VG, CMC. Statistical analysis: ACo, BR, VG and CMC. Manuscript review and approval: ACo, VSa, GLC, DR, LB, EG, DJP, JN, AK, MLo, JWN, CG, SGe, SYK, RJ, TEZ, MRu, FPe, ADF, JVA, MMo, DHO, DS, MJF, MLI, ACo, SR, BH, BV, GMe, FPa, SY, AGu, MG, AD'Alessio, GLB, CAMF, SGr, FG, AD'Incecco, EJ, MVH, AR, AGe, GS, MV, BT, RG, KB, EB, MS, MRo, MF, UM, LL, FM, AAu, MD, AC-F, RC, Vsf, MT, AF, AAd, FZ, MDT, LC, ER, Gmo, LDG, SK, AI, PB, RDMB, DO'R, OF, EK, IM, JB, MMA, API, FA, DLC, GT, GMi, CB, LMe, TG, AS, TB, GLR, AP, ADG, JR, LMo, EN, FB, FAN, A-MD, JGJVA, RF, TAH, KT, ARN, MCG, SP, HW, AHN, BR, PS, CMC and VG. Study supervision: ACo, BR, VG and CMC. Artificial intelligence (AI) was used in this study to perform data analysis. Specifically, AI-based tools and algorithms were employed for data processing, statistical modeling, and/or predictive analysis. All outputs were reviewed and validated by the authors to ensure accuracy and scientific integrity. CMC and VG are joint last authors. Guarantor: ACo.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests ACo received grants for consultancies/advisory boards from MSD, BMS, IQVIA, AstraZeneca, REGENERON, Amgen, Daiichi-Sankyo, Access Infinity, Ardelis Health, Alpha Sight, Guidepoint, Roche; speaker fees from AstraZeneca, Pierre-Fabre, MSD, Sanofi/REGENERON; payment for writing/editorial activity from BMS, MSD, Roche; travel support from Sanofi/REGENERON, MSD. JB declares honoraria/consulting or advisory role from AstraZeneca, BMS, Roche, Access Oncology, travel support from MSD, Roche, Janssen Oncology. GS has received payment or honoraria for advisory boards from Novartis, Roche, Bayer, unrelated to this project. DO'R has received conference attendance support from Takeda, Janssen, Servier, MSD. EB has received grants or contracts from AstraZeneca, Roche and honoraria for lectures from Merck-Sharp & Dome, Astra-Zeneca, Pfizer, Eli-Lilly, Bristol-Myers Squibb, Novartis, Takeda and Roche; EB has been member of Data Safety Monitoring Board or Advisory Board of Merck-Sharp & Dome, Pfizer, Novartis, Bristol-Myers Squibb, Astra-Zeneca, Celltrion and Roche. AAu declares consulting or advisory role for Bristol Myers Squibb, AstraZeneca, Boehringer Ingelheim, Roche, MSD, Pfizer, Eli Lilly, Astellas, Takeda, and Amgen; speaker's bureau for Eli Lilly, and AstraZeneca. AR has received advisory board or speaker bureau honoraria from AstraZeneca, MSD, Novartis, Pfizer, BMS, Takeda, Amgen, Regeneron and Daiichi Sankyo; compensated activity for editorial projects from AstraZeneca, MSD, BMS, Novartis, Roche, and Regeneron. MT received speakers' and consultants' fee from Astra-Zeneca, Pfizer, Eli-Lilly, BMS, Novartis, Roche, MSD, Boehringer Ingelheim, Takeda, Amgen, Merck, Sanofi, Janssen,

Daiichi Sankyo. He also received institutional research grants from Astra-Zeneca, Boehringer Ingelheim and Roche and travel support from Amgen and Takeda. FM received honoraria for advisory board roles with MDS, BMS, Takeda, Roche, Astra-Zeneca, Novartis. PB served as consultant/advisory board for Regeneron, Pierre-Fabre, Janssen, Seagen. DHO declares research funding/grants (to institution) from BMS, Merck, Palobiofarma, Pfizer, Genentech, AstraZeneca, Nuvalent, Abbvie, Onc. AI. BH received grants for consultancies from Boehringer Ingelheim, Astra Zeneca, Merck, BMS, Advaxis, Amgen, AbbVie, Daiichi, Pfizer, GSK, Beigene, Janssen, Black Diamond Therapeutics, Forward Pharma, Numab, Arrivent; speakers fees from Astra Zeneca, Boehringer Ingelheim, Apollomics, Johnson&Johnson, Takeda, Merck, BMS, Genentech, Pfizer, Eli-Lilly, Daiichi; grants for participating on boards from BMS, TPT, Apollomics, eFECTOR, and City of Hope. BT received lecture fees from Pfizer. LC is an employee of Fortrea Inc. BT declares honoraria from Roche. IM declares travel support from Takada, MSD, Pfizer, Oxyvie and speaker fees from Regeneron. A-MD declares research grants from Amgen, the Dutch Cancer Society and HANART, consulting fees from Amgen, Bayer, Boehringer Ingelheim, Sanofi, Roche, Janssen and Astrazeneca, speaker fees from Janssen, Pfizer, Astrazeneca, Lilly and Takeda, advisory board role for Takeda and Roche. GLR declares fees for advisory boards, travel support, consultancies from MSD, BMS, Roche, Sanofi, Regeneron, Lilly, Astrazeneca, Janssen, Pfizer, Novartis, Bayer, Takeda, Amgen, GSK, Daiichi. TAH declares stock interests for GlaxoSmithKline and honoraria from Novartis. BR served as consultant/advisory board for AMGEN, Regeneron, AstraZeneca, Capvision. Speaker fee: AstraZeneca. Received honoraria from Targeted Oncology, SITC. All other authors declare no conflicts of interest associated with the present study.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants. The procedures followed were in accordance with the precepts of Good Clinical Practice and the Declaration of Helsinki. The Pembro-real IT cohort Institutional Review Board (IRB) approval reference is "Comitato Etico per le province di L'Aquila e Teramo, verbale N.15 del 28 Novembre 2019)". For Italian institutions participating to the Pembro-real 5-year cohort the IRB reference is "Comitato Etico Fondazione Policlinico Universitario Campus Bio-Medico, IRB ID approval N.PAR 70.23 OSS, May 17, 2023, registry number: SC 2023.0682" (written informed consent was obtained for patients alive at the time of data collection). For the non-Italian institutions participating to the Pembro-real -year cohort the IRB reference is "Health Research Authority approval of the 22nd of November 2023, REC reference 23/HRA/4467" (written informed consent was waived due to the retrospective and observational nature of the study). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available. The dataset used for this study contains patient-level data that cannot be made available to third parties, although anonymized. Third-party research proposals will be assessed by the study investigators and performed by the study team if accepted. Requests can be made to ACo (a.cortellini@policlinicocampus.it).

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Alessio Cortellini <http://orcid.org/0000-0002-1209-5735>
David J Pinato <http://orcid.org/0000-0002-3529-0103>
So Yeon Kim <http://orcid.org/0000-0003-4102-5880>
Joao V Alessi <http://orcid.org/0000-0002-8072-5946>
Michele Montrone <http://orcid.org/0000-0002-8568-7723>
Dwight H Owen <http://orcid.org/0000-0002-6598-4886>
Mingjia Li <http://orcid.org/0000-0002-3910-4179>
Balazs Halmos <http://orcid.org/0000-0001-7548-8360>
Sai Yendamuri <http://orcid.org/0000-0001-6654-3487>

Antonio D'Alessio <http://orcid.org/0000-0002-9164-3671>
Raffaele Giusti <http://orcid.org/0000-0003-4592-8868>
Emilio Bria <http://orcid.org/0000-0002-2333-704X>
Marcello Tiseo <http://orcid.org/0000-0002-9553-8465>
Giannis Mountzios <http://orcid.org/0000-0002-7780-7836>
Diego Luigi Cortinovis <http://orcid.org/0000-0001-7611-7369>
Laura Mezquita <http://orcid.org/0000-0003-0936-7338>
Arsela Prelaj <http://orcid.org/0000-0002-3863-088X>
Andrea De Giglio <http://orcid.org/0000-0002-0731-5721>
Ernest Nadal <http://orcid.org/0000-0002-9674-5554>
Joachim G J V Aerts <http://orcid.org/0000-0001-6662-2951>
Marina Chiara Garassino <http://orcid.org/0000-0002-3821-5598>
Amin H Nassar <http://orcid.org/0000-0002-4507-2396>
Biagio Ricciuti <http://orcid.org/0000-0002-0651-2678>

REFERENCES

- 1 Remon J, Hendriks LEL, Besse B. Paving the Way for Long-Term Survival in Non-Small-Cell Lung Cancer. *J Clin Oncol* 2021;39:2321-3.
- 2 de Jager VD, Timens W, Bayle A, et al. Developments in predictive biomarker testing and targeted therapy in advanced stage non-small cell lung cancer and their application across European countries. *Lancet Reg Health Eur* 2024;38:100838.
- 3 Garassino MC, Gadgeel S, Speranza G, et al. Pembrolizumab Plus Pemetrexed and Platinum in Nonsquamous Non-Small-Cell Lung Cancer: 5-Year Outcomes From the Phase 3 KEYNOTE-189 Study. *J Clin Oncol* 2023;41:1992-8.
- 4 Novello S, Kowalski DM, Luft A, et al. Pembrolizumab Plus Chemotherapy in Squamous Non-Small-Cell Lung Cancer: 5-Year Update of the Phase III KEYNOTE-407 Study. *J Clin Oncol* 2023;41:1999-2006.
- 5 Reck M, Ciuleanu T-E, Schenker M, et al. Five-year outcomes with first-line nivolumab plus ipilimumab with 2 cycles of chemotherapy versus 4 cycles of chemotherapy alone in patients with metastatic non-small cell lung cancer in the randomized CheckMate 9LA trial. *Eur J Cancer* 2024;211:114296.
- 6 Reck M, Rodríguez-Abreu D, Robinson AG, et al. Five-Year Outcomes With Pembrolizumab Versus Chemotherapy for Metastatic Non-Small-Cell Lung Cancer With PD-L1 Tumor Proportion Score $\geq$ 50. *J Clin Oncol* 2021;39:2339-49.
- 7 Kilickap S, Sezer A, Özgüröglu M, et al. OA11.06 Cemiplimab Monotherapy for First Line Advanced NSCLC Patients with PD-L1 Expression $\geq$ 50%: 5-y Outcomes of EMPOWER-Lung 1. *J Thorac Oncol* 2024;19:S35.
- 8 Goldstraw P, Chansky K, Crowley J, et al. The IASLC Lung Cancer Staging Project: Proposals for Revision of the TNM Stage Groupings in the Forthcoming (Eighth) Edition of the TNM Classification for Lung Cancer. *J Thorac Oncol* 2016;11:39-51.
- 9 Gridelli C, Hainsworth J. Meeting the chemotherapy needs of elderly and poor performance status patients with NSCLC. *Lung Cancer (Auckl)* 2002;38:37-41.
- 10 Cortellini A, D'Alessio A, Cleary S, et al. Type 2 Diabetes Mellitus and Efficacy Outcomes from Immune Checkpoint Blockade in Patients with Cancer. *Clin Cancer Res* 2023;29:2714-24.
- 11 Mallardo D, Woodford R, Menzies AM, et al. The role of diabetes in metastatic melanoma patients treated with nivolumab plus relatlimab. *J Transl Med* 2023;21:753.
- 12 Perrone F, Favari E, Maglietta G, et al. The role of blood cholesterol quality in patients with advanced cancer receiving immune checkpoint inhibitors. *Cancer Immunol Immunother* 2023;72:2127-35.
- 13 Pecci F, Cantini L, Cognigni V, et al. Prognostic Impact of Blood Lipid Profile in Patients With Advanced Solid Tumors Treated With Immune Checkpoint Inhibitors: A Multicenter Cohort Study. *Oncologist* 2024;29:e372-81.
- 14 Cortellini A, Ricciuti B, Tiseo M, et al. Baseline BMI and BMI variation during first line pembrolizumab in NSCLC patients with a PD-L1 expression $\geq$ 50%: a multicenter study with external validation. *J Immunother Cancer* 2020;8:e001403.
- 15 Cortellini A, Bersanelli M, Buti S, et al. A multicenter study of body mass index in cancer patients treated with anti-PD-1/PD-L1 immune checkpoint inhibitors: when overweight becomes favorable. *J Immunother Cancer* 2019;7:57.
- 16 Cortellini A, Bozzetti F, Palumbo P, et al. Weighing the role of skeletal muscle mass and muscle density in cancer patients receiving PD-1/PD-L1 checkpoint inhibitors: a multicenter real-life study. *Sci Rep* 2020;10:1456.

- 17 Cortellini A, Verna L, Porzio G, *et al.* Predictive value of skeletal muscle mass for immunotherapy with nivolumab in non-small cell lung cancer patients: A “hypothesis-generator” preliminary report. *Thorac Cancer* 2019;10:347–51.
- 18 Cortellini A, Ricciuti B, Borghaei H, *et al.* Differential prognostic effect of systemic inflammation in patients with non-small cell lung cancer treated with immunotherapy or chemotherapy: A post hoc analysis of the phase 3 OAK trial. *Cancer* 2022;128:3067–79.
- 19 Banna GL, Cortellini A, Cortinovis DL, *et al.* The lung immunology prognostic score (LIPS-3): a prognostic classification of patients receiving first-line pembrolizumab for PD-L1 $\geq$ 50% advanced non-small-cell lung cancer. *ESMO Open* 2021;6:100078.
- 20 Cortellini A, Brunetti L, Di Fazio GR, *et al.* Determinants of 5-year survival in patients with advanced NSCLC with PD-L1 $\geq$ 50% treated with first-line pembrolizumab outside of clinical trials: results from the Pembro-real 5Y global registry. *J Immunother Cancer* 2025;13:e010674.
- 21 Caruso CM, Guarrasi V, Ramella S, *et al.* A deep learning approach for overall survival prediction in lung cancer with missing values. *Comput Methods Programs Biomed* 2024;254:108308.
- 22 Caruso CM, Soda P, Guarrasi V, *et al.* Not Another Imputation Method: A Transformer-based Model for Missing Values in Tabular Datasets. *arXiv* 2014.
- 23 Caruso CM, Soda P, Guarrasi V, *et al.* MARIA: a Multimodal Transformer Model for Incomplete Healthcare Data. *arXiv* 2024.
- 24 Lundberg SM, Lee S-I. A unified approach to interpreting model predictions. Proceedings of the 31st International Conference on Neural Information Processing Systems; Long Beach, California, USA, 2017
- 25 Ogliari FR, Traverso A, Barbieri S, *et al.* Exploring machine learning tools in a retrospective case-study of patients with metastatic non-small cell lung cancer treated with first-line immunotherapy: A feasibility single-centre experience. *Lung Cancer (Auckl)* 2025;199:108075.
- 26 Koyama J, Morise M, Furukawa T, *et al.* Artificial intelligence-based personalized survival prediction using clinical and radiomics features in patients with advanced non-small cell lung cancer. *BMC Cancer* 2024;24:1417.
- 27 Kumar R, Castillero F, Bhandari S, *et al.* The Hispanic Paradox in Non-Small Cell Lung Cancer. *Hematol Oncol Stem Cell Ther* 2022;15:21–9.
- 28 Price SN, Flores M, Hamann HA, *et al.* Ethnic Differences in Survival Among Lung Cancer Patients: A Systematic Review. *JNCI Cancer Spectr* 2021;5:pkab062.
- 29 Gainor JF, Rizvi H, Jimenez Aguilar E, *et al.* Clinical activity of programmed cell death 1 (PD-1) blockade in never, light, and heavy smokers with non-small-cell lung cancer and PD-L1 expression $\geq$ 50. *Ann Oncol* 2020;31:404–11.
- 30 Cortellini A, De Giglio A, Cannita K, *et al.* Smoking status during first-line immunotherapy and chemotherapy in NSCLC patients: A case-control matched analysis from a large multicenter study. *Thorac Cancer* 2021;12:880–9.
- 31 Bersanelli M, Cortellini A, Buti S. The interplay between cholesterol (and other metabolic conditions) and immune-checkpoint immunotherapy: shifting the concept from the “inflamed tumor” to the “inflamed patient”. *Hum Vaccin Immunother* 2021;17:1930–4.
- 32 Cortellini A, Bersanelli M, Santini D, *et al.* Another side of the association between body mass index (BMI) and clinical outcomes of cancer patients receiving programmed cell death protein-1 (PD-1)/Programmed cell death-ligand 1 (PD-L1) checkpoint inhibitors: A multicentre analysis of immune-related adverse events. *Eur J Cancer* 2020;128:17–26.