



# Ursodeoxycholate inhibits mast cell activation and reverses biliary injury and fibrosis in $Mdr2^{-/-}$ mice and human primary sclerosing cholangitis

Fanyin Meng<sup>1,2,3</sup> · Lindsey Kennedy<sup>1</sup> · Laura Hargrove<sup>3</sup> · Jennifer Demieville<sup>1</sup> · Hannah Jones<sup>2</sup> · Taronish Madeka<sup>2</sup> · Allen Karstens<sup>2</sup> · Kevin Chappell<sup>2</sup> · Gianfranco Alpini<sup>1,2,3</sup> · Amelia Sybenga<sup>2</sup> · Pietro Invernizzi<sup>4,5</sup> · Francesca Bernuzzi<sup>4</sup> · Sharon DeMorrow<sup>1,3</sup> · Heather Francis<sup>1,2,3</sup>

Received: 6 August 2017 / Revised: 10 May 2018 / Accepted: 12 May 2018 / Published online: 24 August 2018  
© United States & Canadian Academy of Pathology 2018

## Abstract

Ursodeoxycholic acid (UDCA) is used to treat biliary disorders; and, bile acids alter mast cell (MC) histamine release. MCs infiltrate  $Mdr2^{-/-}$  mice liver (model of primary sclerosing cholangitis (PSC)). MC-derived histamine increases inflammation, hepatic stellate cell (HSC) activation and fibrosis. The objective was to determine the effects of UDCA treatment on MC infiltration, biliary damage, inflammation and fibrosis in  $Mdr2^{-/-}$  mice and human PSC. Wild-type and  $Mdr2^{-/-}$  mice were fed bile acid control diet or UDCA (0.5% wt/wt). Human samples were collected from control and PSC patients treated with placebo or UDCA (15 mg/kg/BW). MC infiltration was measured by immunohistochemistry and quantitative polymerase chain reaction (qPCR) for c-Kit, chymase, and tryptase. The HDC/histamine/histamine receptor (HR)-axis was evaluated by EIA and qPCR. Intrahepatic bile duct mass (IBDM) and biliary proliferation was evaluated by CK-19 and Ki-67 staining. Fibrosis was detected by immunostaining and qPCR for fibrotic markers. Inflammatory components were measured by qPCR. HSC activation was measured by SYP-9 staining. Inflammation was detected by qPCR for CD68. In vitro, MCs were treated with UDCA (40  $\mu$ M) prior to HA secretion evaluation and coculturing with cholangiocytes or HSCs. BrDU incorporation and fibrosis by qPCR was performed. UDCA reduced MC number, the HDC/histamine/HR-axis, IBDM, HSC activation, inflammation, and fibrosis in  $Mdr2^{-/-}$  mice and PSC patients. In vitro, UDCA decreases MC-histamine release, which was restored by blocking ASBT and FXR $\beta$ . Proliferation and fibrosis decreased after treatment with UDCA-treated MCs. We conclude that UDCA acts on MCs reducing histamine levels and decreases the inflammatory/hyperplastic/fibrotic reaction seen in PSC. Ursodeoxycholic acid (UDCA) is used to treat biliary disorders; and, bile acids alter mast cell (MC) histamine release. Following liver injury like primary sclerosing cholangitis in mice and humans, MCs infiltrate. MC-derived histamine increases biliary damage, fibrosis, and inflammation. UDCA treatment decreases these parameters via reduced MC activation.

**Electronic supplementary material** The online version of this article (<https://doi.org/10.1038/s41374-018-0101-0>) contains supplementary material, which is available to authorized users.

✉ Heather Francis  
hfrancis@medicine.tamhsc.edu  
Heather.Francis@va.gov

<sup>1</sup> Research, Central Texas Veterans Health Care System, Temple, TX, USA

<sup>2</sup> Scott & White Digestive Disease Research Center, Baylor Scott & White Health, Temple, TX, USA

## Introduction

Primary sclerosing cholangitis (PSC) is a chronic cholestatic disease that damages bile ducts inside and outside the liver. During PSC, bile ducts are inflamed, which leads to scarring and narrowing resulting in fibrosis and cirrhosis of

<sup>3</sup> Medical Physiology, Texas A&M Health Science Center, Temple, TX, USA

<sup>4</sup> Humanitas Clinical and Research Center, Rozzano, Milan, Italy

<sup>5</sup> Present address: Program for Autoimmune Liver Diseases, International Center for Digestive Health, Department of Medicine and Surgery, University of Milan-Bicocca, Monza, Italy

the liver and, eventually, liver failure [1]. The only definitive treatment to date is liver transplantation [1, 2]. Multi-drug resistance-2 gene knockout mice ( $Mdr2^{-/-}$ ) develop spontaneous liver fibrosis and mimic some features of human PSC [3, 4].  $Mdr2^{-/-}$  mice have enhanced intrahepatic bile duct mass (IBDM), biliary proliferation and hepatic fibrosis between 4 and 12 weeks of age [4, 5].

Within MCs are intact granules that carry inflammatory mediators like histamine and various cytokines and chemokines [6–8]. MCs infiltrate the liver upon damage, surround bile ducts increasing local histamine levels and enhance biliary mass, proliferation, and hepatic fibrosis [5, 9]. In our previous work, we demonstrated that, in bile duct-ligated (BDL) rats and  $Mdr2^{-/-}$  mice treated with the MC stabilizer, cromolyn sodium, there is reduced biliary damage and liver fibrosis [5, 9]. In MC-deficient mice, IBDM and hepatic fibrosis are reduced following BDL compared to WT BDL mice [10]. In addition, when MC-deficient mice were injected with cultured MCs, these parameters increased demonstrating that MCs regulate biliary injury and hepatic fibrosis [10].

Ursodeoxycholic acid (UDCA) is a natural bile acid that forms about 1–3% of total bile acids in human bile [11]. To date, UDCA (13–15 mg/kg/day) is the choice treatment for primary biliary cholangitis (PBC) patients that have abnormal liver enzymes; however, the role of UDCA in PSC has remained elusive and conflicting [11]. In fact, at high doses (28–30 mg/kg/day) UDCA can be harmful to PSC patients and studies to demonstrate effectiveness at lower doses are inconclusive [1]. UDCA exerts its anticholestatic effects by lowering serum liver enzymes, improving histology and increasing nontoxic bile acid pools, thus delaying the progression to liver transplant [2, 12, 13]. UDCA has been shown to regulate inflammatory cytokine release during cholestatic liver injury [14]. According to the 2009 European Association for the Study of Liver Disease (EASL), UDCA is an approved therapy for PSC patients and numerous patients are prescribed UDCA to ameliorate uncomfortable side effects [15]. The purpose of our study is to understand if UDCA acts on MCs or induces a synergistic interaction between MCs, cholangiocytes and hepatic stellate cells. Since it is unknown if UDCA interacts with MCs to alter PSC progression, we aimed to study these parameters in both  $Mdr2^{-/-}$  mice and human PSC.

## Method and materials

### Reagents and other materials

Chemical grade reagents were purchased from Sigma-Aldrich Co (St. Louis, MO) unless stated otherwise. All mouse and human primers and real-time PCR materials

were obtained from Qiagen (Fredrick, MD). Antibodies for immunohistochemistry and immunofluorescence were purchased from Abcam (Cambridge, MA) or Santa Cruz (Dallas, TX). Histamine EIA kits were purchased from Cayman Chemical (An Arbor, MI) [5, 9, 10]. The apical sodium-bile acid transporter inhibitor was obtained from Tocris Bioscience (Minneapolis, MN) and the farnesoid X receptor (FXR $\beta$ ) inhibitor was obtained from Sigma-Aldrich Co (St. Louis, MO). Kits to measure total bile acids (TBAs) were purchased from Crystal Chemical, Inc (Downers Grove, IL) [5].

### In vivo models

We used the genetically modified mouse model of PSC,  $Mdr2^{-/-}$  mice on FVB/NJ background (wild-type, WT) [3, 5]. All mice were housed in the Baylor Scott and White Health Animal Facility and given free access to drinking water and standard chow. All animals were kept in a temperature-controlled environment with a 12:12 light/dark cycle, and all protocols strictly adhered to regulations as set forth by the local IACUC committee. Male mice (9–11 weeks of age) were fed a bile acid control diet (BAC, which contains the same components as the UDCA chow minus the bile acids)—or UDCA—enriched diet (0.5% wt/wt) [16] for 1 week (10–12 mice per group). Diets were obtained from Dyets, Inc. (Bethlehem, PA) and from all animals we collected serum, isolated cholangiocytes (as previously described [5, 10, 17]) and supernatants (collected after 4–6 h of incubation at 37° [5, 9, 10]) and liver blocks (frozen and paraffin-embedded). We performed hematoxylin and eosin (H&E) staining in livers to evaluate lobular damage, hepatic necrosis and inflammation. TBAs were measured in serum and/or snap liver as previously described [5].

### Human PSC sampling

Human liver sections from normal controls (collected as part of liver cholangiocarcinoma resection and considered non-diseased areas) and PSC patients treated with placebo or UDCA (15 mg/kg/BW) were utilized (Supplementary Table 1). PSC patients presented with a range of early to late PSC as described in our previous work [5]. Liver sections (4–5  $\mu$ m thick) obtained by needle biopsies from three control patients, three early stage PSC (with or without cirrhosis) and three early stage PSC patients treated with UDCA (with or without cirrhosis). The deidentified samples were provided by Dr. Pietro Invernizzi under a protocol approved by Humanitas Research Hospital; the protocol was approved by the Central Texas Veteran's Health Care System Institutional Review Board and Research Development Committee. Using acquired human tissue, H&E was performed to assess liver

damage that was evaluated by a blinded pathologist; patient serum ALT levels were assessed via colorimetric analysis with an AbCam ALT kit (ab105134).

### Detection of MCs, MC markers, HR, and histamine secretion

In all mice, MC presence was detected by immunohistochemistry in liver sections (4–5  $\mu\text{m}$  thick) for mouse MC protease-1 (mMCP-1) and MC markers were measured in total liver by real-time PCR [9, 18]. The HDC/histamine receptor (HR)/HA-axis was measured by real-time PCR and by EIA from all groups of mice [17]. We measured MC number by toluidine blue staining in liver from normal rats fed either taurocholate (TC) or tauroolithocholate (TLC) [19, 20], and in BDL rats fed UDCA [21].

We measured the expression of MC presence by immunohistochemistry for tryptase in human liver sections from control and PSC patients treated with placebo or UDCA. The gene expression of MC markers was also measured in these livers. H1–H4HR expression was measured by real-time PCR and histamine secretion was evaluated by EIA in serum from all patient groups.

### Evaluation of biliary damage in $\text{Mdr2}^{-/-}$ mice and hepatic fibrosis/HSC activation in mice and human PSC

In liver sections from all mice, we performed immunohistochemistry for the biliary marker, CK-19 [9, 17] to measure IBDM and Ki-67 [5] to detect the percentage of proliferating cholangiocytes as previously described by us. We measured fibrosis and HSC activation from all mice and human tissues as previously described by us [5, 9, 10].

Fibrosis was assessed by Masson's Trichrome and Fast Green/Sirius Red (along with semiquantification) [17]. In total liver, we measured the following fibrotic markers by real-time PCR: collagen type-1a and fibronectin-1 [17]. TGF- $\beta$ 1 is a known regulator of HSC-driven fibrosis [5, 13]; therefore, we measured TGF- $\beta$ 1 expression by real-time PCR in all groups of mice. To evaluate HSC activation, synaptophysin 9 (SYP-9) was measured by immunofluorescence in all mice groups co-stained with CK-19 to detect bile ducts [5, 10]. In human tissues, we measured the gene expression of fibronectin-1 [22, 23] and performed Masson's trichrome.

### Measurement of the effects of UDCA treatment on inflammation and extracellular matrix proteins

We measured *CD68* in our mouse tissues as an indicator of inflammatory response by real-time PCR. TNF- $\alpha$  is released

from degranulated MC [18]; therefore, we evaluated the expression of TNF- $\alpha$  in total liver from all mice by real-time PCR. We measured MMP-2, -3, and -9 by real-time PCR in total liver from WT,  $\text{Mdr2}^{-/-}$  + BAC and  $\text{Mdr2}^{-/-}$  + UDCA.

### In vitro evaluation of MC/cholangiocyte/HSC interactions

To determine the direct effects of UDCA on (i) MC-derived histamine release and (ii) biliary and HSC response, we utilized cultured MCs, cholangiocytes, and human HSCs (hHSCs).

Cultured MCs were obtained from ATCC (Manassas, VA) and human HSCs from ScienCell, Carlsbad, CA. Murine cholangiocytes have previously been used by us and all cells were cultured according to previous protocols [5, 9, 10, 17].

To determine how bile acids interact with MCs, we measured the expression of ASBT and FXR $\beta$  by immunofluorescence in cultured MCs. The ASBT antibody was a kind gift from Dr. Paul Dawson (Emory University School of Medicine) and FXR $\beta$  antibodies were purchased from Santa Cruz Biotechnology (Dallas, TX).


To demonstrate that bile acid treatment alters MC-histamine release, MCs were treated with UDCA (40  $\mu\text{M}$ , 30–120 min) in the absence or presence of specific antagonists, guggulsterone (FXR $\beta$  inhibitor, 10  $\mu\text{M}$ ) and SC-435 (ASBT inhibitor, 100 nM) and histamine release was measured by EIA [10]. In separate experiments, MCs were treated with different bile acids including TC (40  $\mu\text{M}$ , 24 h) and TLC (40  $\mu\text{M}$ , 24 h) as well as UDCA (40  $\mu\text{M}$ , 120 min); MC activation, MMP-2, -3, and -9 and TNF- $\alpha$  levels were evaluated by real-time PCR.

Next, cultured cholangiocytes or hHSCs were treated with 0.1% BSA (basal), MC supernatants (basal-treated) and MC supernatants collected following treatment with UDCA (40  $\mu\text{M}$ , 120 min). Biliary proliferation was evaluated by BrDU staining and fibronectin-1 was measured by real-time PCR in treated HSCs [5, 10]. Biliary inflammation was measured by real-time PCR in treated cells for TNF- $\alpha$ , CCL-3, and IL-6.

In our final experiments, hHSCs were treated with 0.1% BSA (basal) or cholangiocyte supernatants collected from WT,  $\text{Mdr2}^{-/-}$  + BAC and  $\text{Mdr2}^{-/-}$  mice treated with UDCA. Following stimulation for 24 h, the expression of fibrosis markers including, TGF- $\beta$ 1, collagen type-1a and fibronectin-1 were evaluated by real-time PCR.

### Statistical methods

All data is expressed as mean  $\pm$  SEM. Groups were analyzed by the Student unpaired *t* test when two groups are



**Fig. 1** **a** Mdr2<sup>-/-</sup> mice fed BAC displayed increased portal area damage including inflammation, necrosis, and increased ductular reaction compared to WT mice. **b** Total bile acids (TBAs) were increased in serum in Mdr2<sup>-/-</sup> mice fed BAC compared to WT and feeding with UDCA decreased TBAs in serum. **c** H&E in human biopsies reveal increased inflammation and necrosis in PSC patients

compared to control, which are reduced in PSC patients treated with UDCA. **d** Serum ALT levels are increased in patients with PSC compared to control and ALT levels are decreased in two patients treated with UDCA. Data are expressed as mean ± SEM of at least eight experiments for TBA. \**p* < 0.05 vs. WT mice; #*p* < 0.05 vs. Mdr2<sup>-/-</sup> mice fed BAC. Images are 20× magnification

analyzed or a two-way ANOVA when more than two groups are analyzed, followed by an appropriate post hoc test. *p* < 0.05 was considered significant.

## Results


### UDCA treatment ameliorates liver injury in Mdr2<sup>-/-</sup> mice and human PSC

Extensive inflammation was found around bile ducts, which was ameliorated in Mdr2<sup>-/-</sup> mice fed UDCA, which were similar to WT (Fig. 1a). In Mdr2<sup>-/-</sup> mice fed UDCA, total bile acid serum content decreased (Fig. 1b). According to pathology reports, the WT mouse liver has no significant histopathological findings; however, the Mdr2<sup>-/-</sup> mouse fed BAC displays moderate to severe cholangitis with prominent duct obliteration and bridging fibrosis that is consistent with human PSC. Some bridging fibrosis and inactive hepatitis was also noted in the Mdr2<sup>-/-</sup> mouse fed BAC. Finally, in the Mdr2<sup>-/-</sup> mouse fed UDCA mild portal

inflammation was detected along with minimal bridging fibrosis and mild cholangitis.

In human PSC, focal necrosis and inflammation increased compared to controls, which were decreased in PSC patients treated with UDCA (Fig. 1c). Further, serum ALT levels increased in PSC patients compared to controls and several patients treated with UDCA had decreased ALT levels, although not all patients responded to UDCA (Fig. 1d). For H&E staining, the pathology reports indicate that there is mild steatohepatitis with limited fibrosis in the human control sample, whereas the PSC liver displays bile duct sclerosing cholangitis with bridging fibrosis and potential bile duct obstruction. The PSC patient treated with UDCA also displays mild sclerosing cholangitis; however, no evidence of obstruction was noted.

To correlate our findings with human dosing, we estimate that mice consumed a relative dose of 20–23 mg/kg BW per day. This calculation was based on the amount of UDCA in the diet, average weight of the mice, and average chow consumed per day. As stated, normal/low dose treatment is indicated at 13–15 mg/kg BW per day and



**Fig. 2 a** Mast cell numbers are increased in *Mdr2*<sup>-/-</sup> mice fed BAC compared to WT and UDCA feeding decreases the presence of mast cells shown by immunohistochemistry for mMCP-1 (marked by red arrows). Semiquantification demonstrates a significant increase in mast cell number in *Mdr2*<sup>-/-</sup> mice fed BAC that is reduced in *Mdr2*<sup>-/-</sup> mice fed UDCA. **b** By immunohistochemistry for tryptase there is an

infiltration of mast cells found surrounding bile ducts in PSC patients treated with placebo compared to normal tissue that were void of mast cells (data not shown). In PSC patients treated with UDCA there were no visible mast cells surrounding the ducts. Images are 20x magnification

higher doses are 28–30 mg/kg BW per day. Therefore, we are slightly over the range of the lower human dose, but we also recognize that there are multiple considerations that might alter our calculations including changes in appetite, which would influence the intake of the mouse.

**UDCA treatment decreases MC presence/activation and inhibits the HDC/histamine/HR-axis in *Mdr2*<sup>-/-</sup> mice and human PSC**

In *Mdr2*<sup>-/-</sup> mice, increased numbers of MCs (marked by red arrows) are found in close proximity to bile ducts. Further, UDCA treatment decreased MC number (Fig. 2a). Semiquantification of the average number of MCs is also shown. c-Kit, chymase and tryptase decreased in *Mdr2*<sup>-/-</sup> mice fed UDCA (Supplementary Fig. 2a). In contrast, TC and TLC feeding increased MC number (marked by red arrows) in normal rats; whereas UDCA decreased MC number in rats subjected to BDL (Supplementary Fig. 1A).


Patients with advanced PSC display a robust infiltration of MCs (marked by red arrows) compared to control (similar to our previous work no MCs were detected in control [5], data not shown) and, in patients with PSC treated with UDCA, there is a marked decrease in MC presence (Fig. 2b). There is an upregulation of MC markers in livers from PSC patients treated with placebo, which are

decreased in PSC patients treated with UDCA (Supplementary Fig. 1B).

In *Mdr2*<sup>-/-</sup> mice fed UDCA, the HDC/HR/HA-axis decreases (Figs. 3a and 4b). The expression of H1–H4HR was upregulated in patients with PSC and treatment with UDCA decreased H2, H3, and H4HR (Fig. 3c). We found increased serum histamine levels in PSC patients, whereas UDCA-treated PSC patients had reduced histamine levels (Fig. 3d).


**UDCA feeding ameliorates biliary hyperplasia, proliferation, hepatic fibrosis, inflammation, and HSC activation in *Mdr2*<sup>-/-</sup> mice and human PSC**

We found that *Mdr2*<sup>-/-</sup> mice fed UDCA have decreased IBDM (Fig. 4a), biliary proliferation (Fig. 4b), and collagen deposition/fibrosis (Fig. 5a, b) compared to *Mdr2*<sup>-/-</sup> mice fed BAC. Bridging fibrosis was increased in the *Mdr2*<sup>-/-</sup> mouse fed BAC compared to WT and this was not as prominent in the *Mdr2*<sup>-/-</sup> mouse fed UDCA. Activation of HSCs was determined by SYP-9 staining and we found an increase in qualitative SYP-9 expression in *Mdr2*<sup>-/-</sup> mice fed BAC that was reduced in *Mdr2*<sup>-/-</sup> mice fed UDCA (Fig. 5c). Further, the expression of collagen type-1a, fibronectin-1 (Fig. 6a) and SYP-9 (Fig. 5c) significantly decreased in *Mdr2*<sup>-/-</sup> mice fed UDCA.




**Fig. 3** **a** HDC, H1-H4HR expression increased in *Mdr2*<sup>-/-</sup> mice fed BAC compared to WT, whereas *Mdr2*<sup>-/-</sup> mice fed UDCA had decreased levels compared to *Mdr2*<sup>-/-</sup> mice fed BAC. **b** Histamine secretion significantly increased in *Mdr2*<sup>-/-</sup> mice fed BAC and secretion diminished in *Mdr2*<sup>-/-</sup> mice fed UDCA compared to BAC fed *Mdr2*<sup>-/-</sup> mice. **c** In human patients with PSC, H1-H4HR expression increased compared to controls and in PSC patients treated

with UDCA, there was a decrease in the expression of H2HR, H3HR, and H4HR. **d** Histamine serum levels were increased in some patients with PSC and levels were decreased in one patient treated with UDCA. Data are expressed as mean ± SEM of at least 12 experiments for real-time PCR and at least 15 experiments for EIA. \**p* < 0.05 vs. WT mice or control patients; #*p* < 0.05 vs. *Mdr2*<sup>-/-</sup> mice fed BAC



**Fig. 4** **a** Bile duct mass and **(b)** Ki-67 were evaluated in liver sections from WT, *Mdr2*<sup>-/-</sup> mice + BAC and *Mdr2*<sup>-/-</sup> mice fed UDCA. We found that bile duct mass (CK-19 staining, red arrows) and Ki-67 increased in *Mdr2*<sup>-/-</sup> mice fed BAC compared to WT mice, whereas in *Mdr2*<sup>-/-</sup> mice fed UDCA, there was decreased bile duct mass (red

arrows) and a lower number of proliferating cholangiocytes (red arrows). Data have been semi-quantified and are expressed as mean ± SEM of at least 10 experiments. \**p* < 0.05 vs. WT mice; #*p* < 0.05 vs. *Mdr2*<sup>-/-</sup> mice fed BAC diet. Images are 20× magnification (inserts are 40× magnification)



**Fig. 5** **a** Staining for Masson's trichrome and **(b)** fast green/sirius red demonstrate an increase in collagen content in Mdr2<sup>-/-</sup> mice fed BAC compared to WT. UDCA feeding decreased collagen content as shown by semi-quantification of Fast Green/Sirius Red staining **(b)**. **c** Qualitative expression of SYP-9 (green staining) increased in Mdr2<sup>-/-</sup>

mice fed BAC compared to WT; whereas, treatment with UDCA decreased qualitative SYP-9 expression (bile ducts are depicted by red staining). Data are expressed as mean  $\pm$  SEM of at least nine experiments. \* $p < 0.05$  vs. WT mice; # $p < 0.05$  vs. Mdr2<sup>-/-</sup> mice fed BAC diet. Images are 40 $\times$  magnification

CD68 expression increased in Mdr2<sup>-/-</sup> mice fed BAC compared to WT and Mdr2<sup>-/-</sup> mice fed UDCA had decreased expression (Fig. 6b). TGF- $\beta$  and TNF- $\alpha$  gene expression increased in Mdr2<sup>-/-</sup> + BAC mice compared to WT; and, when Mdr2<sup>-/-</sup> mice were fed UDCA, TGF- $\beta$  and TNF- $\alpha$  expression decreased (Fig. 6c, d). Finally, we found that UDCA feeding decreased MMP-2, -3, and -9 in Mdr2<sup>-/-</sup> mice compared to BAC feeding (Supplementary Fig. 3A).

In human liver tissue fibronectin-1 expression increased in patients with PSC compared to control. UDCA treatment decreased this marker (Fig. 7a). Further, Masson's trichrome staining revealed increased collagen deposition in PSC patients compared to control; that decreased in PSC patients treated with UDCA (Fig. 7b).

### UDCA acts on MCs and alters biliary proliferation and HSC activation, in vitro


Cultured MCs express ASBT and FXR $\beta$  (Fig. 8a). When MCs were treated with UDCA, histamine secretion significantly decreased and treatment with SC-435 (blocks

ASBT) or guggulesterone (inhibits FXR $\beta$ ) returned histamine secretion to normal levels (Fig. 8b).

We found that UDCA decreased histamine secretion and the expression of c-Kit, tryptase and chymase (Fig. 9a, b). In contrast, treatment with TC, but not TLC increased the expression of MC markers (Supplementary Fig. 4A).


Stimulation with basal-treated MCs induced biliary proliferation, whereas when cholangiocytes were treated with supernatant collected from MCs pretreated with UDCA (Fig. 9c), biliary proliferation decreased (Fig. 9d). Similar to cholangiocytes, fibronectin-1 decreased in HSCs stimulated with UDCA-treated MC supernatants compared to MC basal treatments (Fig. 6e). Further, in vitro, biliary inflammatory markers TNF- $\alpha$ , CCL-3, and IL-6 increased after stimulation with MC supernatants (basal-treated) and were reduced when cholangiocytes were stimulated with MC supernatants pretreated with UDCA (Supplementary Fig. 5).

To demonstrate that MCs, cholangiocytes and HSCs work synergistically during PSC, cultured hHSCs stimulated with cholangiocyte supernatant from Mdr2<sup>-/-</sup> mice fed BAC had increased TGF- $\beta$ 1, collagen type-1a and fibronectin-1 expression compared to either basal or



**Fig. 6 a** The expression of collagen type-1a and fibronectin-1 were increased in total liver mRNA from Mdr2<sup>-/-</sup> mice fed BAC compared to WT and UDCA feeding decreased these fibrotic genes. **b** Real-time PCR for CD68 shows increased expression in Mdr2<sup>-/-</sup> mice fed BAC compared to WT, which is reduced in Mdr2<sup>-/-</sup> mice fed UDCA. **c** The


expression of TGF-β1 and **(d)** TNF-α increased in Mdr2<sup>-/-</sup> mice + BAC compared to WT and was significantly reduced in Mdr2<sup>-/-</sup> mice treated with UDCA. Data are expressed as mean ± SEM of at least nine experiments for real-time PCR. \**p* < 0.05 vs. WT mice; #*p* < 0.05 vs. Mdr2<sup>-/-</sup> mice fed BAC



**Fig. 7 a** The expression of fibronectin-1 was increased in total liver mRNA PSC patients compared to control and UDCA treatment decreased these fibrotic genes in PSC patients. **b** Masson's trichrome staining revealed increased collagen deposition in PSC patients

compared to control and this was reduced in PSC patients treated with UDCA. Data are expressed as mean ± SEM of at least 12 experiments for PCR. \**p* < 0.05 vs. control. Images are 20× magnification

**Fig. 8 a** Cultured mast cells express ASBT and FXR $\beta$  as shown by immunofluorescence and **(b)** when mast cells were treated with UDCA histamine secretion decreased. Pretreatment with inhibitors to ASBT (SC-435) or FXR $\beta$  (guggulsterone) restored histamine secretion to normal levels. Data are expressed as mean  $\pm$  SEM of at least 12 experiments for EIA. \* $p$  < 0.05 vs. basal; # $p$  < 0.05 vs. UDCA treatment



treatment with WT cholangiocyte supernatants (Fig. 10a–c). When hHSCs were stimulated with cholangiocyte supernatants from *Mdr2*<sup>-/-</sup> mice fed UDCA, fibrotic markers significantly decreased (Fig. 10a–c). Finally, in vitro UDCA decreased MC expression of MMP-2, -3, and -9 and TNF- $\alpha$  compared to basal treatment (Supplementary Fig. 3B and C), whereas TC or TLC induced the expression of MMP-2, but not MMP-3 or MMP-9 (Supplementary Fig. 4B).

## Discussion


MCs infiltrate *Mdr2*<sup>-/-</sup> mice and human PSC; and, treatment with UDCA decreases MC activation coupled with reduced biliary damage, proliferation, inflammation and hepatic fibrosis. Further, we found that UDCA reduces histamine, TGF- $\beta$ 1 and inflammatory cytokine release from MCs thus reducing PSC-associated liver injury. In vitro, we found that MCs express specific bile acid transporters and nuclear receptors and inhibition of these components increases MC activity and subsequent histamine release. Further, cholangiocytes induce a response in HSC activity that can also be regulated by UDCA, in vitro.

We have previously reported that *Mdr2*<sup>-/-</sup> mice and late stage PSC patients have increased MC numbers compared to controls [5], and studies supporting our work show

increased MC number and activation in cholangiocarcinoma, hepatocellular carcinoma, fatty liver disease and in hepatitis [24–26]. UDCA also decreases inflammatory MC number within gallbladders of patients with cholesterol-induced gallstones [27]. Further, we found that, in normal rats, TC and TLC increase MC number and in support of our findings, a study found that TC activates ileal secretion in WT mice via degranulation of MCs [28].

Alpini et al. [21] reported that, in BDL rats, UDCA feeding decreased biliary mass, bile secretion, and cholangiocyte proliferation by activation of PKC- $\alpha$ . This was further investigated in BDL-vagotomized rats demonstrating that both UDCA and its taurine conjugate, TUDCA induced protective effects on the biliary tree via calcium-induced signaling [29]. While these studies support the concept that UDCA may reverse liver damage, Fickert et al. [16] described that UDCA feeding exacerbates hepatic damage in *Mdr2*<sup>-/-</sup> mice; however, these studies were performed in different aged mice (4 weeks of age) and there was no evaluation of biliary damage. It is important to note that we utilized 12 week old *Mdr2*<sup>-/-</sup> mice that display a robust PSC phenotype when compared to earlier time points [16].

Further, we found that UDCA decreases HSC activation, TGF- $\beta$ 1 and TNF- $\alpha$  expression coupled with inhibition of hepatic fibrosis. Our previous [5, 10] and current findings





**Fig. 9** **a** Cultured mast cells treated with UDCA (40  $\mu$ M) for 30–120 min had decreased histamine secretion compared to basal treatments. **b** The expression of c-Kit, chymase, and tryptase decreased in mast cells treated with UDCA (40  $\mu$ M, 120 min) shown by real-time PCR. **c** Mast cells were treated with 0.1% BSA (basal) or UDCA (40  $\mu$ M, 120 min) and conditioned medium was collected. Cholangiocytes and HSCs were treated with conditioned medium and biliary proliferation and activation were measured. **d** In cholangiocytes treated with mast cell conditioned medium (basal) there was an increase in cholangiocyte proliferation shown by BrDU staining, whereas cholangiocyte

proliferation decreased following treatment with conditioned medium from mast cells treated with UDCA. **e** HSCs stimulated with conditioned medium from basal-treated mast cells had increased expression of fibronectin-1, which was decreased in HSCs treated with conditioned medium from mast cells treated with UDCA. Data are expressed as mean  $\pm$  SEM of at least 9 experiments for real-time PCR, 12 experiments for EIA and 10 experiments for BrDU labeling. \* $p$  < 0.05 vs. 0.1% BSA (basal); # $p$  < 0.05 + MC (basal) treatment. Images are 20 $\times$  magnification

describe a system whereby MCs are recruited to the damaged liver, interact with proliferating cholangiocytes inducing activation of HSCs, thus promoting hepatic fibrosis. In support of this, in rats with CCl<sub>4</sub>-induced liver fibrosis, treatment with UDCA improved histopathological scores, decreased tissue hydroxyproline and inhibited MMP activation with overall amelioration of hepatic fibrosis [30]. Further, in nonalcoholic steatohepatitis, UDCA was found to be beneficial at repressing hepatic fibrosis and HSC activation by counteracting intestinal barrier dysfunction [31]. While the usage of UDCA for reducing fibrosis in PSC has been controversial, UDCA is useful in PBC management by decreasing inflammation and fibrosis [12]. Further, the derivative of UDCA, nor-UDCA, shows promising results in both animals and humans to reduce and ameliorate fibrosis in PSC and PBC [1, 2].

MCs release numerous inflammatory mediators including histamine, TGF- $\beta$ 1 and TNF- $\alpha$ , all of which were reduced in Mdr2<sup>-/-</sup> mice fed UDCA. During ulcerative colitis there is an increase in inflammatory cytokines, that are mostly ablated in MC-deficient rats [32]. Further, TNF- $\alpha$  increases in lung tissues following autologous liver transplantation; however, when animals were treated with the MC stabilizers, cromolyn sodium, or ketotifen prior to transplantation, inflammatory mediators significantly reduced [33]. Additionally, BDL rats treated with UDCA displayed a significant decrease in TGF- $\beta$ 1 expression compared to control rats [34] and He et al. [35] found that the addition of retinoic acid to UDCA treatment decreased TGF- $\beta$ 1 and collagen deposition in BDL rats. The link between UDCA and inflammation has also been demonstrated in fatty liver disease. Pathil et al. [36] found that

**Fig. 10** The expression of (a) collagen type-1a, (b) TGF- $\beta$ 1, and (c) fibronectin-1 were increased in hHSCs treated with cholangiocyte supernatant from Mdr2<sup>-/-</sup> mice fed BAC compared to basal and WT supernatant treatments; however, expression of these fibrotic markers were decreased when hHSCs were stimulated with cholangiocyte supernatants from Mdr2<sup>-/-</sup> mice fed UDCA. Data are expressed as mean  $\pm$  SEM of at least six experiments for real-time PCR. \*  $p < 0.05$  vs. WT mice; #  $p < 0.05$  vs. Mdr2<sup>-/-</sup> mice fed BAC



**Fig. 11** Working model depicting a summary of the findings. Mdr2<sup>-/-</sup> mice fed BAC have upregulated mast cell number, HDC/HA/HR-axis, biliary hyperplasia, HSC activation, fibrosis, inflammation and serum TBAs. These parameters are all reduced in Mdr2<sup>-/-</sup> mice fed UDCA.

Further, patients treated with UDCA have decreased mast cell number, inhibited HA/HR axis, HSC activation and fibrosis and inflammation compared to placebo-treated patients. Finally, *in vitro*, UDCA acts on mast cells to decrease biliary proliferation and HSC activation

treatment with a conjugate of UDCA improved liver histology, steatosis and decreased proinflammatory cytokines in mice fed a high fat diet. In our study, we found that UDCA decreased MMP expression. Similar to this, it has

been previously demonstrated that UDCA induced beneficial effects on liver ALT levels and inhibited MMPs leading to the destruction of extracellular matrix in the damaged liver [14].

Cholangiocytes express FXR $\beta$  and ASBT [37]. Similarly, we found that MCs express these components (along with others, data not shown) and may use these to allow UDCA (or other bile acids) to induce effects on MC degranulation. Whereas, UDCA has been demonstrated to be a weak ligand for FXR in some studies [38], in our work, we found that blocking FXR $\beta$  restored MC-histamine release (inhibited by UDCA treatment) suggesting that, in MCs, the UDCA/FXR interaction is important. Further, Mueller et al. [39] recently reported that UDCA exerts FXR antagonistic effects on patients with NAFLD. ASBT is a critical regulator of conjugated bile acids especially in cholangiocyte regulation [40, 41] and we demonstrated that inhibition of ASBT or FXR returns histamine levels back to normal following UDCA treatment. In support of our studies, it has been previously demonstrated that bile acids influence MC degranulation [42] and we found that UDCA decreases MC-histamine release, MC marker expression, biliary proliferation, and HSC activation, *in vitro*. Interestingly, a recent study has shown that the ASBT inhibitor itself may be a promising tool for reducing PSC progression and clinical trials are underway [43]. In this work the authors find that treatment with SC-435 decreases inflammation, damage and fibrosis in the Mdr2<sup>-/-</sup> mouse model, which are interesting since we found that blocking ASBT returns histamine levels back to normal following UDCA treatment, *in vitro*. Considering that the authors used *in vivo* models and our studies on the ASBT inhibitor were *in vitro*, more experimental analysis should be performed to fully understand if ASBT alters MCs during disease progression.

In conclusion, our studies are the first to demonstrate that UDCA decreases MC number and histamine release, thereby inhibiting PSC-induced inflammation. Our novel data suggest an alternative strategy for UDCA action that also includes regulation of MC function (Fig. 11). Further studies are warranted to fully understand the implications of bile acid regulation on MC activation including determining the effects of nor UDCA, which has been shown to be a promising therapy for PSC patients [44].

**Funding** Portions of this work were supported by (i) a VA Merit Awards (1I01BX003031, H.F.; 4I01BX000574, G.A.; 1I01BX001724, F.M.; and 1I01BX002638, S.D.) from the United States Department of Veteran's affairs, Biomedical Laboratory Research and Development Service and R01 grants from NIH NIDDK (DK108959, H.F. and DK082435, S.D.); (ii) funds from the PSC Partners Seeking a Cure (H.F. and S.D.); (iii) a Baylor Scott and White Research Mentor Award (H.F.); and (iv) the Dr. Nicholas C. Hightower Centennial Chair of Gastroenterology from Baylor Scott & White Health (GA). This material is the result of work supported with resources and the use of facilities at the Central Texas Veterans Health Care System, Temple, Texas. The content is the responsibility of the author(s) alone and does not necessarily reflect the views or policies of the Department of Veterans Affairs or the United States Government.

## Compliance with ethical standards

**Conflict of interest** The authors declare that they have no conflict of interest.

## References

- Williamson KD, Chapman RW. New therapeutic strategies for primary sclerosing cholangitis. *Semin Liver Dis.* 2016;36:5–14.
- Halilbasic E, Fuchs C, Hofer H, et al. Therapy of primary sclerosing cholangitis—today and tomorrow. *Dig Dis.* 2015;33(Suppl 2):S149–63.
- Popov Y, Patsenker E, Fickert P, et al. Mdr2 (Abcb4)<sup>-/-</sup> mice spontaneously develop severe biliary fibrosis via massive dysregulation of pro- and antifibrogenic genes. *J Hepatol.* 2005;43:1045–54.
- Trauner M, Fickert P, Wagner M. MDR3 (ABCB4) defects: a paradigm for the genetics of adult cholestatic syndromes. *Semin Liver Dis.* 2007;27:77–98.
- Jones H, Hargrove L, Kennedy L, et al. Inhibition of mast cell-secreted histamine decreases biliary proliferation and fibrosis in primary sclerosing cholangitis Mdr2<sup>-/-</sup> mice. *Hepatology.* 2016;64:1202–16.
- Bachelet I, Levi-Schaffer F, Mekori YA. Mast cells: not only in allergy. *Immunol Allergy Clin North Am.* 2006;26:407–25.
- Beaven MA. Our perception of the mast cell from Paul Ehrlich to now. *Eur J Immunol.* 2009;39:11–25.
- Boyce JA. Mast cells: beyond IgE. *J Allergy Clin Immunol.* 2003;111:24–32.
- Kennedy LL, Hargrove LA, Graf AB, et al. Inhibition of mast cell-derived histamine secretion by cromolyn sodium treatment decreases biliary hyperplasia in cholestatic rodents. *Lab Invest.* 2014;94:1406–18.
- Hargrove L, Kennedy L, Demieville J, et al. BDL-induced biliary hyperplasia, hepatic injury and fibrosis are reduced in mast cell deficient Kitw-sh mice. *Hepatology.* 2017;65:1991–2004.
- Beuers U, Trauner M, Jansen P, et al. New paradigms in the treatment of hepatic cholestasis: from UDCA to FXR, PXR, and beyond. *J Hepatol.* 2015;62:S25–37.
- de Vries E, Beuers U. Management of cholestatic disease in 2017. *Liver Int.* 2017;37(Suppl 1):S123–9.
- Penz-Osterreicher M, Osterreicher CH, Trauner M. Fibrosis in autoimmune and cholestatic liver disease. *Best Pract Res Clin Gastroenterol.* 2011;25:245–58.
- Buryova H, Chalupsky K, Zbodakova O, et al. Liver protective effect of ursodeoxycholic acid includes regulation of ADAM17 activity. *BMC Gastroenterol.* 2013;13:155.
- European Association for the Study of the L. EASL Clinical Practice Guidelines: management of cholestatic liver diseases. *J Hepatol.* 2009;51:237–67.
- Fickert P, Zollner G, Fuchsbichler A, et al. Ursodeoxycholic acid aggravates bile infarcts in bile duct-ligated and Mdr2 knockout mice via disruption of cholangioles. *Gastroenterology.* 2002;123:1238–51.
- Graf A, Meng F, Hargrove L, et al. Knockout of histidine decarboxylase decreases bile duct ligation-induced biliary hyperplasia via downregulation of the histidine decarboxylase/VEGF axis through PKA-ERK1/2 signaling. *Am J Physiol Gastrointest Liver Physiol.* 2014;307:G813–823.
- Halova I, Draberova L, Draber P. Mast cell chemotaxis—chemoattractants and signaling pathways. *Front Immunol.* 2012;3:119.
- Alpini G, Glaser SS, Ueno Y, et al. Bile acid feeding induces cholangiocyte proliferation and secretion: evidence for bile acid-regulated ductal secretion. *Gastroenterology.* 1999;116:179–86.

20. Alpini G, Ueno Y, Glaser SS, et al. Bile acid feeding increased proliferative activity and apical bile acid transporter expression in both small and large rat cholangiocytes. *Hepatology*. 2001;34:868–76.
21. Alpini G, Baiocchi L, Glaser S, et al. Ursodeoxycholate and tauroursodeoxycholate inhibit cholangiocyte growth and secretion of BDL rats through activation of PKC alpha. *Hepatology*. 2002;35:1041–52.
22. Carpino G, Franchitto A, Morini S, et al. Activated hepatic stellate cells in liver cirrhosis. A morphologic and morphometrical study. *Ital J Anat Embryol*. 2004;109:225–38.
23. Carpino G, Morini S, Ginanni Corradini S, et al. Alpha-SMA expression in hepatic stellate cells and quantitative analysis of hepatic fibrosis in cirrhosis and in recurrent chronic hepatitis after liver transplantation. *Dig Liver Dis*. 2005;37:349–56.
24. Farrell DJ, Hines JE, Walls AF, et al. Intrahepatic mast cells in chronic liver diseases. *Hepatology*. 1995;22:1175–81.
25. Francis H, Meininger CJ. A review of mast cells and liver disease: what have we learned? *Dig Liver Dis*. 2010;42:529–36.
26. Hodges K, Kennedy L, Meng F, et al. Mast cells, disease and gastrointestinal cancer: A comprehensive review of recent findings. *Transl Gastrointest Cancer*. 2012;1:138–50.
27. Carotti S, Guarino MP, Cicala M, et al. Effect of ursodeoxycholic acid on inflammatory infiltrate in gallbladder muscle of cholesterol gallstone patients. *Neurogastroenterol Motil*. 2010;22:866–73. e232
28. Hardcastle J, Hardcastle PT, Chapman J, et al. Taurocholic acid-induced secretion in normal and cystic fibrosis mouse ileum. *J Pharm Pharmacol*. 2001;53:711–9.
29. Marzioni M, Francis H, Benedetti A, et al. Ca<sup>2+</sup>-dependent cytoprotective effects of ursodeoxycholic and tauroursodeoxycholic acid on the biliary epithelium in a rat model of cholestasis and loss of bile ducts. *Am J Pathol*. 2006;168:398–409.
30. Tasci I, Mas MR, Vural SA, et al. Rat liver fibrosis regresses better with pegylated interferon alpha2b and ursodeoxycholic acid treatments than spontaneous recovery. *Liver Int*. 2006;26:261–8.
31. Namisaki T, Noguchi R, Moriya K, et al. Beneficial effects of combined ursodeoxycholic acid and angiotensin-II type 1 receptor blocker on hepatic fibrogenesis in a rat model of nonalcoholic steatohepatitis. *J Gastroenterol*. 2016;51:162–72.
32. Chu HQ, Li J, Huang HP, et al. Protective effects of tranilast on oxazolone-induced rat colitis through a mast cell-dependent pathway. *Dig Liver Dis*. 2016;48:162–71.
33. Zhang A, Chi X, Luo G, et al. Mast cell stabilization alleviates acute lung injury after orthotopic autologous liver transplantation in rats by downregulating inflammation. *PLoS ONE*. 2013;8:e75262.
34. Yang YY, Huang YT, Lee KC, et al. Chronic administration of ursodeoxycholic acid decreases portal pressure in rats with biliary cirrhosis. *Clin Sci (Lond)*. 2009;116:71–79.
35. He H, Mennone A, Boyer JL, et al. Combination of retinoic acid and ursodeoxycholic acid attenuates liver injury in bile duct-ligated rats and human hepatic cells. *Hepatology*. 2011;53:548–57.
36. Pathil A, Mueller J, Warth A, et al. Ursodeoxycholy l lysophosphatidylethanolamide improves steatosis and inflammation in murine models of nonalcoholic fatty liver disease. *Hepatology*. 2012;55:1369–78.
37. Halilbasic E, Claudel T, Trauner M. Bile acid transporters and regulatory nuclear receptors in the liver and beyond. *J Hepatol*. 2013;58:155–68.
38. Makishima M, Okamoto AY, Repa JJ, et al. Identification of a nuclear receptor for bile acids. *Science*. 1999;284:1362–5.
39. Mueller M, Thorell A, Claudel T, et al. Ursodeoxycholic acid exerts farnesoid X receptor-antagonistic effects on bile acid and lipid metabolism in morbid obesity. *J Hepatol*. 2015;62:1398–404.
40. Alpini G, Glaser S, Baiocchi L, et al. Secretin activation of the apical Na<sup>+</sup>-dependent bile acid transporter is associated with cholehepatic shunting in rats. *Hepatology*. 2005;41:1037–45.
41. Fava G, Marzioni M, Francis H, et al. Novel interaction of bile acid and neural signaling in the regulation of cholangiocyte function. *Hepatol Res*. 2007;37(Suppl 3):S420–429.
42. Quist RG, Ton-Nu HT, Lillienau J, et al. Activation of mast cells by bile acids. *Gastroenterology*. 1991;101:446–56.
43. Miethke AG, Zhang W, Simmons J, et al. Pharmacological inhibition of apical sodium-dependent bile acid transporter changes bile composition and blocks progression of sclerosing cholangitis in multidrug resistance-2 knockout mice. *Hepatology*. 2016;63:512–23.
44. Fickert P, Hirschfield GM, Denk G, et al. nor-Ursodeoxycholic acid improves cholestasis in primary sclerosing cholangitis. *J Hepatol*. 2017;67:549–58.