

Query Details

Back to Main Page

1. Author names: Please confirm if the author names are presented accurately and in the correct sequence (given name, middle name/initial, family name). Also, kindly confirm the details in the metadata are correct.

OK
2. Please check and confirm that the authors and their respective affiliations have been correctly identified and amend if necessary.

OK
3. References [14] and [20] are same, we have deleted the duplicate reference and renumbered accordingly. Please check and confirm.

OK
4. Please check the layout of Tables and correct if necessary.

OK
5. Please provide a definition for the significance of [bold] in the table 2, 3 and 4

Statistically significant p-values are highlighted in bold to distinguish them from non-significant values
6. Data availability statement is preferred for this journal. If applicable, please provide

OK. This is the Data Availability Statement; you can insert it where indicated in the paper.

The data generated from this survey are available from the EuGMS Special Interest Group (SIG) on Delirium upon reasonable request and subject to approval by the SIG.
7. Journal instruction requires a Ethical approval and Informed consent section; however, none was provided. In this regard, please provide the missing section.

At the beginning of the Methods section, we added a statement on ethical approval for the survey and a statement confirming that participants agreed to participate and accepted the conditions of participation. This topic was also discussed with the reviewers during the peer-review process.

Research Paper

Bridging the gap in delirium management: a European comparison of geriatric practices

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

Alessandra Coin ✉
Email : alessandra.coin@unipd.it
Affiliationids : Aff1 Aff2, Correspondingaffiliationid : Aff1
Silvia Sturani Affiliationids : Aff1
Christian Pozzi Affiliationids : Aff3
Maela Caudal Affiliationids : Aff4
Duygu Erbas Sacar Affiliationids : Aff5
Massimiliano Fedecostante Affiliationids : Aff6
Maria Cristina Ferrara Affiliationids : Aff7

Antonio Cherubini Affiliationids : Aff6 Aff8

Alessandro Morandi Affiliationids : Aff9 Aff10

Gülistan Bahat Affiliationids : Aff11

Giuseppe Bellelli Affiliationids : Aff7 Aff12

Suzanne Timmons Affiliationids : Aff13

the S. I. G. Delirium of the European Geriatric Medicine Society

Aff1 Department of Medicine (DIMED), University of Padova, Padua, Italy

Aff2 Unit of Geriatrics, University Padova Hospital, Padua, Italy

Aff3 Centre of Competence on Ageing, University of Applied Sciences and Arts of Southern Switzerland (SUPSI), Lugano, Switzerland

Aff4 Altersmedizinisches Zentrum (Center of Geriatric Medicine), St. Marien-Hospital, Cologne, Germany

Aff5 Department of Internal Medicine, Division of Geriatrics, Kartal Dr. Lutfi Kirdar City Hospital, Istanbul, Türkiye

Aff6 **Geriatricia, Accettazione Geriatrica e Centro di Ricerca Per L'invecchiamento** Geriatricia, Accettazione Geriatrica e Centro di Ricerca per l'Invecchiamento, IRCCS INRCA, Ancona, Italy

Aff7 School of Medicine and Surgery, University of Milano-Bicocca, Milan, Italy

Aff8 Department of Clinical and Molecular Sciences, Università Politecnica delle Marche, Ancona, Italy

Aff9 Department of Clinical and Experimental Science, University of Brescia, Brescia, Italy

Aff10 Azienda Speciale Cremona Solidale, Cremona, Italy

Aff11 Division of Geriatrics, Department of Internal Medicine, Istanbul Medical Faculty, Istanbul University, Capa, Istanbul, Türkiye

Aff12 Acute Geriatric Unit, IRCCS Foundation San Gerardo dei Tintori, Monza, Italy

Aff13 Centre for Gerontology and Rehabilitation, School of Medicine, University College Cork, Cork, Ireland

Received: 9 February 2026 / Revised: 2 May 2026 / Accepted: 3 June 2026

Key summary points

Aim

To compare diagnostic and therapeutic approaches to delirium across European countries with well-established versus less-developed Geriatric Medicine (GM), to identify similarities, differences, and potential gaps in care.

Findings

Countries with well-established GM more frequently reported structured delirium screening, established management protocols, and the use of non-pharmacological interventions. In contrast, countries with less-developed GM showed higher reliance on pharmacological treatments and physical restraints.

Message

Marked differences in delirium management persist across Europe and appear to be associated with the level of development of Geriatric Medicine. These findings highlight the need for harmonized European guidelines to promote equitable, evidence-based delirium care.

Abstract

Aim

To compare diagnostic and therapeutic approaches to delirium across European countries with well-established (WE) versus less-developed (LE) geriatric medicine (GM), to identify similarities, differences and potential gaps in care.

Methods

A cross-sectional online survey was conducted among European geriatricians via the European Geriatric Medicine Society and professional networks from July 2023 to March 2024. **AQ1** Responses on delirium diagnosis and management were analyzed quantitatively, with comparisons between WE and LDGM countries.

Results

Of 196 hospital-based respondents, 79.1% were from WEGM countries and 20.9% from LDGM countries. Structured delirium screening was significantly more common in WEGM nations, especially in geriatric (88.4% vs 43.9%, $p < 0.001$) and orthopedic wards. Non-pharmacological strategies—such as hydration support (76.8% vs 52.5%, $p = 0.002$) and family involvement (81.5% vs 60.4%, $p = 0.005$)—were also more frequently used. **AQ2** Conversely, physical restraints were more commonly employed in LDGM countries (70.7% vs 41.4%, $p < 0.001$), as were pharmacological treatments, such as trazodone and quetiapine.

Conclusion

Marked differences exist in delirium management across Europe, which may be linked at least in part to GM development, although confounding factors may exist, and causality cannot be assumed. WEGM countries appear to prioritize structured screening and non-pharmacological interventions, whereas LDGM countries rely more on pharmacological treatments. These findings highlight the need for standardized European guidelines to ensure equitable delirium care.

Keywords

Delirium
Geriatric Medicine
Older adults
European survey
Health service
Inequality

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1007/s41999-026-01527-6>.

Introduction

Delirium is an acute geriatric syndrome affecting attention, cognition, and psychomotor activity, with a fluctuating course, which occurs primarily in older adults [1]. It often arises due to underlying medical conditions [2] and is strongly associated with frailty, particularly in acute [3,4,5] and long-term care settings.

Prevalence varies by setting: 1–2% in community-dwelling older adults, rising to 14–24% in hospitalized patients and up to 80% in critical care [6,7,8,9]. Often underrecognized, delirium leads to increased disability, mortality, institutionalization, dementia risk, and substantial healthcare costs [10,11,12].

Accurate investigation requires cognitive assessment and a thorough history. National guidelines recommend routine delirium screening using validated tools, with the Confusion Assessment Method (CAM) and the 4 ‘A’s Test (4AT) being the most widely used [13,14,15,16,17]. Despite recommendations to screen all older emergency admissions for dementia and delirium [14], no consensus exists on standardized evaluation methods [18].

Non-pharmacological interventions effectively reduce delirium risk [14,19] and growing evidence supports multimodal, multidisciplinary approaches as highly effective [20,21,22,23].

In addition to delirium prevention, standardized treatment guidelines are essential. Currently, there are no European-specific guidelines for delirium management. Addressing modifiable causes identified during evaluation is critically important, as even minor interventions can have a significant impact [6,24].

No universally accepted pharmacological treatment exists. The use of antipsychotics and sedatives is not recommended, except in cases of hyperactive delirium, for short durations, and only when the patient is in severe distress or at risk of disrupting life-sustaining treatments. Nonetheless, such medications are still commonly used in clinical practice [25]. Low doses of haloperidol are considered as first-line agent in the symptomatic management of delirium episodes [26]. These data underscore the importance of prioritizing non-pharmacological strategies and individualized risk assessment in the prevention and management of delirium in older adults [27,28]. Variability in both pharmacological and non-pharmacological approaches complicates standardization, further influenced by cultural differences across countries.

Moreover, although several national and international guidelines exist, there is currently no unified European framework specifically addressing delirium management, which may contribute to heterogeneity in clinical practice across countries [24, 29].

Considering the variability in delirium management and the lack of homogeneity in following guidelines across European countries, another relevant aspect to address is the potential variability between nations where Geriatric Medicine is an established medical specialty and those where it is not yet formally recognized or integrated into postgraduate medical education. This study compares diagnostic and therapeutic approaches for delirium between countries with well-established (WE) versus less-developed (LD) Geriatric Medicine (GM) medical specialty, aiming to outline the similarities and differences, and thereby contribute to the recognition of the need for, and the development of, standardized guideline measures across Europe [29].

Methods

The survey was created and revised throughout the members of the research team, who are all geriatricians. Four of the authors considered to be experts in the field (GB, AC, ST, AM) gave feedback on the face and content validity of each single question and on the overall questionnaire.

After discussion in the working group and with the Ethical Committees of the University of Padova and the University of Dublin, it was decided that it was not necessary to have an ethical consent due to the nature of the survey, not involving patients, but only asking opinion of physicians in an anonymous way.

The survey was structured in 4 sections (Supplementary file 1).

At the beginning of the survey, participants were informed about the aim of this survey, an estimation of how much time was needed to fulfill it, and that their participation accepted the conditions. Participation was voluntary, and completion of the survey was considered implied informed consent. Responses reflect clinicians' reported perceptions of practices within their unit or hospital rather than externally verified institutional policies. The survey was mainly multiple-choice questions, but the participant was given the possibility to give more detailed information about some questions. No patients' data were acquired and the answers were anonymous. In the first section, a brief overview of participant characteristics was requested, while respecting their anonymity. In the second section, the participant was requested to answer questions about the different wards present in his/her hospital, the existing delirium diagnostic pathway and delirium-frailty pathway in those wards. Specifically, the considered wards were: geriatric, ICU, emergency, orthopedic, medical ward, surgery ward, rehabilitation and long-term facilities. However, the answers concerning rehabilitation and long-term facilities were not considered for the present manuscript (Supplementary Table 1). The aim in this section was to determine if a structured pathway to identify delirium in the different wards exist.

In the third section, participants were asked about the usual treatment of delirium in their hospital. Here, there was no difference made between the wards. The use of non-pharmacological methods was investigated by multiple-choice questions. For each positive answer, a percentage for the use of this method among patients was asked (< 25%; 25–50%; 50–75%; > 75%). Finally, pharmacological methods were examined. For each medication, the participant could specify the usual dose and the type of administration. In this section, the aim was to identify the usual methods used to treat delirium with pharmacological as well as non-pharmacological methods.

The survey was conducted among geriatricians who have an affiliation to the EuGMS (European Geriatric Medicine Society) or were contacted through professional networks and are working in Europe. It was presented as a Google form in English. It was distributed by email through the EuGMS and cascaded in a snowball approach via professional networks. The authors also distributed the questionnaire through their respective national or in-country professional networks.

The survey was launched in July 2023 and closed in March 2024. Participants were asked to provide answers about their own wards and the other wards of their hospital.

Analysis

After excluding answers from countries outside of Europe and incomplete surveys, responses obtained by participants were divided into two groups according to the presence of a recognized primary specialty in geriatric medicine (GM) in the represented countries [29]: well-established GM (Belgium, Czech Republic, Denmark, Finland, France, Germany, Iceland, Ireland, Italy, Lithuania, Luxembourg, Malta, Norway, Poland, Romania, Spain, Sweden, The Netherlands, United Kingdom) and less-developed GM (Belarus, Greece, Portugal, Slovenia, Switzerland, Türkiye). This classification is based on a previously published European framework reflecting the recognition of geriatric medicine as a primary specialty and its integration into postgraduate training pathways and dedicated geriatric services. Quantitative survey data were analyzed statistically. Nominal data were reported as numbers and percentages. The Chi-Square test was conducted to examine differences between countries. Statistical analyses were performed using SPSS 26.0 (IBM Corp.) It was hypothesized that countries with well-established GM had a better management of delirium, e.g. higher rate of delirium diagnosis, as well as a more frequent non-pharmacological treatment.

Results

In total, 238 physicians accessed and completed the survey. Thirteen responses coming from countries outside Europe and 6 incomplete ones were excluded. Because of the aim of the study, 23 responses from a non-hospital setting were also excluded, giving a final sample of 196 participants. According to the pre-defined classification, 41 (20.9%) participants were from countries with less-established GM while 155 (79.1%) were from countries with well-established GM (Table 1).

Table 1

List of survey participants countries

	N (%)
Less-developed GM (N = 41)	
Belarus	1 (2.4)
Greece	4 (9.8)
Portugal	10 (24.4)
Slovenia	1 (2.4)
Switzerland	12 (29.3)
Turkey	13 (31.7)
Well-established GM (N = 155)	
Belgium	14 (9)
Czech Republic	2 (1.3)
Denmark	12 (7.7)
Finland	2 (1.3)
France	16 (10.3)
Germany	9 (5.8)
Iceland	1 (0.6)
Ireland	6 (3.9)
Italy	17 (11)
Lithuania	1 (0.6)
Luxemburg	1 (0.6)
Malta	2 (1.3)
Norway	12 (7.7)
Poland	8 (5.2)
Romania	2 (1.3)
Spain	19 (12.3)
Sweden	4 (2.6)
The Netherlands	10 (6.5)
United Kingdom	17 (11)

General characteristics of the structures where participants work, the prevalence of screening tools, and the prevalence of protocols and pathways for delirium and frailty according to different medical or surgical wards are reported in Table 2. Participants were mainly from University or University-related hospitals (73.0%), with an equal distribution of the hospital size from < 250 to > 1000 beds. A geriatric ward was present in 79.1% of hospitals, with a significant higher prevalence in countries with well-established GM (88.4% vs. 43.9%; $p < 0.001$). The use of a screening tool was more frequently reported in well-established GM countries both in geriatric wards (71.6% vs. 34.1%; $p < 0.001$) and in orthopedic wards (49.0% vs. 19.5%; $p = 0.002$). Likewise, the presence of a protocol for patients with delirium was more common in well-established GM countries in geriatric wards (52.3% vs. 29.3%; $p < 0.001$) and orthopedic wards (40.6% vs. 17.1%; $p = 0.007$).

Table 2

Survey: characteristics of the hospital facilities where participants work

--	--	--	--

Variables:	Less-developed GM (N = 41)	Well-established GM (N = 155)	P-value
Type of structure	Number (percent)		< 0.001
Community hospital	11 (26.8)	29 (18.7)	
Private hospital	8 (19.5)	5 (3.2)	
University hospital	16 (39)	90 (58.1)	
University-related/affiliated hospital	6 (14.6)	31 (20)	
Size of the hospital (beds)			0.712
< 250	13 (31.7)	37 (23.9)	
250–500	8 (19.5)	39 (25.2)	
500–1000	11 (26.8)	47 (30.3)	
> 1000	9 (22)	32 (20.6)	
Presence of a geriatric ward	18 (43.9)	137 (88.4)	< 0.001
Screening tool for delirium	14 (34.1)	111 (71.6)	< 0.001
Protocol for patients with delirium	12 (29.3)	81 (52.3)	< 0.001
Pathway including frailty assessment	8 (19.5)	52 (33.5)	0.066
Pathway including frailty bundle care plan	8 (19.5)	48 (31)	0.073
Clear link with frailty bundle care plan	4 (9.8)	42 (27.1)	0.017
Referral for follow-up of cognition	9 (22)	56 (36.1)	0.065
Presence of an Emergency Room	35 (85.4)	134 (86.5)	0.858
Screening tool for delirium	10 (24.4)	74 (47.7)	0.034
Protocol for patients with delirium	7 (17.1)	52 (33.5)	0.229
Pathway including frailty assessment	4 (9.8)	36 (23.2)	0.237
Pathway including frailty bundle care plan	5 (12.2)	29 (18.7)	0.785
Clear link with frailty bundle care plan	3 (7.3)	25 (16.1)	0.464
Referral for follow-up of cognition	4 (9.8)	27 (17.4)	0.690
Presence of an ICU	36 (87.8)	136 (87.7)	0.991
Screening tool for delirium	19 (46.3)	67 (43.2)	0.826
Protocol for patients with delirium	11 (26.8)	38 (24.5)	0.240
Pathway including frailty assessment	4 (9.8)	14 (9)	0.676
Pathway including frailty bundle care plan	5 (12.2)	13 (8.4)	0.082
Clear link with frailty bundle care plan	5 (12.2)	12 (7.7)	0.178
Referral for follow-up of cognition	6 (14.6)	11 (7.1)	0.364
Presence of a Medical ward	36 (87.8)	144 (92.9)	0.289
Screening tool for delirium	16 (39)	76 (49)	0.510
Protocol for patients with delirium	14 (34.1)	59 (38.1)	0.174
Pathway including frailty assessment	6 (14.6)	25 (16.1)	0.285
Pathway including frailty bundle care plan	7 (17.1)	27 (17.4)	0.242
Clear link with frailty bundle care plan	5 (12.2)	26 (16.8)	0.619
Referral for follow-up of cognition	10 (24.4)	26 (16.8)	0.169
Presence of a Surgical ward	33 (80.5)	139 (89.7)	0.110
Screening tool for delirium	9 (22)	55 (35.5)	0.193
Protocol for patients with delirium	7 (17.1)	42 (27.1)	0.195
Pathway including frailty assessment	3 (7.3)	21 (13.5)	0.402
Pathway including frailty bundle care plan	3 (7.3)	22 (14.2)	0.532
Clear link with frailty bundle care plan	3 (7.3)	17 (11)	0.307
Referral for follow-up of cognition	4 (9.8)	19 (12.3)	0.722
Presence of an Orthopedic ward	30 (73.2)	130 (83.9)	0.116
Screening tool for delirium	8 (19.5)	76 (49)	0.002

Protocol for patients with delirium	7 (17.1)	63 (40.6)	0.007
Pathway including frailty assessment	3 (7.3)	43 (27.7)	0.047
Pathway including frailty bundle care plan	3 (7.3)	40 (25.8)	0.057
Clear link with frailty bundle care plan	3 (7.3)	37 (23.9)	0.086
Referral for follow-up of cognition	4 (9.8)	40 (25.8)	0.172

The different non-pharmacological and pharmacological management strategies for delirium are shown in Tables 3 and 4. Countries with a well-established GM discipline showed a significantly higher prevalence of non-pharmacological approaches, such as support to increase fluid intake, open or liberal visiting times for families, avoidance of unnecessary bladder tubes or catheters, reduction of drugs potentially associated with delirium, and more frequent (> 50% of times) use of patient mobilization. There was also a significantly less frequent use of physical restraints in countries with well-developed GM compared to those with less-developed GM. Figure 1 shows the geographical distribution of countries based on whether non-pharmacological strategies are reported to be used in more or less than 50% of cases. The responses are visualized using a gradient of blue shades.

Table 3

The table shows responses on the use of non-pharmacological intervention strategies, including the number (and percentage) of affirmative responses for each strategy and those with a utilization rate > 50%

Variables	Less-developed GM (N = 41)	Well-established GM (N = 155)	P-value
Mobilization	35 (85.4)	148 (95.5)	0.061
of which, use > 50%	18 (51.4)	107 (72.3)	0.017
Support to increase mobility	20 (48.8)	92 (59.4)	0.243
of which, use > 50%	13 (65)	54 (58.7)	0.602
Pain assessment	41 (100)	146 (94.2)	0.196
of which, use > 50%	33 (80.5)	121 (82.9)	0.723
Pain management	29 (70.7)	112 (72.3)	0.734
of which, use > 50%	21 (72.4)	79 (70.5)	0.843
Physical restraints	31 (75.6)	105 (67.7)	0.448
of which, use > 50%	12 (38.7)	16 (15.2)	0.005
Reactivate day and night cycle	35 (85.4)	130 (83.9)	0.871
of which, use > 50%	23 (65.7)	89 (68.5)	0.757
Support to increase fluid intake	30 (73.2)	130 (83.9)	0.021
of which, use > 50%	25 (80)	96 (73.8)	0.483
Provision of hearing and mobility aids	33 (80.5)	132 (85.2)	0.174
of which, use > 50%	23 (69.7)	101 (76.5)	0.418
Cognitive stimulation	28 (68.3)	105 (67.7)	0.651
of which, use > 50%	18 (64.3)	52 (49.5)	0.165
Verbal reorientation	39 (95.1)	145 (93.5)	0.941
of which, use > 50%	31 (79.5)	125 (86.2)	0.300
Open or liberal visiting times for families	35 (85.4)	144 (92.9)	0.040
of which, use > 50%	28 (80)	117 (81.3)	0.866
Caregiver education for proactive and informed caregiving	29 (70.7)	117 (75.5)	0.171
of which, use > 50%	17 (58.6)	77 (65.8)	0.469
Transferring to specific delirium rooms	3 (7.3)	25 (16.1)	0.148
of which, use > 50%	0 (0)	11 (44)	0.140
Non-disturbed sleep	27 (65.9)	96 (61.9)	0.575
of which, use > 50%	19 (70.4)	65 (67.7)	0.793
Avoidance of unnecessary bladder tubes or catheters	36 (87.8)	145 (93.5)	0.045
of which, use > 50%	27 (75)	114 (78.6)	0.639
Sharing or communicating patient information about delirium	38 (92.7)	145 (93.5)	0.608

of which, use > 50%	33 (86.8)	124 (85.5)	0.835
Ground-leveled beds	15 (36.6)	71 (45.8)	0.238
of which, use > 50%	5 (33.3)	26 (36.6)	0.810
Group activities for patients	14 (34.1)	49 (31.6)	0.741
of which, use > 50%	6 (42.9)	13 (26.5)	0.240
Animal assisted therapy	3 (7.3)	12 (7.7)	0.919
of which, use > 50%	1 (33.3)	2 (16.7)	0.519
Going outside the unit ward	15 (36.6)	35 (22.6)	0.083
of which, use > 50%	4 (26.7)	6 (17.1)	0.440
Reduction of potentially delirigenic drugs	37 (90.2)	149 (96.1)	0.041
of which, use > 50%	29 (78.4)	128 (85.9)	0.259


Table 4

The table shows responses on the use of pharmacological intervention strategies, including the number (and percentage) of affirmative responses for each strategy and those with a utilization rate > 50%

Variables	Less-established GM (N = 41)	Well-established GM (N = 155)	P-value
Delirious patients receiving pharmacological intervention	41 (100)	152 (98.1)	0.369
of which, use > 50%	29 (70.7)	63 (41.4)	< 0.001
Haloperidol	32 (78)	122 (78.7)	0.927
Clonidine	1 (2.4)	2 (1.3)	0.594
Promazine	0 (0)	7 (4.5)	0.166
Trazodone	23 (56.1)	43 (27.7)	< 0.001
Melperone	1 (2.4)	10 (6.5)	0.321
Risperidone	21 (51.2)	95 (61.3)	0.243
Olanzapine	16 (39)	49 (31.6)	0.370
Lorazepam	12 (29.3)	54 (34.8)	0.502
Dexmedetomidine	1 (2.4)	6 (3.9)	0.660
Diazepam	4 (9.8)	16 (10.3)	0.915
Quetiapine	39 (95.1)	103 (66.5)	< 0.001
Midazolam	4 (9.8)	32 (20.6)	0.109
Distraneurin	4 (9.8)	10 (6.5)	0.465
Melatonin	17 (41.5)	51 (32.9)	0.306
Beta-blocker	1 (2.4)	6 (3.9)	0.660
Pipamperone	2 (4.9)	5 (3.2)	0.612
Levodopa	1 (2.4)	5 (3.2)	0.795
Phenobarbital	1 (2.4)	0 (0)	0.051

Fig. 1


Geographical distribution of non-pharmacological strategies (> or < 50%) by country—blue gradient map. Belarus, Czech Republic, Finland, Iceland, Lithuania, Luxembourg, Malta, Romania, and Slovenia are not represented in the maps because they had fewer than 4 responses



Pharmacological approaches were significantly less often used in countries with a well-established GM (41.4% vs. 70.7%; $p < 0.001$). Among different drugs, trazodone and quetiapine were significantly less used in countries with well-established GM (27.7% vs. 56.1% and 66.5% vs. 95.1%, respectively; all $p < 0.001$). Figure 2 illustrates the geographical distribution of medication usage percentages (the gradient of orange shades reflects the proportion of respondents indicating the use of each medication).

Fig. 2

Geographical distribution of medication usage percentages—orange gradient map. Belarus, Czech Republic, Finland, Iceland, Lithuania, Luxembourg, Malta, Romania, and Slovenia are not represented in the maps because they had fewer than 4 responses



Discussion

The results of this survey highlight significant differences in the diagnosis and management of delirium across European countries, supporting the hypothesis that differences in the level of development of geriatric medicine may influence clinical practice. The presence of a dedicated geriatric ward was significantly more common in well-established GM countries, which likely reflects a stronger institutional focus on geriatric care. This structural difference may contribute to the observed variations in screening and management practices.

A key finding was the higher prevalence of routine delirium screening and established management protocols in well-established GM countries. **AQ3** The use of validated screening tools, particularly in geriatric and orthopedic wards, was significantly more frequent in these countries. This aligns with existing literature, which emphasizes that early identification of delirium is crucial for improving patient outcomes [26]. Screening facilitates timely interventions that can reduce the severity and duration of delirium episodes, potentially decreasing associated morbidity and mortality.

The survey also revealed notable differences in non-pharmacological management strategies. Countries with well-established GM were more likely to implement evidence-based interventions, such as mobilization, hydration support, liberal family visiting policies, and avoidance of potentially deliriogenic medications. These findings are consistent with research indicating that multimodal, non-pharmacological interventions are the most effective strategies for preventing and managing delirium [30,31].

One notable result concerns the use of physical restraints. The survey revealed significant differences between countries, with well-established GM countries demonstrating a significantly lower prevalence of restraint use. Despite strong recommendations against their routine use due to the risk of severe complications, such as increased agitation, functional decline, pressure ulcers, and thromboembolic events [32], physical restraints were still commonly used across all surveyed countries. In particular, their overall utilization was higher than what is typically reported in the literature, which suggests that restraints continue to be a widespread, but suboptimal practice in delirium management [33]. The paradoxical effect of physical restraints, whereby they may worsen delirium symptoms by increasing stress and limiting mobility, has been well-documented [34]. The reduced use of restraints in well-established GM countries suggests greater use of alternative management strategies, such as increased supervision, environmental modifications, and caregiver education. These findings reinforce the need for continued efforts to discourage restraint use and promote evidence-based, patient-centered delirium management.

The observed differences in the prevalence of neuroleptic and sedative use in less-established GM countries indirectly suggest a lower reliance on non-pharmacological approaches, which are more commonly implemented in WEGM countries. However, these findings should be interpreted with caution, as the questionnaire did not distinguish the specific clinical indication for each medication (e.g., delirium treatment, agitation, sleep disturbances, or other conditions) and, therefore, may not fully reflect delirium-specific prescribing practices. This highlights the critical need for targeted education and training in these regions to

discourage pharmacological interventions, except in cases of severely agitated hyperactive delirium, and only after non-pharmacological strategies have been exhausted or when the patient is experiencing significant distress or is at risk of interrupting life-sustaining treatments. Surveys such as this are also valuable for demonstrating the feasibility of alternative approaches in contexts where geriatric care models are less established [29, 34, 35, 36].

These findings underscore the need for standardized European guidelines to harmonize delirium management. The observed discrepancies highlight the potential for improved patient outcomes through wider adoption of structured screening, evidence-based non-pharmacological interventions, and judicious use of pharmacological treatments. **AQ4** Developing consensus-based protocols across Europe could help ensure equitable and high-quality care for older adults at risk of delirium, regardless of geographic location or healthcare system structure.

This study has several limitations. First, the survey captured self-reported information from clinicians regarding practices in their unit or hospital and, therefore, reflects perceived practice rather than externally verified institutional policies. Second, responses were analyzed at the individual level without accounting for potential clustering within countries, as clinical practices may be influenced by national healthcare systems, training structures, and local guidelines. **AQ5** Third, the number of respondents varied substantially across countries, with some countries represented by only a few participants; therefore, country-level findings and visual representations should be interpreted with caution. Fourth, participation was voluntary and distributed through professional networks, introducing the possibility of selection bias; moreover, the survey was conducted in English only, which may also have introduced a selection bias related to language proficiency. Finally, the questionnaire did not specify the clinical indication for pharmacological treatments, limiting the interpretation of these data.

Conclusion

This survey highlights relevant differences in the diagnosis and management of delirium across European countries, reflecting variability in geriatric practices and healthcare organization. Countries with more established geriatric services report more frequent use of routine screening, structured management protocols, and non-pharmacological interventions, together with lower use of physical restraints. **AQ6** In contrast, in settings where geriatric medicine is less developed, delirium care appears more often based on pharmacological or restrictive approaches. Although the observational nature of this study does not allow causal interpretations, the findings suggest the presence of a gap in delirium care across Europe. **AQ7** Query Efforts aimed at greater alignment of diagnostic and management practices may help reduce this gap and promote more consistent delirium care for older hospitalized patients.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Funding

None.

Declarations

Conflict of interest

Alessandro Morandi and Antonio Cherubini are Associate Editors, and Gülistan Bahat is a member of the Editorial Board of *European Geriatric Medicine*; they did not take part in the review or selection process of this article. The authors declare that they have no conflicts of interest to disclose.

Supplementary Information

Below is the link to the electronic supplementary material.

Supplementary file1 (DOCX 15 KB)

Supplementary file2 (PDF 1177 KB)

References

1. American Psychiatric Association, American Psychiatric Association Task Force on DSM-IV (1994) Diagnostic and statistical manual of mental disorders: DSM-IV, 4th edn. American Psychiatric Association, Washington, DC

2. Evensen S, Saltvedt I, Lydersen S, Wyller TB, Taraldsen K, Sletvold O (2019) Delirium motor subtypes and prognosis in hospitalized geriatric patients – a prospective observational study. *J Psychosom Res* 122:24–28
3. Geriatric Medicine Research Collaborative (2023) Increasing frailty is associated with higher prevalence and reduced recognition of delirium in older hospitalised inpatients: results of a multi-centre study. *Eur Geriatr Med* 14(2):325–332
4. Cechinel C, Lenardt MH, Rodrigues JAM, Binotto MA, Aristides MM, Kraus R (2022) Frailty and delirium in hospitalized older adults: a systematic review with meta-analysis. *Rev Lat Am Enfermagem* 17(30):e3687
5. Persico I, Cesari M, Morandi A, Haas J, Mazzola P, Zambon A, Annoni G, Bellelli G (2018) Frailty and delirium in older adults: a systematic review and meta-analysis of the literature. *J Am Geriatr Soc* 66(10):2022–2030. <https://doi.org/10.1111/jgs.15503>
6. Inouye SK, Westendorp RGJ, Saczynski JS (2014) Delirium in elderly people. *Lancet* 383:911–922
7. Bellelli G, Morandi A, Di Santo SG et al (2016) “Delirium Day”: a nationwide point prevalence study of delirium in older hospitalized patients using an easy standardized diagnostic tool. *BMC Med* 14:106
8. Han JH, Wilson A, Ely EW (2010) Delirium in the older emergency department patient: a quiet epidemic. *Emerg Med Clin N Am* 28(3):611–631
9. Fuchs S, Bode L, Ernst J et al (2020) Delirium in elderly patients: prospective prevalence across hospital services. *Gen Hosp Psychiatry* 67:19–25
10. Davis DH, Muniz-Terrera G, Keage HA et al (2017) Association of delirium with cognitive decline in late life: a neuropathologic study of 3 population-based cohort studies. *JAMA Psychiat* 74:244–251
11. Wilson JE, Mart MF, Cunningham C et al (2020) Delirium. *Nat Rev Dis Primers* 6:90
12. Bellelli G, Brathwaite JS, Mazzola P (2021) Delirium: a marker of vulnerability in older people. *Front Aging Neurosci* 30(13):626127
13. American Geriatrics Society expert panel on postoperative delirium in older adults (2015) American geriatrics society abstracted clinical practice guideline for postoperative delirium in older adults. *J Am Geriatr Soc* 63(1):142–150
14. (2023) Delirium: prevention, diagnosis and management in hospital and long-term care. London: National Institute for Health and Care Excellence (NICE)
15. Han JH, Schnelle JF, Ely EW (2014) The relationship between a chief complaint of “altered mental status” and delirium in older emergency department patients. *Acad Emerg Med* 21:937–940
16. Inouye SK, van Dyck CH, Alessi CA, Balkin S, Siegel AP, Horwitz RI (1990) Clarifying confusion: the confusion assessment method. a new method for detection of delirium. *Ann Intern Med* 113:941–948
17. Bellelli G, Morandi A, Davis DH, Mazzola P, Turco R, Gentile S, Ryan T, Cash H, Guerini F, Torpilliesi T et al (2014) Validation of the 4AT, a new instrument for rapid delirium screening: a study in 234 hospitalised older people. *Age Ageing* 43:496–502
18. Zucchelli A, Apuzzo R, Paolillo C, Prestipino V, De Bianchi S, Romanelli G, Padovani A, Marengoni A, Bellelli G (2021) Development and validation of a delirium risk assessment tool in older patients admitted to the emergency department observation unit. *Aging Clin Exp Res* 33(10):2753–2758

19. Golubovic J, Neerland BE, Aune D, Baker FA (2022) Music interventions and delirium in adults: a systematic literature review and meta-analysis. *Brain Sci* 12(5):568
20. Chalfont G, Milligan C, Simpson J (2020) A mixed methods systematic review of multimodal non-pharmacological interventions to improve cognition for people with dementia. *Dement Lond Engl* 19(4):1086–1130
21. Cuevas-Lara C et al (2019) Effectiveness of occupational therapy interventions in acute geriatric wards: a systematic review. *Maturitas* 127:43–50
22. Salvi F, Young J, Lucarelli M, Aquilano A, Luzi R, Dell'Aquila G, Cherubini A (2020) Non-pharmacological approaches in the prevention of delirium. *Eur Geriatr Med* 11(1):71–81
23. Sonia M, Kaur S, Kothari N (2025) Prevention of delirium in the intensive care unit through nonpharmacological intervention: an umbrella review. *Indian J Crit Care Med* 29(1):75–83
24. Marcantonio ER (2011) In the clinic: delirium. *Ann Intern Med* 154(11):ITC6–ITC1
25. Tomlinson EJ, Schnitker LM, Casey PA (2024) Exploring antipsychotic use for delirium management in adults in hospital, sub-acute rehabilitation and aged care settings: a systematic literature review. *Drugs Aging* 41(6):455–486
26. Campbell N, Boustani MA, Ayub A, Fox GC, Munger SL, Ott C, Guzman O, Farber M, Ademuyiwa A, Singh R (2009) Pharmacological management of delirium in hospitalized adults—a systematic evidence review. *J Gen Intern Med* 24(7):848–853
27. Burry L, Mehta S, Perreault MM, Luxenberg JS, Siddiqi N, Hutton B, Fergusson DA, Bell C, Rose L (2018) Antipsychotics for treatment of delirium in hospitalised non-ICU patients. *Cochrane Database Syst Rev* 6:CD005594
28. Soiza RL, Myint PK (2019) The Scottish intercollegiate guidelines network (SIGN) 157: guidelines on risk reduction and management of delirium. *Medicina (Kaunas)* 55(8):491
29. Stuck AE, Masud T (2022) Health care for older adults in Europe: how has it evolved and what are the challenges? *Age Ageing* 51(12):afac287
30. Tahir TA, Eeles E, Karapareddy V, Muthuvelu P, Chapple S, Phillips B, Adyemo T, Farewell D, Bisson JI (2010) A randomized controlled trial of quetiapine versus placebo in the treatment of delirium. *J Psychosom Res* 69(5):485–490
31. Grover S, Mahajan S, Chakrabarti S, Avasthi A (2016) Comparative effectiveness of quetiapine and haloperidol in delirium: a single blind randomized controlled study. *World J Psychiatry* 6(3):365–371
32. Hatta K, Kishi Y, Wada K, Odawara T, Takeuchi T, Shiganami T, Tsuchida K, Oshima Y, Uchimura N, Akaho R, Watanabe A, Taira T, Nishimura K, Hashimoto N, Usui C, Nakamura H (2014) Antipsychotics for delirium in the general hospital setting in consecutive 2453 inpatients: a prospective observational study. *Int J Geriatr Psychiatry* 29(3):253–262
33. Cui N, Qiu R, Zhang Y, Chen D, Zhang H, Rao H, Jin J (2021) Why are physical restraints still in use? a qualitative descriptive study from Chinese critical care clinicians perspectives. *BMJ Open* 11(11):e055073
34. Berger S, Grzonka P, Amacher SA, Hunziker S, Frei AI, Sutter R (2023) Adverse events related to physical restraint use in intensive care units: a review of the literature. *J Intensive Med* 4(3):318–325
35. Briskman I, Dubinski R, Barak Y (2010) Treating delirium in a general hospital: a descriptive study of prescribing patterns and outcomes. *Int Psychogeriatr* 22(2):328–331

36. Yates Z, Lee P, Espat NN, Zagales R, Hernandez N, Amin Q, Ford A, Tweedie C, Elkbuli A (2025) Delirium in critically ill geriatric surgical patients: a systematic review of screening, risk factors, diagnosis, and management. *J Trauma Nurs* 32(4):169–179