

Original Articles

Rooting as indicator of wild boar density: environmental drivers and spatial variation across protected areas

Martina Calosi^a, Niccolò Fattorini^{a,b,*}, Rosa Maria Di Biase^{b,c}, Agnese Marcelli^c, Caterina Pisani^{b,c}, Chiara Gabbrielli^a, Sonia Aleotti^a, Mattia Galdangelo^a, Francesco Ferretti^{a,b}

^a Research Unit of Behavioural Ecology, Ethology and Wildlife Management, Department of Life Sciences, University of Siena, Via P.A. Mattioli 4, 53100 Siena, Italy

^b NBFC, National Biodiversity Future Center, Palermo 90133, Italy

^c Department of Economics and Statistics, University of Siena, Piazza San Francesco 7, 53100 Siena, Italy



ARTICLE INFO

Keywords:

Human-wildlife conflicts
Impact
Mediterranean
Population density
Predictive map
Rooting
Ungulate
Sus scrofa
Wildlife management

ABSTRACT

Wild ungulates play crucial roles in food webs and their impacts can propagate across trophic levels. The recent spread of wild boar *Sus scrofa* is generating concerns worldwide for its potential negative impacts on biodiversity and human activities. Under particular conditions, its foraging activity through digging the topsoil (i.e., rooting) may affect endangered plant/animal species or agriculture. Identifying environmental drivers of rooting is crucial to address spatially-explicit measures to reduce negative impacts, as well as to clarify the often unclear link between rooting and wild boar density. We performed six-year intensive spring-summer surveys (763 sampling plots; 3343 surveys between 2019–2024) in 9 protected areas of central Italy encompassing heterogeneous habitats and a gradient in wild boar densities, to investigate spatial variations in wild boar rooting and relevant drivers. Strong support was obtained for a positive relationship between rooting, at both large (i.e., study area) and fine (i.e., sampling plot) scales, and wild boar density in the area. Results supported lower rooting with increasing landscape diversity. Rooting was affected by topography, being reduced by rock cover and terrain steepness, and increased with distance from the nearest road/railway. Models of fine-scale rooting variation were elaborated to develop high-resolution (10 × 10 m) predictive maps of rooting impact, providing a tool scalable to ecologically comparable areas. Findings support the control of wild boar population density and maintenance of high landscape diversity as measures to reduce wild boar impacts. Finally, the present study also supports that rooting can serve as an effective indicator of within- and between-area variations in wild boar density.

1. Introduction

Wild ungulates have a central role in ecosystem dynamics. They can act as bottom-up restrainers of carnivore abundance as well as top-down regulators of vegetation structure (Hobbs, 1996; Hebblewhite et al., 2005; Ripple et al., 2015; Kaštovská et al., 2024). Their presence may also help to preserve habitat patchwork, e.g., by regulating forest expansion and seed dispersal (Hobbs, 1996; Bueno et al., 2011; Albert et al., 2015; Kaštovská et al., 2024). Nevertheless, under some conditions and especially at high population densities, their activities may become detrimental for the conservation status of particular species, e.g., altering habitat structure and threatening endangered plant and animal taxa (Côté et al., 2004; Barrios-Garcia and Ballari, 2012; Foster

et al., 2014; Barasona et al., 2021). The recent increase in abundance and distribution of wild ungulates across temperate ecosystems has expectedly led to increased interactions with other organisms and human activities. In turn, there is an increasing necessity to identify key indicators that would be able to detect the most at-risk situations and to drive actions for reducing impacts (Caplenor et al., 2017; Pascual-Rico et al., 2021; Carpio et al., 2021).

The wild boar *Sus scrofa* is the most widespread wild ungulate in the world (Barrios-Garcia and Ballari, 2012; Massei et al., 2015). It is characterized by wide plasticity in habitat and food requirements, adapting its diet and space use to local resources and conditions (Herrero et al., 2006; Ballari and Barrios-Garcia 2014; Laguna et al., 2021). This suid is also considered an ecosystem engineer as it can

* Corresponding author at: Research Unit of Behavioural Ecology, Ethology and Wildlife Management, Department of Life Sciences, University of Siena, Via P.A. Mattioli 4, 53100 Siena, Italy.

E-mail address: niccolo.fattorini@unisi.it (N. Fattorini).

<https://doi.org/10.1016/j.ecolind.2025.113806>

Received 18 March 2025; Received in revised form 9 June 2025; Accepted 23 June 2025

Available online 16 July 2025

1470-160X/© 2025 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

modify habitats and biotic communities through its feeding activities (Massei and Genov, 2004; Bueno et al., 2009; Wirthner et al., 2011). The main foraging activity of wild boar is known as “rooting” and consists of using the final part of the snout to dig the soil searching for food (mainly bulbs, rhizomes, and other hypogeous storage organs of plants, but also invertebrates such as insect larvae). Rooting can reach depths of 5–15 cm and can perform superficial tillage over wide areas (Massei and Genov, 2004; Bueno et al., 2009; Barrios-García and Ballari, 2012; Bueno and Jiménez, 2014; Horčíčková et al., 2019). The rooting activity may modify soil structure and moisture (Mohr et al., 2005; Tierney and Cushman, 2006; Bueno et al., 2013), altering nutrient cycles and seed-bank richness (Bueno et al., 2011; Bueno et al., 2013; Palacio et al., 2013), potentially affecting plant and animal species (Barrios-García and Ballari, 2012; Mori et al., 2020; Barasona et al., 2021; Labadessa and Ancillotto, 2023). In ecosystems where the wild boar is native, the activity of this suid should not be considered detrimental *per se* (Sondej and Kwiatkowska-Falinska, 2017; Labadessa and Ancillotto, 2023; Scherer et al., 2025). The negative magnitude of its impact should be scaled considering local characteristics, e.g., the presence of rare/endangered plant and animal species or agricultural activities, and environmental/anthropogenic drivers that could boost rooting. For example, the intensity of rooting may vary according to resource availability, soil conditions, and other environmental characteristics (Welander, 2000; Sütő et al., 2020; Ferretti et al., 2021; Calosi et al., 2024). The occurrence and quality of food resources from the surface to the first soil layers, as well as the proximity of alternative resources such as crops or grasslands, may either attract or divert wild boar attention, locally intensifying or reducing rooting (Welander, 2000; Herrero et al., 2006; Ballari and Barrios-García, 2014; Laguna et al., 2021). Rainfall could boost vegetation growth and create damp soil, which would be richer in undersoil fauna and easier to dig compared to a dry one (Hone 1995; Welander, 2000; Sandom et al., 2013a), generating conditions that favour wild boar sense of smell (Brivio et al., 2017), thus prompting rooting activity (Sandom et al., 2013a; Calosi et al., 2024).

Most importantly, wild boar rooting impact may be exacerbated by population densities (Hone, 2002; Sandom et al., 2013a; Calosi et al., 2024). The relationship between wild boar population density and rooting activity is contradictory, with some studies showing a positive association (Hone, 2002; Sandom et al., 2013a; Calosi et al., 2024) and others failing to support it (Massei et al., 2018; Adams et al., 2019; Ferretti et al., 2021). Disentangling this matter is crucial, considering the recent sharp increase in wild boar densities and distribution, mostly primed by anthropogenic factors (e.g., deliberate local releases and favourable hunting management, anthropogenic foraging opportunities in rural/periurban areas, Massei et al., 2015). The spread of wild boar has been also favoured by the recent climatic trends towards milder winters, and was facilitated by key biological traits such as adaptability and high reproductive potential (Massei et al., 2015; Vetter et al., 2015; Markov et al., 2022). Unravelling the relative role of possible determinants of rooting is a key step forward to address the spatial variation of impacts. Spatially explicit predictions of wild boar impact would be important to identify the appropriate spatial ranges over which management actions should be targeted to mitigate it. Moreover, identifying whether rooting estimates may act as indicators of temporal and/or spatial variations in wild boar population density would be important to support management strategies involving population control to reduce impacts.

Here, we evaluated the effects of wild boar population density, accumulated rainfall, and various environmental factors related to habitat characteristics, on wild boar rooting intensity at two different spatial scales. The ultimate goal was to develop a set of variables suitable to identify the major environmental drivers of rooting at a fine scale, and then to use them to build predictive maps. By conducting six year-intensive surveys in nine protected areas of central Italy, we achieved both fine-scale and large-scale estimates of wild boar rooting impact. Surveys were conducted in early summers, i.e., at the end of the

vegetative season in our areas, to assess and model the environmental drivers of wild boar impact on habitats. Model outputs were used to produce high-resolution predictive maps of wild boar rooting impact to provide novel tools to identify the most at-risk situations and to assist actions aimed at mitigating wild boar impacts.

We expect that (1) rooting intensity, both at the large and fine scales, would be positively related to wild boar density in the area (Calosi et al., 2024). Then, we expect that local environmental features would also shape spatial dynamics of rooting independently from population densities (Bueno et al., 2009; Ferretti et al., 2021). In particular, rooting impact would be boosted by: (2a) milder soil and terrain conditions, i.e., rooting would decrease with increasing rock cover and terrain steepness; (2b) greater amount of accumulated rainfall, as affecting vegetation growth and soil moisture; (2c) habitat homogeneity, because a greater landscape diversity would be expectedly able to provide a wider range of alternative food resources and cover sites, thus reducing the rooting activities; (2d) proximity to persistent wet assets. Finally, we expect that (3) human presence would not reduce wild boar rooting activity, as surveys were conducted in low-populated protected areas, as well as because behavioural flexibility may lead wild boar to shift its temporal activity rather than moving to avoid human interactions (Ohashi et al., 2013; Podgórski et al., 2013; Johann et al., 2020).

2. Materials and methods

2.1. Study areas

The study was performed for six years (2019–2024), from late spring (mid-June) to early summer (July), in nine protected areas of central Italy with extents ranging from 10.5 to 90 km² and an elevation range of 0–1,453 m above sea level (Acquerino-Cantagallo Nature Reserve – ACNR; Alpe della Luna Nature Reserve – ALNR; Alto Merse Nature Reserve – AMNR; Basso Merse Nature Reserve – BMNR; Foresta di Berignone Nature Reserve – FBNR; Foresta di Monterufoli e Caselli Nature Reserve – FMCNR; Maremma Regional Park – MRP; Monte Penna Nature Reserve – MPNR; Sasso di Simone Nature Reserve – SSNR; Fig. 1). Main geographic and environmental features of study areas are summarized in Table 1. All the study areas are located in the Mediterranean bioregion, with generally dry summers and mild winters. Their distribution covers longitudinal and elevational climatic gradients, with the Mediterranean climate gradually acquiring from sub-Mediterranean/temperate traits to continental ones with increasing elevation and distance from the coastline, as approaching the Apennine chain (Pesaresi et al., 2014).

The study areas are mainly covered by broadleaved woodlands (67.8 %, on average), followed by agricultural lands (10.9 %), conifers (5.1 %), shrublands (4.3 %), mixed woodlands (3.7 %), and grasslands (3.5 %) (Fig. 1). Artificial surfaces, bare and rocky substrate, and water bodies cover an average of 3.7 % of the land. Wetlands cover the 0.04 % on average and are present only in MRP. Other ungulates inhabiting the study areas are the roe deer *Capreolus capreolus*, occurring in all study areas, the fallow deer *Dama dama*, in all areas except ACNR and MPNR, and the red deer *Cervus elaphus*, only in ACNR, AMNR, FMCNR, MPNR, and ALNR. Livestock is present locally (cattle: MRP, SSNR; horses: MRP, MPNR, ALNR, AMNR; donkeys: MPNR, ALNR; goats: ALNR; sheep: MPNR, MRP). The wolf *Canis lupus* occurs in all areas and the wild boar has been shown as its major prey (Mori et al., 2016, for a review in Italy; Capitani et al., 2004; Mattioli et al., 2004; Lovari and Sangiuliano, 2006; Ferretti et al., 2019; Lazzeri et al., 2024, for wolf diet in our study region). Recreational hunting is not allowed in any study area, although wild boar may be harvested for population control in some areas (selective culling: FMCNR, FBNR, MRP, SSNR; trapping: MRP) depending on local population density.




Fig. 1. Study area – Left panel: Composition of Land Use (% classes) in each study area; right panel: Location of sampling plots for rooting estimates and land use in each study area: a) Foresta di Monterufoli e Caselli Nature Reserve – FMCNR; b) Foresta di Berignone Nature Reserve – FBNR; c) Acquerino-Cantagallo Nature Reserve – ACNR; d) Sasso di Simone Nature Reserve – SSNR; e) Alpe della Luna Nature Reserve – ALNR; f) Monte Penna Nature Reserve – MPNR; g) Maremma Regional Park – MRP; h) Basso Merse Nature Reserve – BMNR; i) Alto Merse Nature Reserve – AMNR.

Table 1

Summary of geographic and environmental features derived at the study area-scale. Location is given for the centroid of each study area (WGS84). The number of plots includes the number of sampling plots used either for density (all the plots) or rooting (plots in natural/seminatural habitats) estimation. Study areas: Acquerino-Cantagallo Nature Reserve – ACNR; Alpe della Luna Nature Reserve – ALNR; Alto Merse Nature Reserve – AMNR; Basso Merse Nature Reserve – BMNR; Foresta di Berignone Nature Reserve – FBNR; Foresta di Monterufoli e Caselli Nature Reserve – FMCNR; Maremma Regional Park – MRP; Monte Penna Nature Reserve – MPNR; Sasso di Simone Nature Reserve – SSNR.

Study area	Centroid coordinates	Elevation range of the study area (m a.s.l.)	Mean slope (%)	Extent (km ²)	No. plots (Density – Rooting)
ACNR	44.006895°N; 11.054998°E	367–1219	25.4	18.6	60 – 60
ALNR	43.650348°N; 12.166402°E	549–1452	47.0	15.4	48 – 48
AMNR	43.196382°N; 11.202669°E	190–488	14.7	19	65 – 65
BMNR	43.115059°N; 11.333300°E	103–416	11.1	16.4	55 – 50
FBNR	43.331584°N; 10.950866°E	92–554	26.4	22	100 – 100
FMCNR	43.255552°N; 10.783597°E	72–597	30.3	48.3	139 – 136
MRP	42.644144°N; 11.094017°E	0–414	15.2	89	271 – 171
MPNR	42.775285°N; 11.690544°E	484–1104	33.5	10.5	75 – 75
SSNR	43.759573°N; 12.290271°E	692–1207	23.0	16.4	50 – 45

2.2. Rooting and density estimates

Wild boar rooting and population densities were estimated through a design-based, stratified sampling strategy repeatedly described in previous works conducted in the same areas (see Fattorini et al., 2011; Ferretti et al., 2016; Fattorini and Ferretti, 2020; Ferretti et al., 2021; Ferretti et al., 2023, for more details). Sampling plots were circular areas with a 5 m radius, distributed with an effort of ~ 1 plot/0.3 km² in each study area (863 plots, overall). Simulations to assess how the uncertainty of ungulate density estimates varied with the number of sampling plots were previously conducted in our largest study area, supporting the effectiveness of this sampling effort in achieving adequate precision (Fattorini et al., 2011). The density estimation was conducted through faeces counts, using the faecal accumulation rate technique (Mayle et al., 1999). This technique requires visiting each sampling plot twice: a first survey is required for the clearance of previously deposited wild boar faeces; a second survey is conducted after ~ 35–40 days (Ferretti et al., 2016; Fattorini and Ferretti, 2020; Ferretti et al., 2021) to count all wild boar faeces accumulated in plots since the clearance day. An unbiased estimator of faeces abundance and a conservative estimator of its standard error (Fattorini et al., 2011; Fattorini and Ferretti, 2020) were used to estimate the density of wild boar faeces. Wild boar densities were derived by using the wild boar daily defecation rate estimated in summer on a semi-captive Mediterranean wild boar herd localized ~ 40–110 km away from our study areas (6.7 faeces/individual/day; Fattorini and Ferretti, 2020). Although an area-specific defecation rate of wild boar populations would have improved our investigation, the value we used is the only estimate available for Mediterranean populations, in summer, and the one based on the highest number of individuals sampled (Table S4, Supplementary Material).

Rooting intensity was estimated during the clearance surveys,

between mid-June and July, in the same sampling plots used for density estimation. The percentage of plot covered by rooting signs attributable to wild boar was visually estimated and used as an index of rooting intensity (e.g., uprooting of soil and plants and lack of vegetation cover, together with hoofprints and other signs of presence; Adams et al., 2019). Rooting signs may persist for weeks or even months, but we performed rooting estimation after the end of the rainy season, under relatively stable weather at the peak of vegetative growth (late spring-early summer), ensuring that most visible signs reflected relatively recent activity. Visual estimates of impact indices are common in assessments of ungulate impacts (e.g., Felton et al., 2022; Treichler et al., 2023). The same 2–5 trained operators conducted the surveys across study areas and years. At the beginning of each field season, all operators performed joint calibration surveys to harmonize estimation criteria and minimize inter-observer bias. In addition, the operators were supported by preprinted reference images of circular plots covered by simulated percentages of rooted ground, and by reference paper sheets measuring 0.785 m² in size, thus able to quantify 1 % rooting extent in plots (sampling plot size: ~ 78.5 m²). Nonetheless, incorporating formal metrics of inter-observer agreement in future studies could help rule out potential estimation bias. The percentage of rooted area within the circular plots was estimated in classes: 0 %; above 0 % but less than 1 %; then considering integer values: 1–5 %; 6–10 %, 11–15 %; and so on, with a range of 5 % up to the 96–100 % class (Fattorini and Ferretti, 2020; Ferretti et al., 2021). For the statistical analyses, each rooting class was represented by its midpoint value (e.g., 0 % for the 0 % class, 0.5 % for > 0–1 % class, 3 % for 1–5 % class, 8 % for 6–10 % class, 13 % for 11–15 % class, as per the above-defined rooting classes). These midpoint values were used as continuous variables in the models to approximate rooting intensity (e.g., Fattorini and Ferretti, 2020; Ferretti et al., 2021). We considered only those plots located in natural or seminatural habitats (763 plots) to avoid including potential ground alterations by human agricultural activities.

Rooting was estimated on the clearance surveys because our major interest was monitoring this impact during the critical phase of plant growth. In Mediterranean biomes, the peak of plant growth is followed by a season with low rainfall and high temperatures. Disturbances during this time can hinder plant cover recovery, potentially causing long-lasting damage to ecosystems and habitat loss (Belnap, 1995). Although rooting estimates and faeces counts were recorded at different times, the temporal gap between the two surveys was relatively short and constant across years (35–40 days), suggesting relative demographic stability in wild boar populations (i.e., after the birth season, which occurs in spring; Scandura et al., 2022). Considering the short time window, both density and rooting estimates likely refer to the same population and can be meaningfully compared.

2.3. Environmental predictors

For analyses of environmental covariates of wild boar rooting at the fine-scale, rock cover percentage was visually estimated in each plot following the same evaluation criteria used for rooting estimates. Elevation (metres a.s.l.) and terrain slope (%) of sampling plots were obtained from a high-resolution digital elevation model provided by SITA-Regione Toscana (<https://geoblog.regione.toscana.it/-/open-geodata>; resolution: 10 × 10 m), using Qgis 3.38.

For land use classes, the CORINE land cover (CLC) classification was considered by integrating local refinements and fine-scale improvements within the Tuscany region boundaries made by SITA-Regione Toscana (land use and cover – UCS vector: <https://dati.toscana.it/dataset/ucs>; reference year: 2019, after implementing the CLC version 2018). For areas close to the Tuscany region boundaries, the UCS layer was extended for 5 km outside regional limits, merging the UCS with the CLC vector provided by the Copernicus Land Monitoring service (<https://land.copernicus.eu/en/products/corine-land-cover/clc2018>; reference year: 2018). The CLC land use categories were re-classified into 10

broader land-cover classes to enhance ecological interpretability and reduce model complexity, following a hierarchical logic based on the official CLC nomenclature. Specifically, the following classes were considered at the third level of the CLC nomenclature: i) grasslands and meadows, i.e., natural/semi-natural meadows with no scattered trees or bushes; ii) shrublands, i.e., transitional habitats between woodland and open areas such as meadows with scattered trees and/or bushes, or abandoned olive groves; iii) mixed woodlands; iv) woodlands dominated by conifers; v) woodlands dominated by broadleaves. The following classes were grouped according to the first CLC level: vi) agricultural lands, such as cereal crops, vineyards, arable lands, and set-asides; vii) wetlands; viii) water bodies; ix) rock and bare soil; x) artificial surfaces. For the correspondence of the broader classes to the original CLC classes, see Table S2 (Supplementary Material). The ten classes were used to calculate the Shannon diversity index adopted as a proxy of heterogeneity in landscape composition potentially affecting wild boar rooting behaviour. Conversely, only the first seven classes were used to assess the percentage of the area covered by each land use class around the sampling plots, as proxies for the cover of natural/seminatural habitats available to wild boar. Both landscape diversity and cover were calculated in circular buffers of 1100 m radius around the central point of each plot. The buffer size was determined by following Calosi et al. (2024), as the mean of the annual home ranges (weighted by number of individuals sampled) obtained in our largest study area (Massei et al., 1997) and in an area geographically close and ecologically comparable to the study areas (Boitani et al., 1994). As discriminating sex and age class of the individual wild boar that performed rooting is not possible, the smaller home range, i.e., the female one (3.9 km²– radius: 1100 m; males: 7.1 km² radius: 1500 m), was considered as representative for the maximal area likely used by the majority of wild boar (Calosi et al., 2024). This approach included both the focal plot and the surrounding habitats likely influencing the behaviour of the wild boar responsible for the observed rooting.

In line with this, distances from the central point of each plot to the edges of key landscape features were calculated to account for the potential influence of these elements on rooting behaviour. Distances (m) from the central point of each plot to the nearest road/railway, as a proximity index to human disturbance, and the nearest persistent wet asset, as a proxy for water availability, were calculated using the shortest path (point to layer) feature in Qgis 3.38. The former was based on the road and rail network layer and the latter on a new layer created by merging the hydrographic network layer, the water infrastructure layer and inland waters of the above-mentioned UCS layer, each provided by SITA-Regione Toscana and spanning the Tuscany region (road/rail: <http://dati.toscana.it/dataset/grafico-civici>; water sources: <https://www.regione.toscana.it/-/reticolo-idrografico-e-di-gestione>). For areas close to the Tuscany region boundaries, the layers were extended for 5 km beyond Tuscany region boundaries. The road/rail layer was extended by merging the DBPrior10K vector provided by CISIS (Centro Interregionale per I Sistemi Informatici Geografici e Statistici; <http://www.centrointerregionale-gis.it/DBPrior/Dati/DBPriorItalia.zip>), and the water source layer was extended by merging the water infrastructure vector provided by the Italian national geoportal (<https://geodati.gov.it/geoportale/>; Table S3, Supplementary Material). Water sources are an important limiting factor for wild boar (Massei et al., 1997). Ephemeral water bodies such as small ponds and ditches were not georeferenced, possibly leading to underestimating water availability. Nevertheless, the surveys were performed in summer, a season characterized by high aridity in our study region. These conditions do not favour the creation of puddles. Consequently, persistent water resources (e.g., ponds, rivers, and lakes) likely represent the main wet asset available to wild boar during the survey period.

Rainfall has a central role in wild boar ecology (Thurjell et al., 2014; Brivio et al., 2017), especially by influencing environmental parameters such as soil moisture and vegetation growth, which in turn may affect wild boar rooting behaviour (Welandar, 2000; Calosi et al., 2024). Thus,

cumulative amounts of rainfall (millimetres) were evaluated at three different temporal scales to investigate either the direct effect on rooting impact by moisturizing the soil (7 days) or the indirect effect on rooting through enhanced vegetation growth (15–30 days). The maximum temporal window was selected based on a global analysis of the effects of accumulated rainfall on vegetation growth, while considering the vegetation types present in the study areas (~30 days; Ding et al., 2020). Rainfall data were gathered by the professional weather stations (ARPAE Emilia Romagna, for ALNR: <https://simc.arpae.it/dext3r>; Servizio Idrologico Regione Toscana, for all the others: <https://www.sir.toscana.it>) closest to the centroid of each study area (Table S1, Supplementary Material). Rainfall data were downloaded and elaborated, obtaining 7, 15, and 30 days-cumulative rainfall before the survey date (rooting estimation) of each sampling plot.

Together with the distance from the nearest road/railway, the human population density was also considered as a proxy for the degree of human presence. The Global Human Settlement raster provided by the Copernicus Emergency Management Service, indicating the number of residential people per raster cell (Schiavina et al., 2023; resolution: 100 × 100 m) was used to obtain the average number of people living in the 1100 m radius circular buffer around each plot.

2.4. Statistical analyses

Statistical analyses were performed through generalized linear mixed models (GLMMs; Zuur et al., 2009). Separate analyses were performed for rooting estimates at the (a) large- (study area) and (b) fine- (sampling plot) scales.

- (a) At the study area-scale, the response variable was the probabilistic rooting estimate for each study area and year resulting from the plot-level estimates. This value was then converted into a proportion and was modelled against the estimate of wild boar population density in the same area (fixed effect; continuous, as number of individuals/km²). Rooting intensity was modelled through beta errors (link function: logit), as recommended for continuous proportions bounded through the 0–1 range (ordered beta parametrization: Kubinec, 2023). Since the response variable represents an estimate with associated uncertainty, it was weighed by the inverse of the standard error of the rooting proportion estimate, in turn accounting for the precision of rooting estimates. As the number of data-points is given by 39 rooting estimates, the model would not allow the inclusion of further predictors. Considering that the surveys were repeated within the same study years and areas, the variables ‘Year’ and ‘Study area’ were included as random intercepts. This model specification, where the study area is treated as a random intercept, accounts for both the within- and between-area effects. Thus, such model would provide the effect of wild boar population density on rooting independently of the study area, meaning that, if a positive density effect on rooting is found, density can predict rooting both within a study area (i.e., between-years in the same area) and between different study areas (i.e., on an ‘average’ year). To further assess this possibility, the model was re-fitted through a mathematical specification able to distinguish within- versus between-area effects (‘within-group centring’; Van de Pol and Wright, 2009). Thus, wild boar population density was centred on its study area-mean and used as a predictor, and the mean population density in each study area was also added as a further predictor. The other model components were kept identical.
- (b) At the sampling plot-scale, three global models were run separately, one for each period over which cumulative rainfall was considered (7, 15, and 30 days-cumulative rainfall). This approach allowed testing direct and indirect effects of rainfall on rooting intensity, which could not be handled together in the same model due to high collinearity (7 days vs 15 days: $r = 0.85$;

7 days vs 30 days: $r = 0.73$; 15 days vs 30 days: $r = 0.77$). Rooting intensity estimated in each sampling plot (response variable) was converted into a proportion and modelled through beta errors. The ultimate goals were to identify the major environmental drivers of rooting at a fine scale, and then to use this set of drivers to build predictive impact maps. Thus, collinearity among variables was checked to avoid potentially strong correlations among local environmental characteristics. Only variables showing a correlation of $|r| < 0.4$ with others were retained as fixed effects (Fig. S1, Supplementary Material). As a result, shrubland cover was excluded due to collinearity with landscape diversity and distance from persistent wet assets; elevation was excluded due to collinearity with grasslands cover; broadleaves woodland cover was excluded as it represented the most common habitat type among the study areas and showed collinearity with agricultural lands, conifer woodland cover and landscape diversity (Fig. S1, Supplementary Material). Additionally, wetland cover was excluded as present only in MRP. Therefore, the fixed effects included in the global models were: (1) wild boar population density (continuous, as number of individuals/km²); (2) 7, 15 or 30 days-cumulative rainfall before the survey date (continuous, in millimetres); (3) rock cover (continuous, as % in each plot); (4) distance from the nearest persistent wet asset (continuous, in metres); (5) distance from the nearest road/railway (continuous, in metres); (6) slope (continuous, as %); cover of: (7) mixed woodlands, (8) grasslands and meadows, (9) agricultural lands, and (10) conifers within the buffer (predictors 7 to 10: continuous, as % land class cover in 1100 m circular buffers); (11) landscape diversity within the buffer (continuous, as Shannon index in 1100 m circular buffers); (12) human population density within the buffer (continuous, as average number of people living in each buffer). To account for repeated rooting estimates obtained within the same study year, study area, and sampling plot, ‘Year’, ‘Study area’, and ‘ID plot’ were included as random intercepts. In each global model, there was no multicollinearity among explanatory variables (all VIFs ≤ 1.8) and covariates were scaled to increase coefficients’ interpretability and model convergence, as well as to compare effect sizes of different predictors.

For each global model in both (a) and (b), an all-subset model selection was performed, as each combination of predictors could represent a different *a priori* hypothesis that could not be discarded in advance (Harrison et al., 2018). Thus, all possible combinations among predictors, as well as the null model (i.e., the model with only the random intercepts), were ranked and weighted from each global model (in *a*, only the null model was the possible competing model). Ranking was based on the AICc value of each model and the relevant Δ AICc, i.e. its difference with the model with the lowest AICc value. Models were selected if they showed Δ AICc < 2 and, following the “nesting rule” (Harrison et al., 2018), when they were not more complex than any simpler alternative with a lower AICc value. Thus, either a single top-ranked model (Table S5, Supplementary Material) or a set of top-ranked competing models (Table S6, Supplementary Material) was obtained from the model selection procedure, depending on the analysis. Model weight was standardised within the subset of selected models. Coefficients of predictors and 95 % confidence intervals were estimated from the best model. The effects of predictors were assessed by checking whether 95 % confidence intervals of coefficients overlapped ‘0’. Best models were validated by visually checking residual patterns (Zuur et al., 2009). GLMMs and model selection were run respectively through the packages *glmmTMB* (Brooks et al., 2017) and *MuMIn* (Bartoń, 2023), using the R software version 4.4.0 (R Core Team, 2023).

2.5. Predictive maps of wild boar rooting

The second aim of this study was to map the wild boar rooting impact continuously across space in the study areas, to identify the most-at-risk sites. A model-assisted approach based on predictive variables available for any location of the study areas was adopted. The only predictive variable visually estimated *in situ*, and as such not available wall-to-wall, was the rock cover. Therefore, all the global models and model selections adopted in the study for assessing drivers of rooting (see [section 2.4](#)) were re-fitted without considering rock cover among predictors. Although rock cover was a driver of rooting intensity (see [section 3.2](#)), its exclusion did not affect model selection, which supported the same best model, only differing for the lack of rock cover (cf. [Table S6-S7, Supplementary Material](#)). Moreover, the best models fitted with and without rock cover provided identical predictive capacity (Pearson correlation between observed and predicted values, for both models: $r = 0.64$; $R^2 = 0.42$), and predicted values from the two models were almost perfectly positively correlated ($r = 0.99$; $R^2 = 0.98$). Thus, excluding rock cover from predictors did not qualitatively affect the predictive capacity of the maps. Maps were built at the finest possible resolution considering constraints in computational effort, but attempting to achieve the closest possible resolution to the original data extent. Thus, raster cell (i.e., pixel) resolution was set at 10×10 m, i.e., 100 m^2 . To the best of our knowledge, there is no map concerning wild boar impact available at such a fine resolution. The rooting impact was predicted for each study area and year separately. Because the use of the plot identity as a random intercept would have precluded the interpolation in raster cells outside the sampling plots, predictor coefficients were estimated from the best model without including this random intercept (mapping model, [Table S8, Supplementary Material](#)).

At each raster cell, the area-specific raster layers of predictor variables in the mapping model were used, exploiting its mathematical equation (Equation 1, [Supplementary Material](#)). The model intercept was adjusted for each area and year by considering the random intercept values estimated by the mapping model in the same study area (Equation 1, [Supplementary Material](#)), as well as by holding the area-specific value of wild boar population density at its mean value, calculated across the study years ([Fig. S2, Supplementary Material](#)).

Once predictions at each raster cell were achieved for each study area and year by exploiting the mapping model, the residuals were calculated for each sampled location (i.e., sampling plot), where both predicted and observed values were available. We followed the novel approach by [Fattorini et al. \(2024\)](#) to spatially interpolate the residuals at non-sampled locations by the inverse distance weighting interpolation, with the smoothing parameter selected from sample data by a leave-one-out technique. Finally, the interpolated residuals were added to the model predictions to achieve the mapped values. Contrary to geostatistical techniques that necessitate several more or less realistic assumptions, this mapping strategy is design-based, avoiding any reliance on model assumptions. Also, the statistical properties of the maps are objective, as they stem from the sampling scheme actually adopted in the field. In particular, under tessellation stratified sampling, the resulting maps have been proven consistent, i.e., they converge to the true maps as the number of sampling locations increases ([Fattorini et al., 2024](#)).

After obtaining area- and year-specific maps, since rooting was estimated on natural/seminatural habitats, portions of the study areas covered by water bodies, anthropic surfaces, and agricultural lands were cut out.

3. Results

3.1. The relationship between rooting and density across protected areas

Estimates of wild boar population density (39 estimates) varied from 1.7 to 22.4 individuals/ km^2 , and those of rooting impact at the study

area scale from 0.6 % to 12.9 % of rooted area (for study year and study area-specific estimates, see [Fig. S2, Supplementary Material](#)). The model including the wild boar population density as predictor was selected over the null model ([Table S5, Supplementary Material](#)), supporting a positive relationship between wild boar density and rooting, and showing more than a two-fold increase of rooting impact when wild boar population density increased from 1.7 to 22.2 individuals/ km^2 ([Fig. 2a; Table 2a](#)). The same model re-fitted to distinguish within- (i.e., centred density) versus between-area (i.e., average density) effects provided identical results, confirming that the positive effect of wild boar population density on rooting occurred both within the same area (thus, between years) and between different study areas, with the former effect being stronger ([Table S9, Supplementary Material](#)).

3.2. Environmental drivers of fine-scale rooting estimates

At the plot sampling-scale, the percentage of ground with rooting varied from 0 to 93 % (3343 plot-surveys), with 36.0 % of the estimates yielding no rooting impact, 50 % of the estimates showing rooting intensities above 0 % but less than 10 %, and 14 % of the estimates showing rooting impact above 10 %. Two top-ranked models were selected to explain factors influencing wild boar rooting at the plot-scale for each cumulative rainfall temporal window ([Table S6](#)). The best model was the same for all the three analyses, supporting the effects of (1) wild boar population density, (2) rock cover, (3) distance from the nearest persistent wet asset, (4) distance from the nearest road/railway, (5) slope, (6) the % of mixed woodlands, (7) the % of agricultural lands, and (8) landscape diversity. Rooting increased with increasing wild boar population density, which had the strongest effect among predictors, indicating more than a twofold increase when density increased from 1.7 to 22.4 individuals/ km^2 ([Fig. 2b; Table 2b](#)). In addition, rooting impact decreased with increasing rock cover within the plot ([Table 2b; Fig. 3a](#)), terrain steepness within the plot ([Table 2b; Fig. 3b](#)), landscape diversity within the buffer ([Table 2b; Fig. 3c](#)), and distance from the nearest road/railway ([Table 2b; Fig. 3d](#)), although the latter showed a weak effect size. The percentage of mixed forests and that of agricultural lands within the buffer, as well as the distance from the nearest persistent wet asset, were selected in the best model, but confidence intervals overlapped “0”, indicating them as uninformative predictors ([Table 2b](#)).

3.3. Map of spatial variations in rooting activity

Area-specific, spatially-explicit maps of wild boar rooting impact were obtained by averaging year-specific maps for each study area ([Fig. 4](#)). Year-specific spatially-explicit maps are depicted in [Supplementary Material \(Fig. S3 and S4\)](#).

4. Discussion

The recent and ongoing spread of wild boar across temperate ecosystems is raising concerns on the potential negative impacts of this suid on natural habitats as well as human activities ([Massei and Genov, 2004; Barrios-Garcia and Ballari, 2012](#)). Identifying determinants of wild boar impact on habitats is crucial to define adequate mitigation measures. Our results provided strong support for the roles of wild boar population density, landscape heterogeneity, and topography in influencing rooting activity. These findings can be used to obtain site-specific information driving spatially-explicit management actions.

4.1. Wild boar population density: a critical driver of rooting pressure

The results supported a positive relationship between wild boar population density and rooting, both at the study area and sampling plot scales (prediction 1). Data were based on six-year intensive sampling, during which rooting and density estimates were collected within the same sampling plots. This methodological consistency likely




Fig. 2. Wild boar population density effect – Relationships between rooting (a) at the study area-scale and (b) at the sampling plot-scale and wild boar population density in the area, as estimated by GLMMs. Marks along the x-axis show the distribution of observed values for each covariate. Lines: predicted values. Bands: 95% confidence intervals.

Table 2

Parameters estimated from top-ranked GLMMs predicting wild boar rooting (a) at the study area-scale and (b) at the sampling plot-scale: coefficients (β) and 95% confidence intervals (95% CIs). An asterisk marks the coefficients whose 95% CIs do not include 0. Variance of random intercepts in the model (σ^2) is also shown.

Response variable	Predictors	β	95 % CIs
(a) Proportion of rooting at area-scale $\sigma_{\text{Area}}^2 = 0.574$ $\sigma_{\text{Year}}^2 = 0.220$	(Intercept)	-3.392	-3.806, -2.978*
	Wild boar population density	0.261	0.246, 0.275*
(b) Proportion of rooting at plot-scale $\sigma_{\text{Area}}^2 = 0.348$ $\sigma_{\text{Year}}^2 = 0.154$ $\sigma_{\text{ID Plot}}^2 = 0.554$	(Intercept)	-2.999	-3.273, -2.724*
	% Mixed woodlands	0.061	-0.003, 0.125
	% Agricultural lands	0.056	-0.016, 0.128
	Wild boar population density	0.190	0.130, 0.250*
	Nearest wet asset distance	-0.056	-0.130, 0.018
	Nearest road/railways distance	-0.099	-0.175, -0.022*
	Landscape diversity	-0.181	-0.282, -0.080*
	Slope	-0.125	-0.192, -0.059*
Rock cover	-0.226	-0.304, -0.148*	

strengthened the observed relationship, which otherwise might have appeared weaker using shorter time series or smaller samples of study areas (Nakagawa and Cutchill, 2007; e.g., Massei et al., 2018; Adams et al., 2019; Ferretti et al., 2021). The wild boar is a gregarious species and higher population densities often lead to larger aggregations of individuals (Massei et al., 1997; Barrios-Garcia and Ballari, 2012). Local clustering of individuals can amplify the effects of behaviours such as rooting and trampling, intensifying impacts on soil and vegetation (Massei et al., 1997; Hone et al., 2002; Barrios-Garcia and Ballari, 2012). The magnitude of these impacts, as well as thresholds of population density beyond which impacts may become threatening for ecosystems, would be expected to depend on local environmental carrying capacity, which may vary across seasons and years (Massei et al., 1997; Barrios-Garcia and Ballari, 2012). Based on our results, moderate-to-high wild boar population densities (~15–20 ind./km² in summer, i.e., after births) were associated with significant rooting activity (~5–8 % of rooted area), which may pose a threat to vulnerable habitats. The occurrence of detrimental effects should depend on the local ecological context, as well as on the local abundance and distribution of focal taxa affected by rooting (e.g., orchids or other vulnerable plants; threatened birds, smaller vertebrates or invertebrates). The effect of population density could be exacerbated by environmental factors that favour rooting, such as abundant food resources underground or specific habitat characteristics (Baubet et al., 2004; Bueno et al., 2009; Lombardini et al., 2017). Nevertheless, density-independent negative effects of rooting should not be ruled out. For instance, even a small number of

individuals may exert negative impacts on rare and localized taxa (e.g., orchid species) and their habitats. In addition, unmeasured variables such as predator pressure and population structure may mediate the observed relationship between density and rooting. These factors could influence wild boar population dynamics and behaviour (Lingle, 2001; Scandura et al., 2022), locally altering the rooting intensity. For example, predation pressure may exert a top-down control on prey density, while increased predation risk may lead to larger group sizes in ungulates (Lingle, 2001), thus modifying the spatiotemporal distribution and magnitude of their impacts. Individual traits such as sex, age, and physiological stage may also affect foraging behaviour. Males typically live solitary or in small groups whereas sows tend to aggregate in larger family groups, particularly while weaning their offspring (Scandura et al., 2022). Also, time spent foraging differs with age: piglets may allocate about twice the time to foraging activities more than adults (Spitz and Janeau, 1995). Consequently, females with piglets may exert a cumulatively greater rooting impact than males, due to both greater time spent foraging and larger group size typical of maternal groups (Spitz and Janeau, 1995; Scandura et al., 2022). Future studies should investigate whether such unaccounted demographic and behavioural factors mediate the observed density-effect on rooting impact. Furthermore, an increased sampling effort in areas with low wild boar density and rooting pressure, where the number of '0 counts' increases, thus inflating the variance of estimates (Fattorini and Ferretti, 2020), may improve the investigation of the relationship between density and rooting. Eventually, we acknowledge that our density estimates




Fig. 3. Environmental predictors effects — Relationships between rooting at the sampling plot-scale and (a) rock cover, (b) terrain steepness, (c) landscape diversity, and (d) distance from the nearest road/railway, as estimated by GLMMs. Marks along the x-axis show the distribution of observed values for each covariate. Lines: predicted values. Bands: 95% confidence intervals. Confidence intervals for all the above predictors do not overlap zero.

rely on defecation rates derived from a semi-captive population (Fattorini and Ferretti, 2020), which might not fully reflect those in wild populations. Although the use of defecation rates estimated under captive or semi-captive conditions is a common approach to monitor ungulate density through faeces counts (Adams et al., 2019; Plhal et al., 2014), future studies could enhance the accuracy of our density estimates by directly obtaining defecation rates from wild individuals (e.g., through field tracking or isotopic labelling), thereby strengthening the observed rooting-density relationship.

4.2. Environmental drivers of rooting pressure

Rooting decreased with increasing rock cover and terrain steepness, confirming prediction (2a) (Ferretti et al., 2021; Calosi et al., 2024). Rocky terrains and steep slopes can enhance water percolation, which in turn reduces soil moisture (Penna et al., 2009; Lange et al., 2010). This may harden the first layers of soil, locally hampering wild boar digging activity (Sandom et al., 2013a).

Accumulated rainfall was not selected in the best model, not supporting an effect of rainfall on rooting activity, contrary to prediction (2b). Rainfall can affect wild boar ecology (e.g., Fernandez-Llario and Mateos-Quesada, 2005; Thurjfell et al., 2014; Colomer et al., 2024). Previous analyses conducted on a single habitat type and throughout the year, thus covering a gradient from rainy to drier periods, showed that rooting activity was favoured by precipitation (Calosi et al., 2024). The

weather homogeneity within a single, relatively dry season, such as late spring-early summer, may have limited the potential of detecting the effects of rainfall on rooting activity. At the multi-habitat scale of our investigation, wild boar dependence on rain-mediated food resources could be buffered by the presence of habitats rich in food resources year-round (e.g., woodland for understorey resources or agricultural land for herbaceous supply) which could be selected in arid periods, mitigating the effect of rainfall (Herrero et al., 2006; Keuling et al., 2009; Ruf et al., 2023). Future analyses incorporating rainfall and soil moisture data at a finer spatial scale could help reveal relevant effects that may have been underestimated at the broader scale of our investigation.

Results showed weak (i.e., confidence intervals overlapping '0') support for the effects of specific land use classes on rooting, such as percentages of agricultural land and woodland within wild boar home range. They supported higher rooting activity in areas with lower landscape diversity. Those results partially confirmed prediction (2c), suggesting that the wild boar is not strictly tied to specific land uses but rather responds to overall landscape composition and specific local conditions (Briedermann, 1990; Morelle et al., 2016; Fettebert et al., 2017; Yang et al., 2024). Habitat composition shapes wild boar ranging patterns (Keuling et al., 2009; Fettebert et al., 2017; Yang et al., 2024). Ecological plasticity enables wild boar to adjust their movements by selecting cost-effective food resources and suitable resting sites (Keuling et al., 2009; Morelle et al., 2016; Fettebert et al., 2017; Scandura et al., 2022). Heterogeneous landscapes are expected to provide abundant and




Fig. 4. Predictive maps of impact – Predictive maps of wild boar rooting obtained by averaging year-specific maps within each study area. Rooting estimates (%) are displayed in 1 % classes up to > 10 % for a more immediate interpretation. **Study areas:** NR – Nature Reserve; RP – Regional Park; a) Acquerino-Cantagallo NR; b) Alto Merse NR; c) Alpe della Luna NR; d) Basso Merse NR; e) Foresta di Berignone NR; f) Maremma RP; g) Monte Penna NR; h) Foresta di Monterufoli e Caselli NR; i) Sasso di Simone NR.

diverse food resources to wild animals, thereby influencing both spatial behaviour and foraging strategies. For example, the availability of heterogeneous resources at close range may reduce the need for animals to range over large areas (Ward and Saltz, 1994; Mangipanea et al. 2018). The role of greater landscape heterogeneity and presence of ecotones in

reducing individual movements (e.g., Lucherini and Lovari, 1996; Säid and Servanty, 2005; Lovari et al., 2013) and in promoting dietary diversity (Minder, 2011; Mangipanea et al., 2018) has been reported for several mammalian species. In wild boar, these patterns may reflect two complementary mechanisms: (1) movement constraints, whereby

increased local resource availability may reduce the need for extensive ranging and intensive rooting to meet nutritional demands; and (2) dietary shifts, as individuals may rely on a broader range of easily accessible food resources, thus reducing the reliance in underground items obtained through rooting (Massey et al., 1997; Schley and Roper, 2003; Scandura et al., 2022). These findings emphasize the importance of accounting for habitat heterogeneity when evaluating rooting dynamics, as local environmental conditions can significantly influence the extent of rooting behaviour. Additionally, seasonal patterns may alter the observed relationships. Previous research has reported different rooting intensity driven by seasonal shifts in food availability and soil conditions (e.g., Massey et al., 1997; Welander, 2000; Calosi et al., 2024). For instance, high mast availability in autumn may mitigate rooting activity, thereby reducing the impact on ground vegetation independently of the surrounding landscape mosaic. Conversely, increased precipitation during the same season may promote rooting by enhancing wild boar olfactory capabilities and making the soil easier to dig (Welander, 2000; Sandom et al., 2013a). Future research comparing seasonal patterns of rooting activity across climatic gradients could help disentangle the relative contributions of local and seasonal effects.

Contrary to prediction (2d), water availability appeared to constrain rooting activity weakly, as the confidence intervals of the estimates overlapped zero. Wild boar select habitats along water edges throughout the year for wallowing in mud and resting (Thurfjell et al., 2009). These habitats are particularly crucial in summer, as mud-bathing helps with thermoregulation (Bracke, 2011; Olczak et al., 2015; Ruf et al., 2023). The results suggest that wild boar may fulfil its foraging needs independently of water availability, possibly by accessing wet areas separately from feeding activities. Notably, the model excluding rock cover (i.e., the one used for producing the impact maps), supported increased rooting in woodland-dominated landscapes and approaching the closest persistent wet assets. These results suggest that rock cover interacts with other variables, potentially masking their effects, and support a role of woodland and water availability in shaping rooting behaviour in rock-free terrains.

4.3. Human activities and rooting pressure

Results supported that rooting activity was not influenced by human population density within the wild boar home range, and rooting activity increased closer to the nearest road or railway – albeit with a relatively weak effect size. This confirms prediction (3), suggesting that wild boar may take advantage of human-made features such as forest roads for cost-effective, easier, and faster movements between resting and foraging sites (Thurfjell et al., 2015). The study areas are all protected, with very low human population density (~0.1–3.8 inhabitants/km²; data from Schiavina et al., 2023), while roads predominantly follow forest tracks, and mostly have restricted vehicle access. As a result, vehicle traffic is minimal, and roads could be perceived as safe corridors that facilitate movement, particularly during nighttime when people are absent (Keuling et al., 2009; Thurfjell et al., 2015; Podgórski et al., 2013). Previous studies showed that human activities such as intensive agricultural and recreational practices (e.g., hiking or hunting) can alter wild boar behaviour and habitat use (Podgórski et al., 2013; Brivio et al., 2017; Brogi et al., 2020). However, wild boar behavioural flexibility could allow this suid to shift its temporal activity patterns in response to human pressures (Keuling et al., 2009; Podgórski et al., 2013; Gordigiani et al., 2022), for example by foraging closely to roads during the night. Future research should investigate wild boar activity patterns in relation to proximity to human settlements to shed light on this possibility.

4.4. Assessing and mapping rooting pressure on habitats

Results suggest that the distribution and intensity of rooting activity were context-dependent and influenced by multiple factors across

protected areas. Rooting activity should not be interpreted as a stand-alone negative indicator of habitat alteration. Moderate levels of rooting may generate positive effects such as soil aeration, control of the spread of pioneer species, and maintenance of open habitats (Groot Bruinderink and Hazebroek, 1996; Barrios-Garcia and Ballari, 2012; Barrios-Garcia et al., 2023). Rather, negative impacts may arise when rooting is both intense and frequent, particularly in combination with specific environmental conditions (Burrascano et al., 2015). In the short term, it may cause a decline in focal plant species of the impacted habitat, facilitating the establishment of ruderal or nitrophilous species. When the alteration persists over time, it could lead to shifts in plant species composition (Burrascano et al., 2015), to the extinction of habitat-specific focal species (Sandom et al., 2013b; Burrascano et al., 2015), ultimately resulting in a reduction of the habitat surface (e.g., Bratton, 1974; Barrios-Garcia and Ballari, 2012). A reduction or alteration of 1 % of surface in rare or declining habitats has been considered as a threshold for identifying significant pressures for conservation, deserving appropriate mitigation measures (Case C-258/11, European Union Court of Justice, 2013). For example, repeated rooting by high wild boar densities may threaten habitats such as some grassland types, particularly during the growing season, potentially leading to a significant reduction in ground vegetation cover (Calosi et al., 2024, for EU priority grasslands, codes '6210' and '6220'). If these changes affect characteristic species, the local conservation status of that habitat might be at risk, potentially leading to its extinction when already rare or fragmented. Suggestively, preliminary surveys conducted in priority-protected grasslands of one of our study areas (MPNR), showed that vegetation cover was much lower in rooted compared to unrooted patches, and species composition in rooted patches shifted toward ruderal, low-quality taxa, not typical of the same grassland habitat (our unpublished data). In the long-term, such a shift in plant species composition may trigger cascading effects on animal communities, particularly those species that depend on specific host plants for reproduction and survival (e.g., insects; Howe et al., 1981; Singer et al., 1984; Scandurra et al., 2016). In this context, monitoring vegetation shifts in response to rooting is crucial to better understand these potential effects and inform conservation strategies.

To the best of our knowledge, this is the first study producing spatially-explicit maps of wild boar rooting impact on natural and seminatural habitats in protected areas (see e.g. Yang et al., 2024, for risk maps of wild boar crop damage). This approach can be calibrated over other protected areas with comparable land use composition and climatic conditions, in combination with local wild boar population density estimates, to identify the most endangered sites within areas. Maps can be generated using widely available land cover data (e.g. CLC; Copernicus, 2023), digital elevation models, and by exploiting the model proposed in this study at least for ecologically comparable areas (Table S8, Supplementary Material) or by adapting it to local environmental conditions (i.e., integrating relevant site-specific predictors). Our approach could be further improved by validating the maps through independent rooting surveys, which were not available in our study. Such limitation could be addressed in future research by combining complementary methodologies, such as drone-based aerial surveys or systematic ground measurements, enhancing the accuracy of our mapping approach.

Maps would provide conservationists, practitioners, and stakeholders with a tool to identify spatial variations in rooting pressure, helping prioritize conservation efforts in sites that host threatened plant or animal species or that are particularly sensitive to damage (Yang et al., 2024). In turn, evidence-based management and conservation strategies would be supported to mitigating wild boar negative impacts on ecosystems. Nonetheless, we recommend caution in generalising the transferability of our specific findings, which were obtained in protected areas, i.e., areas sparsely populated by humans, encompassing a mosaic of natural and semi-natural habitats dominated by woodland, within the Mediterranean bioregion along an elevation-dependent climate

gradient. Consistency with other ecological contexts, such as human-dominated areas and different bioregions or seasons, remains to be evaluated.

4.5. Conclusions and implications

Our study provides insights into the environmental drivers of wild boar rooting during late spring and early summer in Mediterranean protected areas. First, our findings support that rooting at both local and broader spatial scales can be used as an effective indicator of changes in wild boar density. This result suggests that estimates of rooting impact can serve as a reliable and cost-effective proxy for evaluating the within-area, between-years variation in wild boar densities, if standardised and conducted in the same season. This finding also supports management strategies aimed at controlling population densities to mitigate wild boar impacts on habitats. Yet, the observed density–impact relationship was conditional on the two spatial scales of rooting evaluated (Table 2), calling for future investigations conducted along a scale gradient. Second, results emphasise the role of landscape diversity as a potential buffer against wild boar impact through rooting. Consequently, strategies that promote or restore ecosystem diversity could serve as an indirect approach to reducing wild boar pressure on habitats. Third, we provide a framework to produce predictive maps potentially helpful to identify the most at-risk locations in natural/semi-natural habitats. In turn, targeted management actions could be planned. These may involve site-specific control of wild boar density, preventive measures based on small fences to prevent wild boar rooting around focal plants or plant groups where feasible, or through specific dissuasive actions in sites and periods of higher threat to particular habitats.

Fundings

This multi-year study has been supported by Regione Toscana – Settore Tutela della Natura e del Mare; Ente Parco Regionale della Maremma; Tuscany Environment Foundation, under the MAPAT project; National Biodiversity Future Center - NBFC, project funded under the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.4 - Call for tender No. 3138 of 16 December 2021, rectified by Decree n.3175 of 18 December 2021 of Italian Ministry of University and Research funded by the European Union – NextGenerationEU, Award Number: Project code CN_00000033, Concession Decree No. 1034 of 17 June 2022 adopted by the Italian Ministry of University and Research, CUP B63C22000650007, Project title “National Biodiversity Future Center – NBFC; project PRIN 2022 – PNRR n. P20228RERY, funded by the Italian Ministry of University and Research under the PNRR, CUP B53D23023510001- Codice Progetto Prot. P20228RERY_001 Mission 4 Component 2 (M4C2) – Investment 1.1: Fondo per il Programma Nazionale Ricerca (PNR) e progetti di Ricerca di Significativo Interesse Nazionale (PRIN) del Piano Nazionale di Ripresa e Resilienza (PNRR) funded by the European Union “Next Generation EU”, 2022. MC was supported through a fellowship funded by Regione Toscana – FSC – Fondo per lo Sviluppo e la Coesione – POR FSE 2014-2020 – GiovaniSi.

CRedit authorship contribution statement

Martina Calosi: Writing – review & editing, Writing – original draft, Investigation, Data curation, Formal analysis, Conceptualization. **Niccolò Fattorini:** Writing – review & editing, Formal analysis, Data curation, Conceptualization, Investigation. **Rosa Maria Di Biase:** Writing – review & editing, Software, Formal analysis. **Agnese Marcelli:** Writing – review & editing, Software, Formal analysis. **Caterina Pisani:** Software, Writing – review & editing, Formal analysis. **Chiara Gabbriellini:** Investigation. **Sonia Aleotti:** Investigation. **Mattia Galdangelo:** Investigation. **Francesco Ferretti:** Supervision, Data curation, Writing – review & editing, Formal analysis, Conceptualization.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

We are indebted to G. Ruberti, S. Rusci, L. Venturi, and E. Giunta, for their support to our study. Financial support was provided by Regione Toscana—Settore Tutela della Natura e del Mare (RTSTNM), by the Maremma Regional Park (MRP) and by the Tuscany Environment Foundation; we are grateful to these Agencies, to Ente Terre Regionali Toscane, and to the many landowners who allowed us to conduct fieldwork. NF, RDB, CP, and FF acknowledge the support of NBFC to Univ of Siena/Department of Life Sciences and Department of Economics and Statistics, funded by the Italian Ministry of University and Research, PNRR, Missione 4 Componente 2, “Dalla ricerca all’impresa”, Investimento 1.4, Project CN00000033. MC was supported through a fellowship funded by Regione Toscana – FSC – Fondo per lo Sviluppo e la Coesione – POR FSE 2014-2020 – GiovaniSi. Special thanks go to D. Nonis, L. Tonini, and to all the staff of RTSTNM and MRP, for their logistic and technical support. We are grateful to all the students who participated in field surveys, to O. Tomassini and G. Bonari, who provided critical comments on an early draft, and to C. Angiolini and L. De Simone for sharing unpublished preliminary results on the effects of rooting on plant communities and comments on the relevant text. We thank Servizio Idrologico Regione Toscana and Arpa Emilia-Romagna (<https://www.arpae.it/it/temi-ambientali/meteo>) for providing us with meteorological data. We are grateful to two anonymous reviewers for improving earlier drafts with their comments.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ecolind.2025.113806>.

Data availability

Data are available from the authors upon reasonable request

References

- Albert, A., Auffret, A.G., Cosyns, E., et al., 2015. Seed dispersal by ungulates as an ecological filter: a trait-based metaanalysis. *Oikos* 124, 1109–1120.
- Adams, P.J., Fontaine, J.B., Huston, R.M., Fleming, P.A., 2019. Quantifying efficacy of feral pig (*Sus scrofa*) population management. *Wildl. Res.* 46, 587–598.
- Ballari S.A., Barrios-García M.N., 2014. A review of wild boar *Sus scrofa* diet and factors affecting food selection in native and introduced ranges. *Mammal Rev.* 44, 124–134.
- Barasona, J.A., Carpio, A., Boadella, M., et al., 2021. Expansion of native wild boar populations is a new threat for semi-arid wetland areas. *Ecol. Ind.* 125, 107563.
- Barrios-García, M.N., Ballari, S.A., 2012. Impact of wild boar (*Sus scrofa*) in its introduced and native range: a review. *Biol. Invasions* 14, 2283–2300.
- Barrios-García, M.N., Gonzalez-Polo, M., Simberloff, D., et al., 2023. Wild boar rooting impacts soil function differently in different plant community types. *Biol. Invasions* 25, 583–592.
- Bartoń, K., 2023. MuMIn: Multi-Model Inference. R package version 1.47. 5. 2023.
- Baubet, E., Bonenfant, C., Brandt, S., 2004. Diet of the wild boar in the French Alps. *Galemys* 16, 101–113.
- Belnap, J., 1995. Surface disturbances: their role in accelerating desertification. *Environ. Monit. Assess.* 37, 39–57.
- Bracke, M.B.M., 2011. Review of wallowing in pigs: description of the behaviour and its motivational basis. *Appl. Anim. Behav. Sci.* 132, 1–13.
- Brooks, M.E., et al., 2017. GlmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modelling. *The R Journal.* 9, 378–400.
- Boitani, L., et al., 1994. Spatial and activity patterns of wild boars in Tuscany Italy. *J. Mammal.* 75, 600–612.
- Bratton, S.P., 1974. The effect of the European wild boar (*Sus scrofa*) on the high-elevation vernal flora in Great Smoky Mountains National Park. *Bulletin of the Torrey Botanical Club* 101, 198–206.
- Briedermann, L., 1990. Schwarzwild, 2nd edn. VEB Deutscher Landwirtschaftsverlag, Berlin, Germany.

- Brivio, F., Grignolio, S., Brogi, R., et al., 2017. An analysis of intrinsic and extrinsic factors affecting the activity of a nocturnal species: the wild boar. *Mamm. Biol.* 84, 73–81.
- Brogi, R., Grignolio, S., Brivio, F., Apollonio, M., 2020. Protected areas as refuges for pest species? The case of wild boar. *Global Ecol. Conserv.* 22, e00969.
- Bueno, C.G., Alados, C.L., Gómez-García, D., Barrio, I., García-González, R., 2009. Understanding the main factors in the extent and distribution of wild boar rooting on alpine grasslands. *J. Zool.* 279, 195–202.
- Bueno, C.G., Azorín, J., Gómez-García, D., Alados, C.L., Badía, D., 2013. Occurrence and intensity of wild boar disturbances, effects on the physical and chemical soil properties of alpine grasslands. *Plant and Soil* 373, 243–256.
- Bueno, C.G., Jiménez, J.J., 2014. Livestock grazing activities and wild boar rooting affect alpine earthworm communities in the Central Pyrenees (Spain). *Appl. Soil Ecol.* 83, 71–78.
- Bueno, C.G., Reiné, R., Alados, C.L., Gómez-García, D., 2011. Effects of large wild boar disturbances on alpine soil seed banks. *Basic Appl. Ecol.* 12, 125–133.
- Burrascano, S., Copiz, R., Del Vico, E., et al., 2015. Wild boar rooting intensity determines shifts in understorey composition and functional traits. *Community Ecol.* 16, 244–253.
- Calosi, M., Gabbrivelli, C., Lazzeri, L., Fattorini, N., Cesaretti, G., Burrini, L., Petrillo, O., Ferretti, F., 2024. Seasonal and ecological determinants of wild boar rooting on priority protected grasslands. *Environ. Manag.* 74, 268–281.
- Caplenor, C.A., Poudyal, N.C., Muller, L.L., Yoest, C., 2017. Assessing landowners' attitudes toward wild hogs and support for control options. *J. Environ. Manage.* 201, 45–51.
- Capitani, C., Bertelli, L., Varuzza, P., Scandura, M., Apollonio, M., 2004. A comparative analysis of wolf (*Canis lupus*) diet in three different Italian ecosystems. *Mamm. Biol.* 69, 1–10.
- Carpio, A.J., Apollonio, M., Acevedo, P., 2021. Wild ungulate overabundance in Europe: contexts, causes, monitoring and management recommendations. *Mammal Rev.* 51, 95–108.
- Colomer, J., Massei, G., Roos, D., Rosell, C., Rodriguez-Teijeiro, J.D., 2024. What drives wild boar density and population growth in Mediterranean environments? *Sci. Total Environ.* 931, 172739.
- Copernicus Land Monitoring Service, 2023. CORINE Land Cover (CLC) inventory. Available at: <https://land.copernicus.eu/pan-european/corine-land-cover> [Accessed 05/03/2024].
- Côté, S.D., Rooney, T.P., Tremblay, J.P., Dussault, C., Waller, D.M., 2004. Ecological impacts of deer overabundance. *Ann. Rev. Ecol. Syst.* 35, 113–147.
- Ding, Y., Li, Z., Peng, S., 2020. Global analysis of time-lag and -accumulation effects of climate on vegetation growth. *Int. J. Appl. Earth Obs. Geoinf.* 92, 102179.
- European Union Court of Justice, 2013. Case C-258/11 Peter Sweetman and Others v An Bord Pleanála. Request for a preliminary ruling from the Supreme Court (Ireland). Environment — Directive 92/43/EEC — Article 6 — Conservation of natural habitats — Special areas of conservation — Assessment of the implications for a protected site of a plan or project — Criteria to be applied when assessing the likelihood that such a plan or project will adversely affect the integrity of the site concerned — Lough Corrib site — N6 Galway City Outer Bypass road scheme.
- Fattorini, L., Ferretti, F., Pisani, C., Sforzi, A., 2011. Two-stage estimation of ungulate abundance in Mediterranean areas using pellet group count. *Environ. Ecol. Stat.* 18, 291–314.
- Fattorini, N., Ferretti, F., 2020. Estimating wild boar density and rooting activity in a Mediterranean protected area. *Mamm. Biol.* 100, 241–251.
- Fattorini, L., Franceschi, S., Pisani, C., 2024. Design-based consistent strategies exploiting auxiliary information in environmental mapping. *J. Agric. Biol. Environ. Stat.* <https://doi.org/10.1007/s13253-024-00664-4>.
- Felton, A.M., Hedwall, P.O., Felton, A., Widemo, F., Wallgren, M., Holmström, E., et al., 2022. Forage availability, supplementary feed and ungulate density: associations with ungulate damage in pine production forests. *For. Ecol. Manage.* 513, 120187.
- Fernandez-Llario, P., Mateos-Quesada, P., 2005. Influence of rainfall on the breeding biology of Wild boar (*Sus scrofa*) in a Mediterranean ecosystem. *Folia Zool.* 54, 240.
- Ferretti, F., Fattorini, L., Sforzi, A., Pisani, C., 2016. The use of faeces counts to estimate relative densities of wild boar in a Mediterranean area. *Popul. Ecol.* 58, 329–334.
- Ferretti, F., Lazzeri, L., Fattorini, N., 2023. A test of motion-sensitive cameras to index ungulate densities: group size matters. *J. Wildl. Manag.* 87, e22356.
- Ferretti, F., Lazzeri, L., Mori, E., Cesaretti, G., Calosi, M., Burrini, L., Fattorini, N., 2021. Habitat correlates of wild boar density and rooting along an environmental gradient. *J. Mammal.* 102, 1536–1547.
- Ferretti, F., Lovari, S., Mancino, V., Burrini, L., Rossa, M., 2019. Food habits of wolves and selection of wild ungulates in a prey-rich Mediterranean coastal area. *Mamm. Biol.* 99, 119–127.
- Fettebert, J., Baubet, E., Slotow, R., Fischer, C., 2017. Landscape effects on wild boar home range size under contrasting harvest regimes in a human-dominated agro-ecosystem. *Eur. J. Wildl. Res.* 63, 32.
- Foster, C.N., Barton, P.S., Lindenmayer, D.B., 2014. Effects of large native herbivores on other animals. *J. Appl. Ecol.* 51, 929–938.
- Gordigiani, L., Viviano, A., Brivio, F., Grignolio, S., Lazzeri, L., Marcon, A., Mori, E., 2022. Carried away by a moonlight shadow: activity of wild boar in relation to nocturnal light intensity. *Mammal Res.* 67, 39–49.
- Groot, B.G., Hazebroek, E., 1996. Wild boar (*Sus scrofa scrofa* L.) rooting and forest regeneration on podzolic soils in the Netherlands. *For. Ecol. Manage.* 88, 71–80.
- Harrison, X.A., Donaldson, L., Correa-Cano, M.E., Evans, J., Fisher, D.N., Goodwin, C.E. D., Robinson, B.S., Hodgson, D.J., Inger, R., 2018. A brief introduction to mixed effects modelling and multi-model inference in ecology. *PeerJ* 6, e4794.
- Hebblewhite, M., White, C.A., Nietvelt, C.G., McKenzie, J.A., Hurd, T.E., Fryxell, J.M., Bayley, S.E., Paquet, P.C., 2005. Human activity mediates a trophic cascade caused by wolves. *Ecology* 86, 2135–2144.
- Herrero, J., García-Serrano, A., Couto, S., Ortuño, V.M., García-González, R., 2006. Diet of wild boar *Sus scrofa* L. and crop damage in an intensive agroecosystem. *Eur. J. Wildl. Res.* 52, 245–250.
- Hobbs, N.T., 1996. Modification of ecosystems by ungulates. *J. Wildl. Manag.* 60, 695.
- Hone, J., 1995. Spatial and temporal aspects of vertebrate pest damage with emphasis on feral pigs. *J. Appl. Ecol.* 32, 311–319.
- Hone, J., 2002. Feral pigs in Namadgi National Park, Australia: dynamics, impacts and management. *Biol. Conserv.* 105, 231–242.
- Horčíčková, E., Brůna, J., Vojta, J., 2019. Wild boar (*Sus scrofa*) increases species diversity of semi-rural grassland: field experiment with simulated soil disturbances. *Ecol. Evol.* 9, 2765–2774.
- Howe, T.D., Singer, F.J., Ackerman, B.B., 1981. Forage relationships of European wild boar invading northern hardwood forest. *J. Wildl. Manag.* 45, 748–754.
- Johann, F., Handschuh, M., Linderoth, P., Dormann, C.F., Arnold, J., 2020. Adaptation of wild boar (*Sus scrofa*) activity in a human-dominated landscape. *BMC Ecol.* 20, 4.
- Keuling, O., Stier, N., Roth, M., 2009. Commuting, shifting or remaining? Different spatial utilisation patterns of wild boar *Sus scrofa* L. in forest and field crop during summer. *Mamm. Biol.* 74, 145–152.
- Kaštovská, E., Mastný, J., Konvička, M., 2024. Rewilding by large ungulates contributes to organic carbon storage in soils. *J. Environ. Manage.* 355, 120430.
- Kubinec, R., 2023. Ordered beta regression: a parsimonious, well-fitting model for continuous data with lower and upper bounds. *Polit. Anal.* 31, 519–536.
- Labadessa, R., Ancillotto, L., 2023. Beauty and the beast: multiple effects of wild boar rooting on butterfly microhabitat. *Biodivers. Conserv.* 32, 1189–1204.
- Lange, J., Arbel, Y., Grodek, T., Greenbaum, N., 2010. Water percolation process studies in a Mediterranean karst area. *Hydrol. Process.* 24, 1866–1879.
- Laguna, E., Barasona, J.A., Vicente, J., Keuling, O., Acevedo, P., 2021. Differences in wild boar spatial behaviour among land uses and management scenarios in Mediterranean ecosystems. *Sci. Total Environ.* 796, 148966.
- Lazzeri, L., Belardi, I., Pacini, G., Fattorini, N., Ferretti, F., 2024. Beyond ungulate density: prey switching and selection by the wolf in a recolonised area. *Global Ecol. Conserv.* 54, e03069.
- Lingle, S., 2001. Anti-predator strategies and grouping patterns in white-tailed deer and mule deer. *Ethology* 107, 295–314.
- Lombardini, M., Meriggi, A., Fozzi, A., 2017. Factors influencing wild boar damage to agricultural crops in Sardinia (Italy). *Curr. Ecol.* 63, 507–514.
- Lovari, S., Sangiuliano, A., 2006. Il lupo sul M. Amiata. Progetto sui grandi canidi (lupo, cane) nel territorio dell'Amiata grossetana e senese. Technical Report, Comunità Montana Amiata Grossetana e Ministero del Ambiente. Arcidosso and Rome, Italy. DOI: 10.13140/RG.2.1.3653.2648.
- Lovari, S., Sforzi, A., Mori, E., 2013. Habitat richness affects home range size in a monogamous large rodent. *Behav. Process.* 99, 42–46.
- Lucherini, M., Lovari, S., 1996. Habitat richness affects home range size in the red fox *Vulpes vulpes*. *Behav. Process.* 36, 103–105.
- Mangipanea, L.S., Belant, J.L., Hiller, T.L., et al., 2018. Influences of landscape heterogeneity on home-range sizes of brown bears. *Mamm. Biol.* 88, 1–7.
- Markov, N., Economov, A., Hjeljord, O., et al., 2022. The wild boar *Sus scrofa* in northern Eurasia: a review of range expansion history, current distribution, factors affecting the northern distributional limit, and management strategies. *Mammal Rev.* 52, 519–537.
- Massei, G., Coats, J., Lambert, M.S., et al., 2018. Camera traps and activity signs to estimate wild boar density and derive abundance indices. *Pest Manag. Sci.* 74, 853–860.
- Massei, G., Genov, P.V., 2004. The environmental impact of wild boar. *Galemys* 16, 135–145.
- Massei, G., Genov, P.V., Staines, B.W., Gorman, M.L., 1997. Factors influencing home range and activity of wild boar (*Sus scrofa*) in a Mediterranean coastal area. *J. Zool.* 242, 411–423.
- Massei, G., Kindberg, J., Licoppe, A., et al., 2015. Wild boar populations up, numbers of hunters down? A review of trends and implications for Europe. *Pest Manag. Sci.* 71, 492–500.
- Mattioli, L., Capitani, C., Avanzinelli, E., Gazzola, A., Apollonio, M., 2004. Predation by wolves (*Canis lupus*) on roe deer (*Capreolus capreolus*) in north-eastern Apennine Italy. *J. Zool.* 264, 249–258.
- Mayle, B.A., Peace, A.J., Gill, R.M.A., 1999. How many deer? A field guide to estimating deer population size. Edinburgh, United Kingdom: Forestry Commission (No. 18, pp. viii+96).
- Minder, I., 2011. Local and seasonal variations of roe deer diet in relation to food resource availability in a Mediterranean environment. *Eur. J. Wildl. Res.* 58, 215–225.
- Mohr, D., Cohnstaedt, L.W., Topp, W., 2005. Wild boar and red deer affect soil nutrients and soil biota in steep oak stands of the Eifel. *Soil Biol. Biochem.* 37, 693–700.
- Morelle, K., Fettebert, J., Mengal, C., Lejeune, P., 2016. Invading or recolonizing? Patterns and drivers of wild boar population expansion into Belgian agroecosystems. *Agr. Ecosyst. Environ.* 222, 267–275.
- Mori, E., Benatti, L., Lovari, S., Ferretti, F., 2016. What does the wild boar mean to the wolf? *Eur. J. Wildl. Res.* 63, 9.
- Mori, E., Ferretti, F., Lagrotteria, A., La Greca, L., Solano, E., Fattorini, N., 2020. Impact of wild boar rooting on small forest-dwelling rodents. *Ecol. Res.* 35, 675–681.
- Nakagawa, S., Cuthill, I.C., 2007. Effect size, confidence interval and statistical significance: a practical guide for biologists. *Biol. Rev.* 82, 591–605.
- Ohashi, H., Saito, M., Horie, R., Tsunoda, H., Noba, H., Ishii, H., Kuwabara, T., Hiroshige, Y., Koike, S., Hoshino, Y., Toda, H., Kaji, K., 2013. Differences in the

- activity pattern of the wild boar *Sus scrofa* related to human disturbance. *Eur. J. Wildl. Res.* 59, 167–177.
- Olczak, K., Nowicki, J., Klocek, C., 2015. Pig behaviour in relation to weather conditions – a review. *Annals of Animal Sciences.* 15, 601–610.
- Palacio, S., Bueno, C.G., Azorin, J., Maestro, M., Gómez-García, D., 2013. Wild boar disturbance increases nutrient and C stores of geophytes in subalpine grasslands. *Am. J. Bot.* 100, 1790–1799.
- Pascual-Rico, R., Morales-Reyes, Z., Aguilera-Alcalá, N., et al., 2021. Usually hated, sometimes loved: a review of wild ungulates' contributions to people. *Sci. Total Environ.* 801, 149652.
- Penna, D., Borga, M., Norbiato, D., Dalla, F.G., 2009. Hillslope scale soil moisture variability in a steep alpine terrain. *J. Hydrol.* 364, 311–327.
- Pesaresi, S., Galdenzi, D., Biondi, E., Casavecchia, S., 2014. Bioclimate of Italy: application of the worldwide bioclimatic classification system. *J. Maps* 10, 538–553.
- Pilhal, R., Kamler, J., Homolka, M., 2014. Faecal pellet group counting as a promising method of wild boar population density estimation. *Acta Theriol.* 59, 561–569.
- Podgórski, T., Baś, G., Jędrzejewska, B., Sönnichsen, L., Śniezko, S., Jędrzejewski, W., Okarma, H., 2013. Spatiotemporal behavioral plasticity of wild boar (*Sus scrofa*) under contrasting conditions of human pressure: primeval forest and metropolitan area. *J. Mammal.* 94, 109–119.
- R Core Team, 2023. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. <<https://www.R-project.org/>>.
- Ripple, W.J., Newsome, T.M., Wolf, C., et al., 2015. Collapse of the world's largest herbivores. *Sci. Adv.* 1, e1400103.
- Ruf, T., Vetter, S.G., Painer-Gigler, J., Stalder, G., Bieber, C., 2023. Thermoregulation in the wild boar (*Sus scrofa*). *J. Comp. Physiol. B* 193, 689–697.
- Sandom, C.J., Hughes, J., Macdonald, D.W., 2013a. Rooting for rewilding: quantifying wild boar's *Sus scrofa* rooting rate in the scottish highlands. *Restor. Ecol.* 21 (3), 329–335.
- Sandom, C.J., Hughes, J., Macdonald, D.W., 2013b. Rewilding the scottish highlands: do wild boar, *Sus scrofa*, use a suitable foraging strategy to be effective ecosystem engineers? *Restor. Ecol.* 21 (3), 336–343.
- Säid, S., Servanty, S., 2005. The influence of landscape structure on female roe deer home-range size. *Landsc. Ecol.* 20, 1003–1012.
- Scandura, M., Podgórski, T., Vicente, J., Iacolina, L., 2022. Wild Boar *Sus scrofa* Linnaeus, 1758. In: Corlatti, L., Zachos, F.E. (Eds.), *Terrestrial Cetartiodactyla, Handbook of the Mammals of Europe*. Springer International Publishing, Cham, pp. 1–27.
- Scandurra, A., Magliozzi, L., Fulgione, D., Aria, M., D'aiello, B., 2016. Lepidoptera Papilionoidea communities as a sentinel of biodiversity threat: the case of wild boar rooting in a Mediterranean habitat. *J. Insect Conserv.* 2016, 353–362.
- Scherer, G., Streanga, B., Fartmann, T., 2025. Rare butterfly species vitally depend on soil disturbance by an ecosystem engineer in abandoned calcareous grasslands. *Global Ecol. Conserv.* 58, e03451.
- Schiavina, M., Melchiorri, M., Pesaresi, M., Politis, P., Carneiro Freire, S.M., Maffenini, L., Florio, P., Ehrlich, D., Goch, K., Carioli, A., Uhl, J., Tommasi, P. and Kemper, T., 2023. GHSL Data Package 2023, Publications Office of the European Union, Luxembourg, 2023, doi:10.2760/098587, JRC133256.
- Schley, L., Roper, T.J., 2003. Diet of wild boar *Sus scrofa* in Western Europe, with particular reference to consumption of agricultural crops. *Mammal Rev.* 33, 43–56.
- Singer, F.J., Swank, W.T., Clebsh, E.E.C., 1984. Effects of wild pig rooting in a deciduous forest. *J. Wildl. Manag.* 48, 464–473.
- Sondej, I., Kwiatkowska-Falinska, A.J., 2017. Effects of wild boar (*Sus scrofa* L.) rooting on seedling emergence in Białowieża. *Forest Policy Econ.* 65, 380–389.
- Spitz, F., Janeau, G., 1995. Daily selection of habitat in wild boar (*Sus scrofa*). *J. Zool.* 237, 423–434.
- Sütő, D., Farkas, J., Siffer, S., Schally, G., Katona, K., 2020. Spatiotemporal pattern of wild boar rooting in a Central European dry oak forest. *Eur. J. For. Res.* 139, 407–418.
- Thurjell, H., Ball, J.P., Ahlén, P., Kornacher, P., Dettki, H., Sjöberg, K., 2009. Habitat use and spatial patterns of wild boar *Sus scrofa* (L.): agricultural fields and edges. *Eur. J. Wildl. Res.* 55, 517–523.
- Thurjell, H., Spong, G., Ericsson, G., 2014. Effects of weather, season, and daylight on female wild boar movement. *Acta Theriol.* 59, 467–472.
- Thurjell, H., Spong, G., Olsson, M., Ericsson, G., 2015. Avoidance of high traffic levels results in lower risk of wild boar-vehicle accidents. *Landsc. Urban Plan.* 133, 98–104.
- Tierney, T.A., Cushman, J.H., 2006. Temporal changes in native and exotic vegetation and soil characteristics following disturbances by feral pigs in a California grassland. *Biol. Invasions* 8, 1073–1089.
- Treichler, J.W., VerCauteren, K.C., Taylor, C.R., Beasley, J.C., 2023. Changes in wild pig (*Sus scrofa*) relative abundance, crop damage, and environmental impacts in response to control efforts. *Pest Manag. Sci.* 79, 4765–4773.
- Van de Pol, M., Wright, J., 2009. A simple method for distinguishing within-versus between-subject effects using mixed models. *Anim. Behav.* 77, 753.
- Vetter, S.G., Ruf, T., Bieber, C., Arnold, W., 2015. What is a mild winter? Regional differences in within-species responses to climate change. *PLoS One* 10, e0132178.
- Ward, D., Saltz, D., 1994. Foraging at different spatial scales: dorcas gazelles foraging for lilies in the Negev desert. *Ecology* 75, 48–58.
- Welander, J., 2000. Spatial and temporal dynamics of wild boar (*Sus scrofa*) rooting in a mosaic landscape. *J. Zool.* 252, 263–271.
- Wirthner, S., Frey, B., Busse, M.D., Schütz, M., Risch, A.C., 2011. Effects of wild boar (*Sus scrofa* L.) rooting on the bacterial community structure in mixed-hardwood forest soils in Switzerland. *Eur. J. Soil Biol.* 47, 296–302.
- Yang, G., Peng, C., Yang, X., Guo, Q., Su, H., 2024. Habitat suitability and crop damage risk caused by wild boar in Guizhou Plateau China. *J. Wildl. Manag.* 88, e22542.
- Zuur, A.F., Ieno, E.N., Walker, N.J., Saveliev, A.A., Smith, G.M., 2009. *Mixed effects models and extensions in ecology with R*. Springer, New York.