

HIV INFECTION AMONG ADOLESCENTS AND YOUNG ADULTS
IN ITALY: EPIDEMIOLOGY, VIRO-IMMUNOLOGICAL AND
TREATMENT OUTCOMES

Francesco Maria Fusco , Ilaria Mastrorosa , Francesca Vichi ,
Adriana Cervo , Annalisa Saracino , Leonardo Calza ,
Eugenia Quiros-Roldan , Maria Elisabetta Teti , Andrea Costantini ,
Antonella Cingolani , Enrico Girardi , Andrea Antinori ,
Antonella d'Arminio Monforte , Alessandro Tavelli , for the ICONA
Foundation Study Group

PII: S1201-9712(25)00422-9
DOI: <https://doi.org/10.1016/j.ijid.2025.108200>
Reference: IJID 108200

To appear in: *International Journal of Infectious Diseases*

Received date: 14 September 2025
Revised date: 5 November 2025
Accepted date: 7 November 2025

Please cite this article as: Francesco Maria Fusco , Ilaria Mastrorosa , Francesca Vichi ,
Adriana Cervo , Annalisa Saracino , Leonardo Calza , Eugenia Quiros-Roldan ,
Maria Elisabetta Teti , Andrea Costantini , Antonella Cingolani , Enrico Girardi , Andrea Antinori ,
Antonella d'Arminio Monforte , Alessandro Tavelli , for the ICONA Foundation Study Group, HIV
INFECTION AMONG ADOLESCENTS AND YOUNG ADULTS IN ITALY: EPIDEMIOLOGY, VIRO-
IMMUNOLOGICAL AND TREATMENT OUTCOMES, *International Journal of Infectious Diseases*
(2025), doi: <https://doi.org/10.1016/j.ijid.2025.108200>

This is a PDF of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability. This version will undergo additional copyediting, typesetting and review before it is published in its final form. As such, this version is no longer the Accepted Manuscript, but it is not yet the definitive Version of Record; we are providing this early version to give early visibility of the article. Please note that Elsevier's sharing policy for the Published Journal Article applies to this version, see: <https://www.elsevier.com/about/policies-and-standards/sharing#4-published-journal-article>. Please also note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2025 Published by Elsevier Ltd on behalf of International Society for Infectious Diseases.
This is an open access article under the CC BY-NC-ND license
(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

HIGHLIGHTS

- Milder clinical presentation and more robust immune recovery for AYAs
- Higher risk of TD due to patient choice/adherence issues and LTFU for AYAs
- Dedicated clinical care pathways need to be created for AYAs.

Journal Pre-proof

Downloaded for Anonymous User (n/a) at IRCCS San Gerardo dei Tiri
November 20, 2025. For personal use only. No other uses without permission.

HIV INFECTION AMONG ADOLESCENTS AND YOUNG ADULTS IN ITALY: EPIDEMIOLOGY, VIRO-IMMUNOLOGICAL AND TREATMENT OUTCOMES

Authors

Francesco Maria Fusco^{1§}, Iaria Mastrosera^{2§*}, Francesca Vichi³, Adriana Cervo⁴, Annalisa Saracino⁵, Leonardo Calza⁶, Eugenia Quiros-Roldan⁷, Maria Elisabetta Teti⁸, Andrea Costantini⁹, Antonella Cingolani¹⁰, Enrico Girardi¹¹, Andrea Antinori^{2,12}, Antonella d'Arminio Monforte¹³, Alessandro Tavelli^{13,14} for the *ICONA Foundation Study Group*

§ These authors equally contributed to this work

Affiliations

¹ UOC Infezioni sistemiche e dell'Immunodepresso, PO "D. Cotugno", AO dei Colli, Napoli, Italy; ² Clinical and Research Infectious Diseases Department, National Institute for Infectious Diseases Lazzaro Spallanzani IRCCS, Rome, Italy; ³ Infectious diseases, Florence; ⁴ Department of Infectious Diseases, Azienda Ospedaliero-Universitaria di Modena, Policlinico di Modena, Modena, Italy; ⁵ Clinic of Infectious Diseases, Department of Precision and Regenerative Medicine and Ionian Area, Polyclinic of Bari, University Hospital Polyclinic, University of Bari, Bari, Italy; ⁶ Infectious Diseases Unit, IRCCS Policlinico di Sant'Orsola, Alma Mater Studiorum University of Bologna, Bologna, Italy; ⁷ Department of Clinical and Experimental Sciences, Unit of Infectious and Tropical Diseases, University of Brescia and ASST Spedali Civili di Brescia, Brescia, Italy; ⁸ Clinical Infectious Diseases, Policlinico Tor Vergata of Rome, Rome, Italy; ⁹ Clinical Immunology Unit, Azienda Ospedaliero Universitaria delle Marche - Università Politecnica delle Marche, Ancona, Italy; ¹⁰ Unit of Infectious Diseases, Fondazione Policlinico Universitario Agostino Gemelli IRCCS, Department of Safety and Bioethics, Università Cattolica del Sacro Cuore, Rome, Italy; ¹¹ Scientific Direction, National Institute for Infectious Diseases Lazzaro Spallanzani IRCCS, Rome, Italy; ¹² Health Direction, National Institute for Infectious Diseases Lazzaro Spallanzani IRCCS, Rome, Italy; ¹³ Fondazione ICONA, Milan, Italy; ¹⁴ National PhD Programme in One Health approaches to infectious

diseases and life science research, Department of Public Health, Experimental and Forensic Medicine,
University of Pavia, Pavia, Italy.

***Corresponding author:**

Ilaria Mastrorosa, MD

Via Portuense 292, 00149 Rome, Italy

Tel.: +39 06 55170546

E-mail: ilaria.mastrorosa@inmi.it.

Journal Pre-proof

Downloaded for Anonymous User (n/a) at IRCCS San Gerardo dei T
November 20, 2025. For personal use only. No other uses without pe

ABSTRACT

Objectives. No previous study described adolescents and young adults (AYAs) with HIV in Italy. Aims were to investigate the temporal trend of AYAs enrolment in the ICONA cohort over 2014-2023, and to compare their baseline characteristics and antiretroviral treatment (ART) outcomes with adults.

Methods. All subjects enrolled in ICONA, the Italian cohort enrolling HIV-1 positive individuals from ART-naïve, over 2014-2023 were grouped into: AYAs (18-24 years old) and adults (≥ 25). Study outcomes were: time to first ART start, time to treatment discontinuation (TD) of first-line regimen, cause-specific TD, virologic failure, and Loss To Follow Up (LTFU).

Results. 9,519 participants: 653 AYAs (6.9%) and 8,866 adults (93.1%). Excluding the decline recorded in 2020, the percentage of AYAs enrolled showed a similar trend before- and after-COVID.

Compared with adults, AYAs had a milder clinical presentation, a better viro-immunological status, a higher mean change of CD4 count over 24 months and, after adjusting, a higher risk of TD for patient's choice/adherence-related issues and LTFU.

Conclusions. The less advanced clinical presentation and more robust immune recovery of AYAs, might depend upon more recent HIV infection, better thymic function, and lower immune activation. The higher risk of TD due to patient's choice/adherence-related issues and LTFU among AYAs is a major concern and suggests the need to create *ad hoc* clinical care pathways.

Keywords (6): adolescent and young adults; HIV infection; antiretroviral therapy; virological failure; treatment discontinuation; loss to follow up

MAIN TEXT

Introduction

In Europe, age-specific rates of new HIV diagnoses declined from 2014 to 2020, remained stable between 2020 and 2021, and increased from 2022 across most age groups [1]. Similarly, in Italy, according to the most recent report of the national surveillance system [2], after the overall decrease in the number of new HIV diagnoses observed from 2012 to 2020, a reversal of the trend has been described in the last three years, and, in particular, the proportion of elderly people (over 50 years old) is gradually increasing. Despite this, a part of new diagnoses still occurs among Adolescents and Young Adults (AYAs). According to the Centers for Disease Control and Prevention, adolescents (13 to 19 years of age) and young adults (20 to 24 years of age) accounted for 20% of the 31,800 HIV diagnoses in the United States 2022 [3]. In Europe, between 2022 and 2023, the only groups that continued to see an increase in new HIV diagnoses were women aged 15 to 24, with a 15.8% rise, and men aged 15 to 19, where the rate rose by 20.6% [1]. In Italy, about 7% of new HIV diagnoses occurs in people under the age of 25 [2].

Downloaded for Anonymous User (n/a) at IRCCS San Gerardo dei Tiri
November 20, 2025. For personal use only. No other uses without permission.

Data from literature suggest that this population has some common characteristics [4-10]: AYAs are less likely to report currently taking antiretroviral therapy (ART) and to being adherent to ART, have a higher rate of loss-to-follow-up and are more likely to have a treatment interruption. Consequently, AYAs as population could have lower levels of viral suppression compared to older patients. Moreover, most people who acquire HIV during adolescence and young adulthood get it through sexual transmission and AYAs have peculiar lifestyle behaviours, as active sexual life, inconsistent use of condoms and frequent alcohol or drug use. Finally, high rates of sexually transmitted infections (STIs) among young people may increase the risk of getting or spreading HIV [11]. Consequently, in this population, the over mentioned gaps in HIV care may significantly contribute to disease transmission. These behavioural aspects, together with social and psychological vulnerability, contribute to make HIV infection a real challenge, both for the patient and the treating physician.

Most of the available data come from studies conducted, many of them not so recently, in the United States and Africa, where social determinants and models of access to healthcare services may differ from Europe. Few data are available about this population in Europe, and no studies of AYAs with HIV infection have ever been conducted in Italy.

Herein, we plan to give a real-life picture of AYAs who acquired HIV infection in Italy in the last decade. Specifically, the present study is designed to: (i) investigate the trend over time of AYAs enrolment in the ICONA (Italian Cohort of Naïve to Antiretrovirals) cohort from 2014 to 2023; (ii) evaluate the epidemiological, socio-behavioural, clinical and viro-immunological profiles of AYAs at enrolment compared to adult population; (iii) describe and compare ART treatment initiation, first line ART discontinuation and viro-immunological response in AYAs and adults; (iv) assess and compare the risk of loss to follow-up of AYAs and adults over the period of observation.

Downloaded for Anonymous User (n/a) at IRCCS San Gerardo dei T
November 20, 2025. For personal use only. No other uses without pe

Methods

Study setting and ethical issues

Retrospective analysis of prospectively collected data from the ICONA cohort. The ICONA Foundation cohort is an Italian multicentric observational study, set up in 1997, of HIV-1 positive and antiretroviral-naïve subjects at the time of enrolment, aged ≥ 18 years. More than 21,500 people with HIV (PWH) were enrolled and prospectively followed in 62 adults' Italian Infectious Diseases centres. PWH enrolled in ICONA are representative of the Italian HIV population [12]. In the present study, all individuals newly enrolled in ICONA from January 2014 to December 2023 were included, after providing a written informed consent, in order to compare participants aged 18-24 years at enrolment (Adolescent and Young Adults- AYAs) with subjects aged more than 24 (adults).

See supplementary materials for more details.

Statistical Analysis

Demographic and clinical differences at ART-initiation between the two groups were assessed using chi-square for categorical variables or non-parametric Kruskal-Wallis test for continuous variables.

Proportion of AYAs enrolment over total number of PWH included in ICONA per year has been evaluated. Moreover, in order to evaluate the temporal trends of AYAs proportion and to assess changes related to the COVID-19 pandemic, we performed an Interrupted Time Series Analysis (ITSA). The analysis was conducted using annual data from 2014 to 2023, with the year 2020 identified as the change point and comparing the trends in the two periods. Time to first ART regimen initiation from the date of HIV diagnosis was assessed using unadjusted and adjusted Cox regression models, including only subjects with HIV diagnosis after 2015, year of recommendation of universal ART start in Italy, and with at least one follow-up visit after enrolment or known ART initiation. Time to Virologic Failure (VF: 2 consecutive HIV-RNA ≥ 50 copies/ml after 6 months of the first ART regimen) from date of first ART start and time to Treatment Discontinuation (TD: discontinuation for any reason of at least one drug of the regimen or treatment intensification) were also analysed using standard survival analysis (Kaplan-Meier curves and log-rank test) and unadjusted and adjusted Cox regression models. Survival analysis and unadjusted and adjusted Cox regression models were also used to assess cause-specific risks of first ART discontinuation: TD due to patient's choice or adherence issues, TD due to toxicity/intolerance (gastrointestinal intolerance, neuropsychiatric, renal, metabolic and dermatologic adverse events, allergies, other toxicities) and TD for failure (virological, immunological and/or clinical). For the ART discontinuation endpoints participants' follow-up accrued from the date of first ART start until discontinuation or to the last available clinical visit. For each cause-specific TD endpoint subjects who discontinued for other reasons, were truncated at the date of last clinical follow-up, assuming non-informative censoring. For the virological failure endpoint censoring was applied at the date of participants' last available viral load measure. An intention-to-treat approach (which ignore any ART change) was used for the virological failure analysis.

Furthermore, we assessed time to Loss to Follow Up (LTFU) through unadjusted and adjusted Cox regression models, considering as endpoint no follow-up visit for more than 12 months or with the last visit before January 2023 and excluding participants who died, or moved to another country or

another hospital/center not included in the ICONA cohort network, those incarcerated, or PWH with voluntary withdrawn from the ICONA study within 12 months after the last visit. For this endpoint, we considered only the 16 ICONA centers with an active update of the follow-up visits over the period 2016-2023 with electronic import of laboratory and clinical visits data, in order to minimize possible bias due to missing data or delay in data reporting and to get a more accurate estimate for the overall rate of loss to care. Follow-up accrued at the date of enrolment till last available follow-up in the cohort. A second definition of Definitive LTFU has also been used, considering only the subgroup of LTFU as previously defined that never returned to care.

Finally, immunological response has been investigated as CD4 changes from baseline (ART initiation) to 6 months (± 2 months), 12 months (± 3 months) and 24 months (± 4 months), by means of adjusted linear mixed models with random intercept and slopes per patient. Interaction terms (time \times age group) were created to assess whether these changes differed significantly over time between

AYAs and adults. We further tested whether there was an interaction between the class of the ART regimen on the first line (Integrase Strand Transfer Inhibitors - INSTI - based *vs* non-INSTI based) and AYAs/Adults on the CD4 count recovery on a three-way interaction (time \times age group \times INSTI). The equality of all mean changes in the groups was tested by Wald test. All the models have been adjusted for pre-defined factors considered as confounders of the association between the exposure of interest (AYAs/adults) and each separate outcome. All statistical analyses were performed using STATA (v.18.0, StataCorp, USA) and RStudio (v 12.0). All p-values presented are two-sided and p-value < 0.05 indicated conventional statistical significance.


Results

Descriptive analysis

Over the study period, a total of 9,519 individuals were included in ICONA: 653 AYAs (6.9%) and 8,866 adults (93.1%). From the beginning of the observation period until 2019, the trend of proportion of AYAs enrolled showed a small increase, although the confidence interval included zero (2014–

2019 coefficient = +0.25%/year; 95%CI -0.42, +0.92; $p=0.399$). In 2020, a steep drop of approximately 3.2% points was observed compared to the expected value based on the pre-2020 trend (level change coefficient = -3.23%; 95%CI -6.60, +0.15; $p= 0.058$). No evidence for a change was found in the trend after 2020 compared to the previous period (2014-2019) (change 2020/2023 coefficient - 2014/2019 coefficient = -0.13%/year; 95%CI -1.15, +0.89; $p=0.771$) (Figure 1).

Figure 1. Proportion of adolescents and young adults (AYAs) enrolled in the ICONA cohort over the study period, from 2014 to 2023 (% of total enrollment), and predicted trend of AYAs enrollment estimated by interrupted time series analysis (ITSA). The dashed vertical line marks the start of the post-COVID period (2020-2023).



Downloaded for Anonymous User (n/a) at IRCCS San Gerardo dei Tintori
November 20, 2025. For personal use only. No other uses without permission.

In Table 1 socio-demographic, epidemiological, and clinical-therapeutic characteristics of AYAs are reported, as compared with adult participants, at enrollment. Briefly, AYAs were more frequently men-who-have-sex-with-men (MSM) and non-Italian and less frequently Injective Drug Users (IDU). Time elapsed between HIV diagnosis and enrolment in ICONA cohort was similar between the two groups, but the clinical presentation was different: among AYAs, we observed a lower proportion of AIDS-related events and hospitalizations, higher CD4 counts and CD4/CD8 ratios, and lower HIV-

RNA levels. A smaller proportion of AYAs had a positive HCV serostatus. Conversely, no differences were found in terms of acute HIV infection and concomitant diagnosis of HBV co-infection and STIs. The choice of the first ART regimen was similar between the two groups. Finally, the educational and job profiles were found to be different, with a higher percentage of AYAs who attended college and a lower percentage employed, as compared to adults.

Analysis of study outcomes

Antiretroviral therapy initiation

Among 9,519 participants, 7,453 received an HIV diagnosis as of 2015, when universal access to ART was included in Italian national guidelines, with at least one follow-up visit after enrollment or ART initiation. AYAs had a lower cumulative probability of ART start at 12-months [91.9% (91.9-96.0) for AYAs vs 96.3% (95.8-96.7) than adults, log-rank $p < 0.001$] (Supplemental Figure S1A). The Cox regression model confirms the lower crude risk of starting ART of AYAs than adults (HR 0.83; 95% CI 0.73-0.88; $p < 0.001$), but we observed no evidence for differences between the two groups after adjusting for potential predefined confounders (aHR 0.98; 95% CI 0.68-1.08; $p = 0.68$) (Table 2).

Virological failure

Out of a total of 6,637 individuals who had at least two viral load assessments more than 6 months after starting ART, we observed 695 (10.5%) VF: 34 (7.9%) among 430 AYAs and 661 (10.6%) among 6,207 adults, defined as two consecutive HIV-RNA > 50 copies/mL after 6 months of therapy. At the survival analysis, AYAs and adults had a similar cumulative probability of VF [at 1-year 3.1% (1.8-5.2) for AYAs vs 5.1% (4.6-5.7) for adults, log-rank $p = 0.082$] (Supplemental Figure S1B). Similarly, we found no evidence for differences in the risk of VF between the two study groups at both unadjusted and adjusted analyses (HR 0.74; 95% CI 0.52-1.04, $p = 0.083$; aHR 0.97; 95% CI 0.68-1.39, $p = 0.886$) (Table 2).

Treatment discontinuation

Overall, 4,992 (52.4%) study participants (331 AYAs and 4,661 adults) discontinued at least one drug of their first ART regimen for one of the following reasons: treatment simplification, drug

toxicity/intolerance, failure and patient's choice or adherence issues and unknown/other causes. The different distributions of the causes of TD in the two groups and the Kaplan-Meier curves showing the cumulative probabilities of TD: for any causes, for failure, for patient's choice/non-adherence and for toxicity of AYAs vs adults are reported in Supplemental Figures 1 and 2C-F. In particular, AYAs reported a higher proportion of TD for patient's choice/non-adherence (6.9% vs 3%; $p < 0.001$), and a lower proportion of TD for failure (2.7% vs 5.1%; $p = 0.044$), compared to adults. At the unadjusted and adjusted Cox regression analysis, the risk of TD for any causes was similar between the two study groups (HR 1.03, 95% CI 0.92-1.15, $p = 0.569$; aHR 1.10, 95% CI 0.98-1.23, $p = 0.120$), as well as the risk of TD for failure (HR 0.55; 95% CI 0.28-1.07, $p = 0.078$; aHR 0.75; 95% CI 0.37-1.53, $p = 0.426$); the risk for TD for patient's choice/non-adherence of AYAs was instead confirmed to be higher than in adults (HR 2.34; 95% CI 1.51-3.64, $p < 0.001$; aHR 2.21; 95% CI 1.40-3.49, $p = 0.001$) (Table 2). Finally, we investigated the risk of discontinuation of the first ART regimen for toxicity and again there was no evidence for a difference between AYAs and adults (HR 0.81; 95% CI 0.59-1.10, $p = 0.175$; aHR 0.96; 95% CI 0.70-1.32, $p = 0.81$) (Table 2), even when analyzed according to the class of the first ART regimen (p -value interaction=0.826).

Loss to follow up


The risk of LTFU was assessed by restricting the data derived from 16 centers of the ICONA network with active update of follow-up visits over the period 2016-2023; a total of 2,266 study participants was included for the analysis of this outcome. After applying the definition of being lost to follow-up detailed in the method section, we observed a total of 606 (26.7%) LTFU, 71 among 162 AYAs (43.8%) and 535 LTFU among 2,104 adults (25.4%) ($p < 0.001$), mostly distributed in 2019 and 2022 (Supplemental Figures 3). At the survival analysis, AYAs had a higher cumulative probability of LTFU than adults [at 1-year 23.9% (17.7-31.7%) for AYAs vs 12.6% (7.8-10.4) for adults, log-rank $p < 0.001$] (Supplemental Figure S1G). The higher risk of LTFU was observed among AYAs, even after controlling for several potential confounding factors (HR 1.90; 95% CI 1.48-2.43, $p < 0.001$; aHR 1.49; 95% CI 1.15-1.92, $p = 0.002$) (Table 2). Restricting the definition of definitive LTFU to

individuals with no other follow-up visit recorded since their last visit, we found a total of 359 (15.8%) definitive LTFU, 42 among 162 AYAs (25.9%) and 317 (15.1%) among 2,104 adults ($p < 0.001$). Again, AYAs showed a higher cumulative probability of definitive LTFU than adults [at 1-year 15.7% (10.7-22.7) for AYAs vs 7.4% (6.3-8.7) for adults, log-rank $p < 0.001$] (Supplemental Figure S1H) confirmed at the unadjusted analysis and, marginally, a 38% higher risk after adjusting for the same set of covariates (HR 1.75; 95% CI 1.27-2.42, $p < 0.001$; aHR 1.38; 95% CI 0.99-1.92, $p = 0.055$) (Table 2).

Immunological response

A total of 6,900 study participants had at least two CD4 count determinations, the first at baseline (ART starting) and the other during the following 24 months. The estimated mean change of CD4 count by the linear mixed model from baseline to each time point after starting ART was significantly higher among AYAs compared to adults, after controlling for gender, calendar year of ART starting and class of first ART regimen as well as the trends of CD4 recovery overtime ($p < 0.001$). The test for the three-way interaction (age group \times time \times ART class) was statistically significant ($p = 0.008$), suggesting that the effect of age (AYAs vs adults) on CD4 recovery over time differed across the classes of first line regimen (INSTI vs non-INSTI class). Indeed, a wider increase in CD4 count from baseline to the other 3 timepoints was observed comparing AYAs with adults of the INSTI group (Figure 2, Table 3).

Figure 2. Immunological response in adolescents and young adults (AYAs) vs adults after initiation of the first-line antiretroviral therapy (ART) - linear mixed model adjusted for gender, calendar year of ART starting, and class of first ART regimen (Integrase Strand Transfer Inhibitors – INSTI – vs non-INSTI class) with random intercept and slope per patient.



Downloaded for Anonymous User (n/a) at IRCCS San Gerardo dei Tiri
November 20, 2025. For personal use only. No other uses without permission.

Discussion

This study describes the epidemiological, socio-behavioural, clinical and viro-immunological characteristics of AYAs (Adolescents and Young Adults aged 18-24) enrolled in the Italian ICONA cohort in the period 2014-2023 and compares this population with adults. At the best of our knowledge, it represents the first wide-scale description of AYAs living with HIV in Italy.

The comparison among the two populations shows a higher number of MSM and foreign-born among AYAs, and a higher number of IDUs and HCV-Ab positive among adults, reflecting the demographic evolution occurring among PWH in Italy. Indeed, MSM have progressively become the most represented population over the recent years in Italy, especially among younger patients, while the number of IDUs among PWH has progressively decreased [2]. Moreover, the different rate of IDU may explain the difference in HCV *serostatus*, considering that the use of injective drugs is the most frequent mode of HCV transmission [13]. In the meanwhile, in recent years, an increasing number of foreign-born, mostly belonging to the younger age groups, settled in Italy, and this is also reflected in the Italian epidemiological scenario as well as in the ICONA cohort [2]. The other significant difference between the AYAs and adults' groups is the severity of the clinical picture at enrollment;

it is less advanced among AYAs, with a lower rate of AIDS events and hospitalization, a higher CD4 count and a lower level of HIV-RNA. These differences are probably due to the presumable shorter duration of infection, to the more preserved immune system among AYAs [14], and to the higher prevalence of the population with awareness about HIV risk, such as MSM, who undergoes more frequently to HIV testing, thus favouring early detection of the disease. Interestingly, the physicians' choice of the first ART regimen was similar between the two study groups, in accordance with current national and international guidelines.

Overall, the results on the proportion of AYAs enrolled in ICONA suggest a transient impact associated with the COVID-19 pandemic period, without significant modifications in the subsequent trend compared to pre-COVID years. There are no unequivocal reasons to explain this variation. The drop in the proportion of AYAs observed after the onset of the COVID-19 pandemic may be related to the pandemic itself. Indeed, during those years, access to HIV testing was significantly hampered for multiple reasons—including health system strain, lockdowns, and reduced outreach services—leading to a preferential diagnosis of more severe and symptomatic cases, which are more frequent among adult patients. [15]

With regard to the first key outcome of the study, we found no differences between AYAs and adults, enrolled as of 2015, in the time to cART initiation, a result that reflects the universal recommendation to start cART as soon as possible after HIV diagnosis, regardless of CD4 cell count. Similarly, no differences emerged in the time to VF and to first TD, but some differences occurred about reasons for TD: among AYAs, discontinuation were less frequent for VF and more frequent for patients' choice and/or adherence issues. The latter result may be due to disadvantaged social and economic situation among AYAs, and to a difficult access to care due to administrative and legal barriers, especially among foreign-born individuals, and these difficulties may contribute to their poor adherence to ART, as their main focus is not addressing health issues. Moreover, significant differences emerged about LTFU rates, with AYAs significantly at higher risk of being LTFU. This finding was confirmed independently by the definition used to measure LTFU rates. The higher

mobility for educational and working reasons may justify this result among AYAs. In particular, young foreign-born individuals often change location or move to other countries during their migration project, looking for better opportunities, and the high presence of foreign-born among our AYAs population may have strongly contributed to this finding [16-17]. This reason may explain the lower rates of LTFU in 2020-2021, when lockdown was in force in Italy, such reflecting limitations in mobility. Finally, as the overmentioned baseline *status*, immunological response in AYAs after initiation of the first-line ART was stably and significantly higher at 6, 12 and 24 months in AYAs than in adults. This finding is more remarkable among those using INSTI. The presumable shorter duration of HIV infection, as well as a better thymic function, with a stable production of naïve T cells, a lower immune-activation and systemic inflammation comorbidity-driven, and the higher CD4 nadir may explain the faster and robust recovery of CD4 [18]. The more rapid virological suppression induced by INSTI may contribute to better immunological recovery among those using this drug class [19].

Downloaded for Anonymous User (n/a) at IRCCS San Gerardo dei Tiri
November 20, 2025. For personal use only. No other uses without permission.

Most of the results from our analyses are consistent with similar studies performed elsewhere in the world. Although our study does not specifically measure adherence, the greater number of TD related to adherence problems is indicative of how common this issue is among AYAs and deserves special attention. A lower adherence and a higher risk of TD in AYAs have already been described in studies from the United States, Australia and South Africa [4-7]. A large systematic review and meta-analysis including 50 studies found an overall suboptimal adherence to ART in AYAs, as low as 62.3% [20]. Similarly, and due to a likely common greater difficulty of retention in care, higher LTFU rates have been reported among AYAs in the United States, Australia and Perù [4,6,7,9,21,22]. On the other hand, overall the Continuum of Care (CoC) among AYAs included in our population was good: time to ART initiation, total TD and rates of virological suppression were not significantly different in the two study groups. Consequently, the CoC is better preserved than those described in other similar studies: inconsistent care among AYAs, with lower rates of viral suppression, has been reported from Nigeria, USA, Canada, South Africa and Perù [4-6,9, 23,24]. Of note, some of these studies include

patients perinatally infected, too. These patients, excluded from our cohort, which includes only ART-naïve PWH >18 years old, experienced the critical transition from paediatric to adult services, therefore their results about TD and LTFU rates cannot be entirely comparable with ours, since the populations are different. Similar barriers in adherence and retention have been documented in other chronic conditions affecting adolescents and young adults, such as tuberculosis (TB), diabetes, and mental health disorders. In these settings, common critical points include the transition from family-supported to autonomous care, competing educational or social priorities, limited awareness of long-term health consequences, and stigma. Lessons from multidisciplinary and youth-friendly models developed in these areas - such as integrated TB/HIV services, peer-support programs, and community-based adherence support - may help in developing tailored strategies for AYAs living with HIV [25-28].

This study has some limitations, including the observational nature of the data source, which is prone to bias due to unmeasured confounders, and the incomplete information available about adherence and other factors such as comorbidities and/or concomitant medications. Moreover, we cannot know if those LTFU in our cohort are effectively lost or in care in another clinical centre not included in the ICONA cohort.

Nonetheless, this study has a number of remarkable strengths. Firstly, it is the first description of the real-life situation of AYAs with HIV infection in Italy. Moreover, given the wide diffusion of centres participating to ICONA in Italy, the results of this study should be considered as representative of the Italian population of HIV-infected AYAs. On the contrary, these results should not be generalized for other European countries, as many factors, including the differences among health systems and the proportion of foreign-born among AYAs, can strongly influence the study outcomes.

In conclusion, the results of this study highlight that AYAs with HIV have special needs, particularly regarding the adherence to ART and the risk of being LTFU. Less social and economic stability, and increased mobility for educational and working reasons, might contribute to these findings. Specific interventions, such as rapid linkage to care, immediate ART start, “ad-hoc” clinical pathways and

supportive care are essential to enhance awareness among AYAs about their health needs and increase retention in care, especially among those with higher risk of being LTFU and to experience VF, like young foreign-born AYAs.

Declaration of competing interests

FMF received honoraria from ViiV Healthcare and Gilead Sciences for advisory board and support for travel and arrangement for congresses and meetings from ViiV Healthcare, GSK, Gilead Sciences, MSD. IM reports payments to their institution from Gilead Sciences, speakers' honoraria/educational activities for ViiV Healthcare and Gilead Sciences and advisor for Gilead Sciences. EQR received travel grants from Gilead Sciences and ViiV Healthcare. Ace received consultancy fees from ViiV Healthcare and Gilead Sciences. ACo received honoraria from Gilead Sciences for Advisory Board. ACi received funding for scientific advisory boards, travel, or speaker honoraria from Gilead Sciences, ViiV Healthcare, Janssen - Cilag, and MSD. EG received a research grant from Gilead Sciences and speaker fees from ViiV healthcare and Gilead Science. AA served as a paid consultant to Astra Zeneca, Bavarian Nordic, Gilead Sciences, GSK, Janssen-Cilag, MSD, Moderna, Pfizer, and ViiV Healthcare and received institutional research grants from Astra Zeneca, Gilead Sciences and ViiV Healthcare. FV, AS, LC, MET, AdM and AT declare no competing interests.

Acknowledgements

Funding

No specific funding for this project. The Icona Foundation is supported by unrestricted grants from Gilead Sciences, ViiV Healthcare and Merck Sharpe & Dohme. The funders of the ICONA Foundation had no role in the study design, data collection, analysis, decision to publish, or preparation of this study. The research contribution of Mastroianni I, Girardi E and Antinori A was partially granted by the Ricerca Corrente of the Italian Ministry of Health, Linea 2-progetto 2 (INMI L. Spallanzani IRCCS).

Acknowledgements**ICONA FOUNDATION STUDY GROUP**

BOARD OF DIRECTORS: A d'Arminio Monforte (President), A Antinori (Vice-President), S Antinori, A Castagna, M Cernuschi, L Cosmaro, A Di Biagio, E Dibrino, E Girardi, S Lo Caputo, GC Marchetti, C Mussini, E Quiros-Roldan, L Sarmati, B Suligo, C Torti.

SCIENTIFIC SECRETARIAT: A d'Arminio Monforte, A Antinori, A Castagna, F Ceccherini-Silberstein, A Cingolani, A Cozzi-Lepri, A Di Biagio, R Gagliardini, A Giacomelli, E Girardi, A Gori, S Lo Caputo, G Marchetti, F Maggiolo, C Mussini, M Puoti, CF Perno, A Tavelli.

STEERING COMMITTEE: A Antinori, A Bandera, S Bonora, A Calcagno, A Castagna, F Ceccherini-Silberstein, A Cervo, A Cingolani, P Cinque, A Cozzi-Lepri, A d'Arminio Monforte, A De Vito, A Di Biagio, A Fortunato, R Gagliardini, A Giacomelli, E Girardi, N Gianotti, A Gori, G Guaraldi, S Lanini, G Lapadula, M Lichtner, A Lai, S Lo Caputo, G Madeddu, F Maggiolo, V Malagnino, G Marchetti, A Mondì, V Mazzotta, C Muccini, C Mussini, S Nozza, CF Perno, S Piconi, C Pinnetti, MC Poliseno, M Puoti, E Quiros Roldan, R Rossotti, S Rusconi, MM Santoro, A Saracino, L Sarmati, V Spagnuolo, N Squillace, V Svicher, L Taramasso, C Tincati, C Torti, A Vergori.

STATISTICAL AND MONITORING TEAM: A Cozzi-Lepri, S De Benedittis, I Fanti, M Giotta, S Lanini, N Lentini, M Muccio, R Pastorino, A Rodano', A Roen, A Tavelli.

COMMUNITY ADVISORY BOARD: S Bazzichetto, M Cernuschi, L Cosmaro, A Perziano, V Calvino, D Russo, M Farinella, N Policek, VL Del Negro.

BIOLOGICAL BANK INMI AND SAN PAOLO: M Augello, S Carrara, S Graziano, G Prota, H Sadeghi, S Truffa, D Vincenti, R Rovito.

PARTICIPATING PHYSICIANS AND CENTERS: Italy A Giacometti, A Costantini, V Barocci (Ancona); A Saracino, C Santoro, E Milano (Bari); L Comi, C Suardi (Bergamo); P Viale, L Badia, S Cretella (Bologna); EM Erne, A Pieri (Bolzano); E Quiros Roldan, E Focà (Brescia); B Menzaghi, C Abeli (Busto Arsizio); L Chessa, F Pes (Cagliari); P Maggi, L Alessio (Caserta); G Nunnari, BM Celesia (Catania); J Vecchiet, K Falasca (Chieti); A Pan, S Dal Zoppo (Cremona); D Segala (Ferrara);

A Bartoloni, B Borch, C Costa, A Castellaccio (Firenze); S Lo Caputo, A Narducci (Foggia); M Bassetti, E Pontali, S Bianchi, N Bobbio (Genova); C. Del Borgo, R. Marocco, G. Mancarella (Latina); S Piconi, C Molteni (Lecco); S Rusconi, G Canavesi (Legnano); G Pellicanò, Russotto Y (Messina); G Marchetti, S Antinori, A Gori, M Puoti, A Castagna, A Bandera, V Bono, MV Cossu, A Giacomelli, R Lolatto, MC Moioli, L Pezzati, S Diotallevi, C Tincati (Milano); C Mussini, M Menozzi (Modena); P Bonfanti, G Lapadula (Monza); V Sangiovanni, I Gentile, V Esposito, N Coppola, FM Fusco, G Di Filippo, V Rizzo, N Sangiovanni, S Martini (Napoli); AM Cattelan, D Leoni (Padova); A Cascio, M Trizzino (Palermo); D Francisci, E Schiaroli (Perugia); G Parruti, F Sozio (Pescara); D Messeri, SI Bonelli (Pistoia); C Lazzaretti, R Corsini (Reggio Emilia); A Antinori, R Cauda, G D'Ettore, L Sarmati, A Latini, A Cingolani, I Mastrosera, S Lamonica, M Capozzi, M Camici, I Mezzaroma, M Rivano Capparuccia, G Iaiani, C Stingone, L Gianserra, J Paulicelli, MM Plazzi, L Bortolani, M Fusto (Roma); I Coledan (Rovigo); G Madeddu, A De Vito (Sassari); M Fabbiani, F Montagnani (Siena); A Franco, R Fontana Del Vecchio (Siracusa); D Francisci, C Di Giuli (Terni); GC Orofino, G Calleri, G Di Perri, S Bonora, D Meloni (Torino); C Tascini, A Londero (Udine); G Battagin, S Nicolè (Vicenza); G Starnini, S Dell'Isola (Viterbo).

Downloaded for Anonymous User (r/a) at IRCCS San Gerardo dei Tintori, November 20, 2025. For personal use only. No other uses without permission.

Authors' contributions

Conception: FMF, IM, FV, AT; Study Design: FMF, IM, FV, EG, AA, ADM, AT; Accessing and verifying data: IM, AT; Statistical Analysis: AT; Acquisition of data: FMF, IM, FV, ACe, EQR, ACo, ACi; Draft of the manuscript: FMF, IM, AT; Patients' enrolment: FMF, IM, FV, ACe, AS, LC, EQR, MET, ACo, ACi, AA; Review of the article and critical revision for important intellectual content: all the authors; Reading and final approval of the submitted version: all the authors.

Data Availability Statement

The datasets generated during the current study are not publicly available because they contain sensitive data that must be treated under data protection laws and regulations. An appropriate data sharing agreement can be arranged after a reasonable request to the corresponding author.

Ethics statement

All the enrolled subjects provided a written informed consent for study participation and processing of personal data. The ICONA Foundation study protocol and the informed consent were approved by the Institutional Review Board (IRB) of each participating centre. The latest amendment of the ICONA Foundation Study was approved centrally by the Lazio Area 4 Territorial Ethics Committee on 01 July 2024 (approval no. 83-2024). All procedures of the study were performed in accordance with the 1964 Helsinki declaration and its later amendments

References

1. European Centre for Disease Prevention and Control/WHO Regional Office for Europe. HIV/AIDS Surveillance in Europe 2024 – 2023 data. Stockholm: ECDC; 2024. Available from: <https://www.ecdc.europa.eu/en/publications-data/hiv-aids-surveillance-europe-2024-2023-data>. Accessed Nov 27, 2024.
2. Centro Operativo AIDS. Sorveglianza HIV/AIDS 2024 (dati 2023). Not Ist Super Sanità. 2024;37(11):3-59. Available from: <https://www.iss.it/documents/20126/0/Notiziario+COA+casi+HIV+e+AIDS+2023.pdf/8b1aaae1-d985-b6bc-bc67-42be27e32886?t=1732285696251>. Accessed Nov 27, 2024.
3. Centers for Disease Control and Prevention. HIV Surveillance Report: Estimated HIV incidence and Prevalence in the United States, 2018-2022. Available from: https://stacks.cdc.gov/view/cdc/156513/cdc_156513_DS1.pdf
4. Beer L, Mattson CL, Shouse RL, Prejean J. Receipt of clinical and prevention services, clinical outcomes, and sexual risk behaviors among HIV-infected young adults in care in the United States. *AIDS Care*. 2016;28(9):1166-1170. doi:10.1080/09540121.2016.1160028.
5. Haghightat R, Toska E, Bungane N, Cluver L. The HIV care cascade for adolescents initiated on antiretroviral therapy in a health district of South Africa: a retrospective cohort study. *BMC Infect Dis*. 2021;21(1):60. doi:10.1186/s12879-020-05742-9.

Downloaded for Anonymous User (n/a) at IRCCS San Gerardo dei Tiri
November 20, 2025. For personal use only. No other uses without permission.

6. Beer L, Mattson CL, Bradley H, Shouse RL; Medical Monitoring Project. Trends in ART Prescription and Viral Suppression Among HIV-Positive Young Adults in Care in the United States, 2009-2013. *J Acquir Immune Defic Syndr.* 2017;76(1):e1-e6. doi:10.1097/QAI.0000000000001427.
7. Hughes C, Puhr R, Ojaimi S, et al. Human immunodeficiency virus-infected young people in Australia: data from the Australian HIV Observational Database. *Intern Med J.* 2018;48(12):1447-1456. doi:10.1111/imj.14040.
8. Crepaz N, Dong X, Hess KL, Bosh K. Brief Report: Racial and Ethnic Disparities in Sustained Viral Suppression and Transmission Risk Potential Among Persons Aged 13-29 Years Living With Diagnosed HIV Infection, United States, 2016. *J Acquir Immune Defic Syndr.* 2020;83(4):334-339. doi:10.1097/QAI.0000000000002277.
9. Niwa M, Matos E, Rodriguez C, et al. Retention and Viral Suppression Among Adolescents Newly Initiating Antiretroviral Therapy in Adult HIV Care in Lima, Peru: A Retrospective Cohort Study. *J Adolesc Health.* 2024;74(2):260-267. doi:10.1016/j.jadohealth.2023.08.041.
10. Meloni ST, Agaba P, Chang CA, et al. Longitudinal evaluation of adherence, retention, and transition patterns of adolescents living with HIV in Nigeria. *PLoS One.* 2020;15(7):e0236801. doi:10.1371/journal.pone.0236801.
11. HIVinfo.NIH.Gov. HIV and specific population. HIV and Adolescents and young adults. Available from: <https://hivinfo.nih.gov/understanding-hiv/fact-sheets/hiv-and-adolescents-and-young-adults>.
12. Vourli G, Pharris A, Cazein F, et al. Are European HIV cohort data within EuroCoord representative of the diagnosed HIV population? *AIDS.* 2019 Jan 27;33(1):133-143. doi:10.1097/QAD.0000000000002034.
13. Trickey A, Fraser H, Lim AG, et al. The contribution of injection drug use to hepatitis C virus transmission globally, regionally, and at country level: a modelling study. *Lancet Gastroenterol Hepatol.* 2019 Jun;4(6):435-444. doi:10.1016/S2468-1253(19)30085-8.

14. Balestre E, Eholié SP, Lokossue A, et al.; International epidemiological Database to Evaluate AIDS (IeDEA) West Africa Collaboration. Effect of age on immunological response in the first year of antiretroviral therapy in HIV-1-infected adults in West Africa. *AIDS*. 2012 May 15;26(8):951-7. doi:10.1097/QAD.0b013e3283528ad4.
15. Rick F, Odoke W, van den Hombergh J, et al. Impact of coronavirus disease (COVID-19) on HIV testing and care provision across four continents. *HIV Med*. 2022 Feb;23(2):169-177. doi:10.1111/hiv.13180.
16. Saracino A, Lorenzini P, Lo Caputo S, et al. Increased risk of virologic failure to the first antiretroviral regimen in HIV-infected migrants compared to natives: data from the ICONA cohort. *Clin Microbiol Infect*. 2016;22(3):288.e1-8. doi:10.1016/j.cmi.2015.10.026.
17. Crockett KB, Schember CO, Bian A, et al. Relationships between patient race and residential race context with missed human immunodeficiency virus care visits in the United States, 2010-2015. *Clin Infect Dis*. 2023;76(12):2163-2170. doi:10.1093/cid/ciad069.
18. Chen J, Titanji K, Sheth AN, et al. The effect of age on CD4⁺ T-cell recovery in HIV-suppressed adult participants: a sub-study from ACTG A5321 and the Bone Loss and Immune Reconstitution (BLIR) study. *Immun Ageing*. 2022;19:4. doi:10.1186/s12979-022-00260-0.
19. Jacobson K, Ogbuagu O. Integrase inhibitor-based regimens result in more rapid virologic suppression rates among treatment-naïve human immunodeficiency virus-infected patients compared to non-nucleoside and protease inhibitor-based regimens in a real-world clinical setting: A retrospective cohort study. *Medicine (Baltimore)*. 2018 Oct;97(43):e13016. doi:10.1097/MD.00000000000013016.
20. Kim SH, Gerver SM, Fidler S, Ward H. Adherence to antiretroviral therapy in adolescents living with HIV: systematic review and meta-analysis. *AIDS*. 2014 Aug 24;28(13):1945-56. doi:10.1097/QAD.0000000000000316.
21. Althoff KN, Rebeiro P, Brooks JT, et al.; North American AIDS Cohort Collaboration on Research and Design (NA-ACCORD). Disparities in the quality of HIV care when using US

- Department of Health and Human Services indicators. *Clin Infect Dis*. 2014 Apr;58(8):1185-9. doi:10.1093/cid/ciu044.
22. Nachega JB, Hislop M, Nguyen H, et al. Antiretroviral therapy adherence, virologic and immunologic outcomes in adolescents compared with adults in southern Africa. *J Acquir Immune Defic Syndr*. 2009 May 1;51(1):65-71. doi:10.1097/QAI.0b013e318199072e.
23. Ahonkhai AA, Banigbe B, Adeola J, et al. Age Matters: Increased Risk of Inconsistent HIV Care and Viremia Among Adolescents and Young Adults on Antiretroviral Therapy in Nigeria. *J Adolesc Health*. 2016 Sep;59(3):298-304. doi:10.1016/j.jadohealth.2016.05.002.
24. Palmer A, Gabler K, Rachlis B, et al.; Canadian Observational Cohort (CANOC) Collaboration. Viral suppression and viral rebound among young adults living with HIV in Canada. *Medicine (Baltimore)*. 2018 Jun;97(22):e10562. doi:10.1097/MD.00000000000010562.
25. Leddy AM, Ricks P, Kates J, Starke JR, Cruz AT. Social determinants of adherence to treatment for children and adolescents/young adults with tuberculosis infection and disease: a narrative review. *J Pediatr Infect Dis Soc*. 2022;11(Suppl 3):S79-S86. doi:10.1093/jpids/piac070.
26. Chiang SS, Tanzer JR, Starke JR, Mandalakas AM. Identifying adolescents at risk for suboptimal adherence to tuberculosis treatment: a prospective cohort study. *PLoS Glob Public Health*. 2024;4(2):e0002918. doi:10.1371/journal.pgph.0002918.
27. Ude A-O, Alhassan D, Hill AJ, et al. Transitioning to adult care in youth-onset diabetes: a scoping review. *BMC Public Health*. 2025;25:1476. doi:10.1186/s12889-025-22956-1.
28. Waumans RC, van der Veen DC, Keijzer H, et al. Understanding and preventing non-adherence to and dropout from treatment for depressive and anxiety disorders in adolescents and young adults. *Front Psychiatry*. 2023;14:1174285. doi:10.3389/fpsy.2023.1174285.

Table 1. Socio-demographic, epidemiological, and clinical-therapeutic characteristics of individuals enrolled in the ICONA cohort between 2014 and 2023, by age group (adolescents and young adults – AYAs - 18-24 years vs adults ≥ 25 years)

	Adults 8,866 (93.1%)	AYAs 653 (6.9%)	Total 9,519 (100.0%)	p
Age, median [IQR]	41 [33-50]	23 [21-24]	39 [31-49]	<0.001
Sex at birth, N (%)				
<i>Female</i>	1,628 (18.4%)	137 (21.0%)	1,765 (18.5%)	0.097
<i>Male</i>	7,238 (81.6%)	516 (79.0%)	7,754 (81.5%)	
Country of birth, N (%)				
<i>Foreign-born</i>	2,333 (26.3%)	277 (42.4%)	2,610 (27.4%)	<0.001
<i>Italy-born</i>	6,533 (73.7%)	376 (57.6%)	6,909 (72.6%)	
Risk factor for HIV acquisition, N (%)				
<i>MSM contacts</i>	4,256 (48.0%)	391 (59.9%)	4,647 (48.8%)	<0.001
<i>Heterosexual contacts</i>	3,502 (39.5%)	198 (30.3%)	3,700 (38.9%)	
<i>IDU</i>	490 (5.5%)	25 (3.8%)	515 (5.4%)	
<i>Other/Unknown</i>	618 (7.0%)	39 (6.0%)	657 (6.9%)	
Months from HIV diagnosis to enrollment, median [IQR]	0.4 [0.1-1.1]	0.5 [0.1-1.2]	0.5 [0.1-1.1]	0.591
Year of enrollment, median [IQR]	2018 [2015-2021]	2017 [2016-2019]	2017 [2015-2020]	0.049
Acute HIV infection, N (%)	154 (1.7%)	15 (2.3%)	169 (1.8%)	0.296
Italian geographic area, N (%)				
<i>Central</i>	3,101 (35.0%)	208 (31.9%)	3,309 (34.8%)	0.103
<i>Northern</i>	4,214 (47.5%)	312 (47.8%)	4,526 (47.5%)	
<i>Southern/Islands</i>	1,551 (17.5%)	133 (20.4%)	1,684 (17.7%)	
Job, N (%)				
<i>Unemployed</i>	1,143 (12.9%)	134 (20.5%)	1,277 (13.4%)	<0.001
<i>Employed</i>	5,674 (64.0%)	356 (54.5%)	6,030 (63.3%)	
<i>Unknown</i>	2,049 (23.1%)	163 (25.0%)	2,212 (23.2%)	
Education, N (%)				
College	2,306 (26.0%)	226 (34.6%)	2,532 (26.6%)	<0.001
Primary School	301 (3.4%)	26 (4.0%)	327 (3.4%)	
Secondary School	1,145 (12.9%)	65 (10.0%)	1,210 (12.7%)	
University	1,124 (12.7%)	84 (12.9%)	1,208 (12.7%)	
Unknown	3,990 (45.0%)	252 (38.6%)	4,242 (44.6%)	
Smoker, n (%)	3,329 (43.0%)	261 (45.6%)	3,590 (43.2%)	0.223
AIDS at diagnosis, N (%)	1,320 (14.9%)	28 (4.3%)	1,348 (14.2%)	<0.001
Hospitalization at enrollment, N (%)	1,404 (15.8%)	38 (5.8%)	1,442 (15.1%)	<0.001
HCV-Ab positive, N (%)	615 (7.7%)	31 (5.2%)	646 (7.5%)	0.025
HBsAg positive, N (%)	365 (4.6%)	21 (3.5%)	386 (4.5%)	0.242
STI at enrolment, N(%)	874 (9.9%)	56 (8.6%)	930 (9.8%)	0.549
CD4 at enrollment, cells/mm³, median [IQR]	329 [121-527]	460 [311 -48]	339 [134-534]	<0.001

Downloaded for Anonymous User (n/a) at IRCCS San Gerardo dei Tintori, November 20, 2023. For personal use only. No other uses without permission.

CD4 at enrollment, cells/mm³, N (%)				
<200 CD4/mm ³	2,748 (33.9%)	62 (10.4%)	2,810 (32.3%)	<0.001
200-349 CD4/mm ³ , N (%)	1,532 (18.9%)	126 (21.1%)	1,658 (19.0%)	
350-499 CD4/mm ³ , N (%)	1,589 (19.6%)	142 (23.8%)	1,731 (19.9%)	
≥500 CD4/mm ³ , N (%)	2,241 (27.6%)	266 (44.6%)	2,507 (28.8%)	
HIV-RNA at enrollment, log₁₀ cps/ml, median [IQR]	4.9 [4.3-5.5]	4.7 [4.1-5.3]	4.9 [4.2-5.5]	<0.001
HIV-RNA at enrollment, log₁₀ cps/ml, median [IQR]				
<100.000 cps/mL	4,433 (54.5%)	401 (66.6%)	4,834 (55.3%)	<0.001
100.000-500.000 cps/mL	2,162 (26.6%)	124 (20.6%)	2,286 (26.2%)	
>500.000 cps/mL	1,544 (19.0%)	77 (12.8%)	1,621 (18.5%)	
CD4/CD8 ratio at enrollment, median [IQR]	0.4 [0.2-0.6]	0.5 [0.3-0.8]	0.4 [0.2-0.6]	<0.001
First line ART, N (%)				0.356
3DR-INSTI	5,245 (62.7%)	362 (60.1%)	5,607 (62.5%)	
3DR-NNRTI	1,150 (13.7%)	90 (15.0%)	1,240 (13.8%)	
3DR-PI	1,184 (14.1%)	78 (13.0%)	1,262 (14.1%)	
Others	67 (0.8%)	6 (1.0%)	73 (0.8%)	
2DR	444 (5.3%)	39 (6.5%)	483 (5.4%)	
4DR	281 (3.4%)	27 (4.5%)	308 (3.4%)	
Abbreviations: AYAs, adolescents and young adults; IQR, interquartile range; N, number of participants; MSM, men-who-have-sex-with-men; IDU, injective drug users; HCV-Ab, antibodies anti-HCV; STI, sexually transmitted infection; cps, copies; ART, antiretroviral treatment; 3DR, 3 drugs regimen; INSTI, integrase strand transfer inhibitor; NNRTI, non-nucleoside reverse transcriptase inhibitor; PI, protease inhibitor; 2DR, 2 drugs regimen; 4DR, 4 drugs regimen.				

Downloaded for Anonymous User (n/a) at IRCCS San Gerardo dei Tintori, November 20, 2023. For personal use only. No other uses without permission.

Table 2. Hazard Ratios (HR) and Adjusted Hazard Ratios (aHR) of the different study outcomes in adolescents and young adults (AYAs) vs adults enrolled in ICONA cohort in 2014-2023.

Outcomes*	N, (%)	HR (95%CI) (AYAs vs Adults)		aHR (95%CI) (AYAs vs Adults)	
		HR	p	aHR	p
Time to ART starting	7453/7643 (97.5%)	0.83 (0.73-0.88)	<0.001	^[1] 0.98 (0.89-1.08)	0.680
Time to VF	695/6637 (10.5%)	0.74 (0.52-1.04)	0.083	^[2] 0.97 (0.68-1.39)	0.886
Time to TD for any reason	4992/8334 (59.9%)	1.03 (0.92-1.15)	0.569	^[3] 1.10 (0.98-1.23)	0.120

Time to TD for toxicity	794/8334 (9.5%)	0.81 (0.59-1.10)	0.175	^[3] 0.96 (0.70-1.12)	0.81 0
Time to TD for failure	246/8334 (3.0%)	0.55 (0.28-1.07)	0.565	^[3] 0.75 (0.37-1.53)	0.47 0
Time to TD for patient's choice/adherence issues	165/8334 (1.9%)	2.34 (1.51-3.64)	<0.001	^[4] 2.21 (1.40-3.49)	0.00 1
Time to LTFU	606/2266 (26.7%)	1.90 (1.48-2.43)	<0.001	^[5] 1.49 (1.15-1.92)	0.00 2
Time to Definitive LTFU	359/2266 (15.8%)	1.75 (1.27-2.42)	0.001	^[5] 1.38 (0.99-1.92)	0.05 5

*See the main text for outcomes' definitions.

Abbreviations: N, number of participants; HR, hazard ratios; CI, confidence interval; AYAs, adolescents and young adults; vs, versus; aHR, adjusted hazard ratios; ART, antiretroviral treatment; VF, virological failure; TD, treatment discontinuation of at least one drug of the first regimen; LTFU, loss to follow-up.

(1) Adjusted for gender, mode of HIV transmission, italian nation of birth, AIDS diagnosis, acute HIV infection diagnosis, HIV-RNA at enrolment, CD4 count at enrolment, calendar year of HIV diagnosis

(2) Adjusted for gender, mode of HIV transmission, italian nation of birth, AIDS diagnosis, HIV-RNA at ART start, CD4 count at ART start, calendar year of ART start, first ART regimen, **HBV co-infection**

(3) Adjusted for gender, mode of HIV transmission, italian nation of birth, AIDS diagnosis, HIV-RNA at ART start, CD4 count at ART start, calendar year of ART start, first ART regimen

(4) Adjusted for gender, mode of HIV transmission, italian nation of birth, AIDS diagnosis, HIV-RNA at ART start, CD4 count at ART start, calendar year of ART start, first ART regimen, job profile, schooling

(5) Adjusted for gender, mode of HIV transmission, italian nation of birth, calendar year of enrollment in ICONA, job profile, schooling.

Downloaded for Anonymous User (n/a) at IRCCS San Gerardo dei Tintori
November 20, 2025. For personal use only. No other uses without permission.

Table 3. Mean CD4 change in adolescents and young adults (AYAs) vs adults after initiation of the first-line antiretroviral therapy (ART).

	Timepoint	Group	Mean CD4 change from ART start, cells/uL [95%CI]		Contrast, AYAs vs adults (change from ART start, cells/uL) [95%CI]		p-value
Overall	6 months [±2]	Adults	+191	[+180, +201]	+66	[+25, +108]	0.002
		AYAs	+257	[+217, +297]			
	12 months [±3]	Adults	+235	[+229, +242]	+29	[+3, +55]	0.031
		AYAs	+264	[+239, +289]			
	24 months [±4]	Adults	+291	[+284, +298]	+29	[+1, +56]	0.044
		AYAs	+320	[+293, +347]			
	Timepoint	Group	Mean CD4 change from ART start, cells/uL [95%CI]		Contrast, AYAs vs adults (change from ART start, cells/uL) [95%CI]		p-value

INSTI	6 months [±2]	Adults	+193	[+180, +206]	+120	[+65, +175]	<0.001
		AYAs	+313	[+259, +367]			
	12 months [±3]	Adults	+241	[+232, +249]	+57	[+22, +92]	0.001
		AYAs	+298	[+264, +332]			
	24 months [±4]	Adults	+297	[+288, +306]	+55	[+18, +92]	0.004
		AYAs	+351	[+316, +387]			
non-INSTI	6 months [±2]	Adults	+188	[+171, 204]	-4.5	[-67, +57]	0.885
		AYAs	+183	[+123, +243]			
	12 months [±3]	Adults	+226	[+215, +236]	-5	[-45, +35]	0.802
		AYAs	+221	[+182, +259]			
	24 months [±4]	Adults	+281	[+270, +393]	-1.5	[-43, +40]	0.941
		AYAs	+280	[+240, +320]			

Declaration of Interest Statement

Downloaded for Anonymous User (n/a) at IRCCS San Gerardo dei Tiri
November 20, 2025. For personal use only. No other uses without permission.

- The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.
- The author is an Editorial Board Member/Editor-in-Chief/Associate Editor/Guest Editor for this journal and was not involved in the editorial review or the decision to publish this article.
- The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

FMF received honoraria from ViiV Healthcare and Gilead Sciences for advisory board and support for travel and arrangement for congresses and meetings from ViiV Healthcare, GSK, Gilead Sciences, MSD. IM reports payments to their institution from Gilead Sciences, speakers' honoraria/educational activities for ViiV Healthcare and Gilead Sciences and advisor for Gilead Sciences. EQR received travel grants from Gilead Sciences and ViiV Healthcare. Ace received consultancy fees from ViiV Healthcare and Gilead Sciences. ACo received honoraria from Gilead Sciences for Advisory Board. ACi received funding for scientific advisory boards, travel, or speaker honoraria from Gilead Sciences, ViiV Healthcare, Janssen - Cilag, and MSD. EG received a research grant from Gilead Sciences and speaker fees from ViiV healthcare and Gilead Science. AA served as a paid consultant to Astra Zeneca, Bavarian Nordic, Gilead Sciences, GSK, Janssen-Cilag, MSD, Moderna, Pfizer, and ViiV Healthcare and received institutional research grants from Astra

Downloaded for Anonymous User (n/a) at IRCCS San Gerardo dei Tiri
November 23, 2025, 1:00:58 PM. For personal use only. No other uses without permission.

Journal Pre-proof