

Response of the Carbonate System to Ocean Alkalinity Enhancement: A Mesocosm-Scale Experiment

Selene Varliero, Piero Macchi,* Daniela Basso, Paraskevi Pitta, Iordanis Magiopoulos, Chiara Santinelli, Giovanni Checcucci, Silvia Valsecchi, Laura Borromeo, and Guido Raos*



Cite This: <https://doi.org/10.1021/acsestwater.6c00342>



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: We present the results of a mesocosm-scale experiment conducted in 2023 in the eastern Mediterranean Sea (Crete, Greece) to evaluate the effects of ocean liming via the addition of calcium hydroxide. We assessed two dosage regimes through repeated additions—low ($20 \mu\text{mol L}^{-1}$ alkalinity increase per addition, reaching a total theoretical alkalinity increment of $120 \mu\text{mol L}^{-1}$) and high ($180 \mu\text{mol L}^{-1}$ per addition, reaching a total theoretical increment of $1080 \mu\text{mol L}^{-1}$)—comparing against control mesocosms over 14 days, monitoring the pH, total alkalinity, dissolved inorganic carbon, and solid precipitates. The results for the treated mesocosms show that the low dosage effectively increased the total alkalinity and maintained it over the duration of the experiment, unlike the high dosage mesocosms, where the alkalinity dropped within a few days due to the secondary precipitation of carbonates and degassing. In particular, the final net changes of alkalinity were $+110 \mu\text{mol L}^{-1}$ for low dosages (close to the theoretical limit) but amounted to about $-330 \mu\text{mol L}^{-1}$ for the high ones. The concurrent changes in pH were equal to $+0.1$ and $+0.4$, while those in dissolved inorganic carbon were equal to -60 and $-620 \mu\text{mol L}^{-1}$, respectively. The fact that, in the low dosage mesocosms, there was a slight reduction in the dissolved inorganic carbon (comparable to that occurring in control mesocosms) despite the total alkalinity increase suggests limited atmospheric CO_2 uptake. We attribute this to restricted air-sea gas exchange caused by the mesocosm lids, which are commonly adopted in these facilities to prevent the entrance of external contaminants. Our study highlights future directions for testing and enhancing the efficiency of CO_2 uptake by seawater, improving ocean liming as a viable climate mitigation strategy.

KEYWORDS: CO_2 storage, climate change mitigation, marine chemistry, carbonate system



INTRODUCTION

The concentration of atmospheric carbon dioxide (CO_2) has risen dramatically in the last centuries, from 283 ppm in 1800 to 422 ppm in 2024.^{1,2} While CO_2 does not have the highest radiative forcing among the greenhouse gases, its reduction represents a priority since it is the most emitted and persistent one over time. In 2019, it accounted for 64% of global CO_2eq (carbon dioxide equivalent) emissions.³

With these premises, keeping the increase of temperature below 2°C will require the removal of 100 to 1000 GtCO_2 ($= 10^{14}$ – 10^{15} kg of CO_2) from the atmosphere over the course of the 21st century, depending also on the concurrent switch to renewable energies.⁴ These ambitious targets demand the deployment of both carbon dioxide removal (CDR) technologies and effective CO_2 storage solutions.³ Many CDR technologies are currently being developed and tested, but no single technology can fulfill these requirements alone, highlighting the need for a portfolio of solutions.

Ocean alkalinity enhancement (OAE) is a promising method with an estimated CO_2 storage potential ranging


from 1 to 100 $\text{GtCO}_2 \text{ y}^{-15}$ and potential long-term storage for several thousands or even millions of years,^{6–8} although these estimates are largely based on modeling studies. It involves all the species that are part of the carbonate system in seawater: in addition to CO_2 , these include bicarbonate (HCO_3^-) and carbonate (CO_3^{2-}) ions (undissociated carbonic acid being negligible).^{9,10} In OAE, CO_2 is absorbed from the atmosphere and reacts with dissolved alkaline substances, forming CO_3^{2-} and HCO_3^- . This increases both the alkalinity and the dissolved inorganic carbon (DIC) of seawater. Consequently, OAE also contributes to mitigating the ongoing process of ocean acidification by enhancing the oceanic buffer capacity.¹¹

Received: March 9, 2026

Revised: September 3, 2026

Accepted: September 4, 2026

OAE was first proposed by Kheshgi, who suggested employing either calcium oxide (CaO) or calcium hydroxide (Ca(OH)₂).¹² When Ca(OH)₂ (also known as slaked lime, SL) is used, the process can also be defined as ocean liming (OL). In this OAE application, the release of hydroxide ions increases the pH. This shifts the carbonate equilibrium toward CO₃²⁻, creating a CO₂ deficit and inducing atmospheric CO₂ absorption:



Afterward, other studies proposed alternative techniques,¹³ including the usage of other minerals such as olivine (Fe,Mg)₂SiO₄¹⁴ and electrochemical production of alkalinity,¹⁵ all of which have both strengths and weaknesses. In this study, we focus on the original implementation of OL through the use of SL.

Following the initial proposals, some experimental work has been carried out to confirm the storage potential of OAEs.^{16,17} In parallel, research is progressing toward modeling the large-scale deployment of OAEs, with a specific outlook on long-term responses of the ocean to the introduction of large amounts of alkaline substances.^{18,19} Ocean liming may also interfere with natural sources of alkalinity, so its effectiveness depends on the location of its deployment.²⁰ These studies are paving the way for a comprehensive assessment of the feasibility, scalability, and environmental impacts of OAEs.

OL temporarily raises the local pH of seawater, potentially causing the precipitation of carbonate minerals such as aragonite and calcite (two common polymorphs of CaCO₃), as well as brucite (Mg(OH)₂).^{17,21} This phenomenon was defined as “runaway precipitation”,¹⁶ and it is undesirable not only due to its unknown effects on marine biota but also because it reduces process efficiency by lowering the alkalinity and releasing some CO₂ back into the atmosphere.^{9,10} In fact, once triggered by a large spike in pH, precipitation does not stop and may proceed up to a point where the seawater alkalinity is lower than the starting alkalinity, resulting in a negative efficiency.²² To mitigate these effects, research focuses on determining optimal dosages of SL that avoid mineral precipitation in transient high-pH, low-dilution scenarios or seeking alternative alkalization technologies that avoid spikes in pH.^{10,23,24}

This paper presents the results of an experiment conducted in 2023 at the Hellenic Centre for Marine Research (HCMR) in Heraklion (Crete, Greece), aimed at investigating OL at a scale beyond laboratory conditions. The objectives and experimental conditions were defined on the basis of a previous experiment, carried out in 2022 in mesocosms at Estación de Ciencias Mariñas de Toralla (ECIMAT) in Vigo (Spain).^{25,26} The two experiments differed in terms of duration, seawater conditions, and dosage levels, allowing us to establish a robust methodology for mesocosm experiments. Both were part of a multidisciplinary effort, involving several research teams. Mesocosms (see Figure 1) were used, as they provide controlled environments that simulate marine systems, avoiding many of the uncertainties associated with open-sea studies. Given that the eastern Mediterranean seawater is oligotrophic, with characteristically low nutrient concentrations,²⁷ a key objective of the study was also to assess the biological effects of OL on plankton communities, coccolithophores, and chlorophyll dynamics. Results from the biological experiments have been published elsewhere.²⁸ This paper



Figure 1. Mesocosm facility at HCMR, Heraklion, Greece (June 2023).

focuses exclusively on the chemical aspects, with particular attention to the carbonate system.

Our experiments aimed to determine the threshold concentration of Ca(OH)₂ that can trigger runaway precipitation in OL applications. To investigate a broad range of concentrations and assess both short- and medium-term effects within a single set of experiments, Ca(OH)₂ dosages were progressively increased in each mesocosm every 2 days. The experiments were designed to mimic the impacts of SL dissolution in the wake of ships²⁹ before the added alkalinity disperses homogeneously on the ocean surface through the action of currents and winds.

MATERIALS AND METHODS

Mesocosm Facilities and Experimental Setup

The experiment used nine land-based mesocosms hosted in large natural seawater pools that stabilized the structures and maintained the temperature constant.

Each mesocosm had a volume of approximately 3 m³ and a depth of ~1.5 m. The exact volume of water was measured during filling on day 0 using a flow meter. Gentle bubbling was maintained for the entire duration of the experiment to ensure good mixing in the water column, avoiding stratification and facilitating equilibration with air. Specific precautions were taken to prevent accidental nutrient contamination of the oligotrophic seawater. As shown in Figure 1, each mesocosm was covered with a lid, allowing limited air exchange while preventing contamination from exogenous matter (e.g., mineral particles carried by wind, leaves, insects, etc.).

Before the first SL addition, a bottle with a funnel on top was placed at the bottom of each mesocosm to collect the possible precipitates. These bottles were retrieved at the end of the experiment.

The main characteristics of the seawater are summarized in Table 1.

On day 0, the mesocosms were filled with seawater collected close to the HCMR facility. At day 1, the first dosages in the high- and low-concentration mesocosms were applied, and subsequent additions were repeated every 2 days until day 11. Samplings were performed daily at 8:00 AM, while additions were carried out every 2 days,

Table 1. Summary of the Experimental Conditions, Including Average Values and Standard Deviations of Key Seawater Characteristics

Origin of seawater	Eastern Mediterranean
Environment	Oligotrophic
Mesocosm volume at day 0 (m ³)	3.10
Average temperature (°C)	21.05 ± 0.99
Average salinity (psu)	38.1 ± 2.9
Initial TA (μmol L ⁻¹)	2740 ± 33
Initial pH	8.20 ± 0.01
Initial DIC (μmol L ⁻¹)	2456 ± 37

typically 2 h after the morning sampling. On days of addition, an extra sampling was conducted 4 h postaddition.

We performed two types of treatments, which differ by the amount of added Ca(OH)₂. Henceforth, we call them high (H) and low (L) dosage treatments. Each of them was replicated in three mesocosms, enabling the testing of a wider range of values while minimizing the risk of uncontrolled or rapid precipitation. Additionally, three mesocosms were not treated with Ca(OH)₂ and used as the control (C). These mesocosms, and the samples extracted from them, are labeled H1–H3, L1–L3, and C1–C3, respectively.

Each dosage was administered in six consecutive additions over 2 weeks. The alkalinities added to each mesocosm are reported in Table 2. Note that the values in the table are theoretical; hence, we denote

Table 2. Total Added Alkalinity (ΔTA_{th}) Reached at Each Day in the L and H Mesocosms, in μmol L⁻¹

Addition no.	Day	ΔTA _{th} (H)	ΔTA _{th} (L)
1	1	180	20
2	3	360	40
3	5	540	60
4	7	720	80
5	9	900	100
6	11	1080	120

them by the symbol ΔTA_{th}. They are equal to twice the nominal concentration of added Ca(OH)₂, assuming its complete dissociation and neglecting any side reactions. The highest tested TA dosage was 180 μmol L⁻¹, a value determined from preliminary laboratory experiments conducted at HCMR a few days before the start of the mesocosm study. These experiments were performed in 10 L containers using single additions of Ca(OH)₂ at different dosage levels, selected on the basis of calculations considering the local seawater chemistry in order to avoid runaway carbonate precipitation. It is important to highlight that the seawater conditions in Crete, characterized by high values of temperature, salinity, and Ca²⁺ concentration, favor calcium carbonate precipitation. Therefore, we chose to apply only modest increases in alkalinity, compared to other studies in the literature.^{16,21,22}

Ca(OH)₂ was kindly provided by Unicalce with a declared purity of 93.96%. It was added in the form of a 1.5 M slurry, prepared with the water of each mesocosm collected during the morning sampling on the same day. Calcium hydroxide was weighed in at the proper amount to be added to each mesocosm and adjusted to account for their different volumes. The slurry was homogenized by mixing seawater and Ca(OH)₂ in a capped glass bottle and vigorously shaking it by hand for approximately 1 min. Immediately after pouring, the contents of the mesocosms were mixed by drastically increasing air bubbling at the surface for about 1 min.

Sample Handling and Measurements

Water sampling from the mesocosms was performed by using a simple low-disturbance system. From the beginning of the experiment, each mesocosm was equipped with a small sampling tube permanently installed in the water column, positioned approximately at mid-depth.

Samples were collected by applying suction with a syringe, allowing seawater to be withdrawn without disturbing the mesocosm environment, and transferred directly into polycarbonate bottles. To minimize contamination, each mesocosm was assigned a dedicated syringe. In addition, the first 30 s of flow was discarded to avoid collecting stagnant water inside the sampling tube, ensuring that only the circulating mesocosm water was sampled.

After the sampling, the water was filtered through a 0.2 μm poly(ether sulfone) membrane. A first set of measurements took place on site, while part of the samples were stored at 4 °C in glass bottles and transported inside a cooler to Politecnico di Milano for subsequent analyses.

On-Field Measurements. At HCMR, the pH was monitored both with field probes and with the bench instrument Thermo Scientific Orion Star A111 pH meter. The pH was measured both before and after filtration. The pH probe was calibrated according to the NIST scale, and the recorded values were then converted to the total scale,³⁰ accounting also for minor temperature differences between the samples. All pH values reported in this paper are expressed on the total scale. The measured conductivity was used to calculate the practical salinity.³¹ The total alkalinity was measured by automatic titration (Hanna Instruments HI84531).

Laboratory Measurements. At the chemistry laboratory of Politecnico di Milano, pH and conductivity were measured using SevenExcellence electrode sensors (METTLER TOLEDO), applying the same calibration methodology as that used for the field measurements. TA was determined by using the same titration procedure adopted on site. The duplicate measurements of pH and TA performed both in the field and in the laboratory were used to assess possible contamination or sample degradation during storage and transportation. Dissolved inorganic carbon (DIC) could not be measured at HCMR and was measured at Politecnico di Milano by acidification followed by nondispersive infrared detection of produced gaseous CO₂ using an Analytik Jena multi N/C 2100S analyzer.³²

The solid particles collected at the end of the experiments were dried and analyzed by X-ray powder diffraction (XRD) using a Bruker D2-phaser diffractometer at Politecnico di Milano. Raman spectra were collected by a Horiba Jobin Yvon LabRAM HR Evolution spectrometer. The system is coupled to an Olympus BXFM petrographic microscope that works in transmitted and reflected light and is connected to a 5Mpx camera (objectives: 5X and 10X; LWD: 50X and 100X; spot size: ~1 μm). The laser in use was a green Nd laser (wavelength: 532 nm), characterized by a focal length of 800 mm and two grids (600 grooves/μm) and a CCD detector (1024 × 256 px, −60 °C). Most of the 9 filters (100%, 50%, 25%, 10%, 5%, 3%, 1%, 0.1%, and 0.01%) were used to diminish laser power, along with EDGE filter. Around 300 Raman spectra were randomly collected directly on the precipitate mounted on a microscope slide. We collected 28 to 41 spectra from each of the three mesocosms for each treatment (C, L, H) and treated them with LabSpec 5 software in order to identify the mineral phases.

Data Analysis

The pH, DIC, and TA data were analyzed to assess the quality of the measurements, monitor the stability of the mesocosms, and calculate the aragonite saturation state (Ω_{ar}) and the equilibrium partial pressure of CO₂ (pCO₂).

We applied statistical inference using control charts like those used in quality analysis of industrial processes.³³ Standard deviation charts, calculated from the three samplings of mesocosms at each time, were used to identify possible outliers in the measurements. Charts of the sample averages were instead used to identify the possible out-of-control status of the mesocosms upon the addition of Ca(OH)₂. For the constructions of the charts, the population standard deviations were calculated from the three control mesocosms, whereas the expected values (specification values) of pH, TA, and DIC were obtained from simulations of the mesocosms using the CO2SYS software,³⁴ given the known additions of Ca(OH)₂. The comparisons between the experimental values and the predictions clearly indicated that the re-equilibration of CO₂ with the atmosphere had not taken




Figure 2. Differences in (a) pH and (b) TA in the samples measured in the field and those delivered to Milano. The blue boxplot extends from the first to the third quartile. The orange line represents the median, and white circles represent outliers.

place (see below). Therefore, the control charts were calculated assuming mesocosms nonequilibrated with the atmosphere, i.e., with a constant DIC.

The averaged pH and TA data were used to calculate the evolution of Ω_{ar} and pCO_2 in the system. The data set, consisting of cleaned and averaged values, was analyzed using the CO2SYS Python library.³⁴ The software used constants from Lueker et al.³⁵ for carbonic acid dissociation, Dickson³⁶ for $KHSO_4$, and Uppström³⁷ for the total boron content (B_T). The calculation of Ω_{ar} in CO2SYS does not consider the variation of Ca^{2+} due to the dissolution of $Ca(OH)_2$ and precipitation of $CaCO_3$, so the value was corrected as suggested by Moras et al.¹⁶

$$[Ca^{2+}]_{corr} = [Ca^{2+}]_{salinity} + \frac{\Delta TA}{2}$$

where $[Ca^{2+}]_{salinity}$ is the calcium concentration calculated by CO2SYS from the salinity.

RESULTS AND DISCUSSION

Before diving into the discussion of the results, let us consider briefly the reliability and reproducibility of the experimental measurements. The standard deviation charts reported in the Supporting Information (SI) indicate good reproducibility among the three mesocosms within each treatment. Only a small number of measurements were identified as unreliable, flagging 7 outliers out of the 120 measurements of pH, DIC, and TA performed after the sample delivery for the L and H treatments. Figure 2 illustrates the differences in TA and pH, as measured in the field at HCMR immediately after sampling and filtration or delivered and subsequently analyzed in the laboratory at Politecnico di Milano. A decrease in pH and an increase in TA (3.35% on average) are observed in the latter. The simultaneous occurrence of these two phenomena is unusual, suggesting that they derive from separate processes. The pH decrease is likely due to CO_2 adsorption, possibly resulting from sample handling in different environments. On the other hand, the TA increase also suggests a rise in DIC but likely in the form of carbonate and bicarbonate ions. The alkalinities calculated from the pH and DIC were systematically higher (by 200–300 $\mu mol L^{-1}$) than those measured in the laboratory (see the Supporting Information). These differences are not negligible, and we hope to investigate their origin in future experiments. Nonetheless, these effects were consistent across all mesocosms, with no preferential treatment bias. The fact that both L and H treatments behaved similarly to the control samples reinforces the hypothesis that

these changes are due to external factors. Consequently, all samples were deemed reliable and analyzed in Milano, where DIC measurements were feasible.

Figure 3 presents the parameters measured in the laboratory at Politecnico di Milano. First of all, we observe that even the control mesocosms show variations over the time of the experiment, in particular a decrease in TA on days 2 and 6. This is remarkable, considering that the laboratory measurements tend to overestimate the TA by about 100 $\mu mol L^{-1}$, compared to those obtained in the field (see again Figure 2). Nonetheless, the L treatment closely follows the behavior of the control mesocosms and, unlike the H treatment, retains higher TA levels without any significant loss on the day after each addition. This also applies to the other measurements. In the L treatment, no significant variation of DIC was observed relative to the control, whereas in the H treatment, DIC displayed a strong decrease beginning on day 6, after the fourth addition.

The loss of stability in the H treatment is also evident in Figure 4, which compares the temporal evolution of the measured TA with the expected increase following alkalinity additions (see the SI for the other parameters). In the L treatment, the mean TA values remain within the control limits for most of the experimental period, indicating stable system behavior. In contrast, the mean TA values in the H mesocosms progressively deviate from the control limits after day 3, falling outside the error range and indicating progressive destabilization of the systems.

Both TA and DIC are key parameters for the marine carbon system. An increase in DIC indicates an uptake of atmospheric CO_2 , while TA is essential for detecting $CaCO_3$ precipitation, as it directly reflects the disappearance of CO_3^{2-} from the solution.⁹ In this experiment, precipitation was visually observed in both H and L mesocosms from the first addition, though to different extents. In fact, some precipitation also occurred in the C mesocosms. To assess this quantitatively, the increase in TA relative to the control was calculated as follows:

$$\Delta TA = TA_{X,i} - TA_{C,i} \quad (2)$$

where X refers to treatment type (L or H) and i denotes the sampling time (in days). A similar equation was applied to calculate ΔDIC , assuming that each mole of $Ca(OH)_2$ added can theoretically uptake 2 mol of CO_2 . The results for both parameters are presented in Figure 5.




Figure 3. (a) pH, (b) DIC, and (c) TA evolution, measured in Milano on samples from the high (H_M), low (L_M), and control (C_M) mesocosms. The lines are averages of the three replicates, and the colored areas represent the standard deviations. White symbols indicate time points with elevated standard deviation of the mean, as defined in the [Materials and Methods](#) section. Vertical lines signal the additions of Ca(OH)₂.

In the H treatment, TA reaches the expected increase 4 h after each addition but declines on the following day, indicating the precipitation of carbonates. Some precipitation was observed already after the first addition, but it initially involved only a minor portion of added TA.




Figure 4. Control charts of sample averages for the (a) L and (b) H mesocosms, showing the temporal evolution of the sample mean of TA values during the experiment. The central line is the bold solid line, while the thin lines indicate the upper and lower control limits (modified by the additions). The dots represent the observed sample means.

The H treatment exhibits a significant decrease in Δ DIC, most likely due to CaCO₃ precipitation, which removes both DIC and TA from the solution. In contrast, DIC levels in the L treatment remain stable, implying very limited precipitation but also little or no absorption of atmospheric CO₂. This behavior could be attributed to inefficient air exchange between the mesocosms and the atmosphere, likely caused by the small surface area available for gas exchange compared to the large volume of the mesocosms.

The lids placed on top of the mesocosms may have affected the CO₂ equilibration efficiency. This factor was not considered ahead of the experiments, as previous studies conducted at the HCMR facility showed stable oxygen levels in the mesocosms, and therefore, gas exchange was not expected to be significantly affected. However, it is important to note that O₂ concentrations in the atmosphere are much higher than CO₂ concentrations, meaning that while oxygen equilibration remained efficient, CO₂ exchange could have been substantially impacted by the lid. In addition, the bubbling applied for water mixing and aeration was intentionally very gentle in order to avoid disturbing the water column and the biological communities. While sufficient for its purpose, this low-intensity bubbling was not adequate to promote efficient CO₂ equilibration. Further investigation might be necessary to fully understand the impact of the lid on CO₂ exchange. Nevertheless, preventive measurements will be necessary in future experiments to avoid contamination of the oligotrophic seawater, ensuring the consistency of the data on marine biota collected by other research teams.

Figure 6 shows the behavior of Ω_{ar} and pCO₂ calculated from the field and laboratory data. The observed variations align with the previously observed TA increases and pH decreases (see Figure 2). In all mesocosms, the initial pCO₂




Figure 5. Evolution of (a) ΔTA and (b) ΔDIC . White symbols indicate points with large standard deviation of the mean. Vertical lines signal the additions of $\text{Ca}(\text{OH})_2$, while the dashed red and green lines represent the theoretical values that $\Delta\text{TA}_{\text{th}}$ and $\Delta\text{DIC}_{\text{th}}$ should reach according to the treatment.




Figure 6. Evolution of Ω_{ar} and pCO_2 , calculated from (a, b) field measurements and (c, d) measurements on delivered samples. Vertical lines signal the additions of $\text{Ca}(\text{OH})_2$, and white symbols indicate points with large standard deviation of the mean.

calculated from the field data was lower than the expected 420 ppm, likely due to restricted CO_2 equilibration in the mesocosms. In contrast, the pCO_2 calculated from the samples analyzed in the laboratory at Politecnico di Milano increased in a way that is consistent with the TA and DIC changes. In all cases, both L and H treatments exhibited lower pCO_2 levels than the control, suggesting that the system would have absorbed more CO_2 without the lids covering the mesocosms,

over a period lasting several days or possibly weeks. These lids were used to prevent contamination from external sources, which was particularly important for the biological measurements, although they may have reduced the efficiency of CO_2 equilibration with the atmosphere. Oxygen exchange was likely less affected by the presence of the lids because atmospheric O_2 concentrations are much higher than those of CO_2 . Increasing the air–water interface of the mesocosms could

further improve the gas exchange efficiency in future experiments.

To gain more insight, we also analyzed the sediments collected at the bottom of the mesocosm. Although some precipitation was visually observed in the L treatment, the amount of material collected at the end of the experiment was insufficient for XRD analysis. Therefore, only samples from H treatment were examined by this technique.

A total of three mineral phases were identified: aragonite, calcite, and halite. Excluding halite (NaCl), which is likely formed during drying of the samples, the other phases are carbonates. Aragonite was the dominant carbonate phase in all samples, accounting for 85–92% of the identified mineral content, whereas calcite was present only in minor proportions (2–5%). A small number of diffraction peaks remain unexplained, particularly in sample H2 (see the SI); however, their low intensity suggests the presence of additional mineral phases only in trace amounts. These phases could not be unambiguously identified and were therefore excluded from the quantitative analysis.

Interestingly, no brucite ($\text{Mg}(\text{OH})_2$) was detected. This is noteworthy, as in previous investigations, we detected its occurrence in both natural and artificial seawater.²² It is likely that magnesium hydroxide formed within the high-concentration slurries but redissolved upon dilution in the mesocosms, as suggested by Ringham et al.¹⁵

Raman spectroscopy was performed on the precipitates collected from all mesocosms. The repartition of the detected minerals is presented in Figure 8. In the C mesocosms, the




Figure 7. Repartition of minerals identified in the XRD analysis from the mesocosms H.

precipitate contained aragonite, calcite, and other minerals in varying proportions, including huntite ($\text{CaMg}_3(\text{CO}_3)_4$), cerussite (PbCO_3), and terrigenous minerals such as quartz, K-feldspar, plagioclase, muscovite, pyrite, magnetite, hematite, and titanite. In the L and H mesocosms, the spectra confirmed the XRD results showing a dominant presence of aragonite with smaller amounts of calcite, while the minerals detected in the control were not detected (see Figure 7 and the Supporting Information). No halite was detected, likely because of methodological limitations and the selection of minerals targeted for analysis.




Figure 8. Repartition of minerals identified in the Raman spectra from the different mesocosms. The category “Other” includes huntite, cerussite, quartz, K-feldspar, plagioclase, muscovite, pyrite, magnetite, hematite, and titanite.

CONCLUSION

We have carried out an experiment intended to test the effects of enhanced alkalization induced by liming. Studies were carried out at the Hellenic Centre for Marine Research (HCMR) in Heraklion (Crete, Greece), using mesocosms. A fundamental knowledge gap was the safe quantity of SL that can be added without triggering the unwanted precipitation of CaCO_3 , which implies a loss of efficiency of the liming process. The results of our experiment enabled us to assess that, for the specific conditions of Crete seawater, an SL addition corresponding to $120 \mu\text{mol L}^{-1}$ TA fulfills the “safety criterion”. Of course, this threshold cannot be considered universally valid because different seawater conditions and methods of $\text{Ca}(\text{OH})_2$ addition could generate localized pH spikes, potentially causing the precipitation of CaCO_3 or even $\text{Mg}(\text{OH})_2$.

Our experiment was not designed to monitor and optimize the uptake of atmospheric CO_2 by the treated seawater, but the results reported here imply that it can be a slow process requiring several days or weeks, even when alkalization is confined to a surface layer with a depth of a few meters. Further experimentation could help optimize delivery techniques for the alkaline plume, thereby enhancing the efficiency of CO_2 uptake by seawater. In particular, future mesocosm designs should aim to improve air-sea gas exchange, avoiding complete covering of the systems and preserving the air–water interface area. Also, it would be worthwhile to plan on-site measurements of the DIC (such as the pH and alkalinity). This would complicate the logistics and increase the cost of the experiments but would improve the overall quality and internal consistency of the data. Once these limitations are overcome, the protocol we have proposed here for experiments and for calculation of the efficiency of OL could also be adopted at other locations, producing a geographical map of safe SL addition thresholds.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acsestwater.6c00342>.

Raw experimental data and their elaboration (XLSX)

Additional data, including control charts, comparisons between measured and calculated TA, and XRD patterns (PDF)

AUTHOR INFORMATION

Corresponding Authors

Piero Macchi – Department of Chemistry, Materials and Chemical Engineering “G. Natta”, Politecnico di Milano, I-20131 Milano, Italy; orcid.org/0000-0001-6292-9825; Email: piero.macchi@polimi.it

Guido Raos – Department of Chemistry, Materials and Chemical Engineering “G. Natta”, Politecnico di Milano, I-20131 Milano, Italy; orcid.org/0000-0001-7011-4036; Email: guido.raos@polimi.it

Authors

Selene Varliero – Department of Chemistry, Materials and Chemical Engineering “G. Natta”, Politecnico di Milano, I-20131 Milano, Italy

Daniela Basso – Department of Earth and Environmental Sciences, University of Milano-Bicocca, 20126 Milano, Italy
Paraskevi Pitta – Institute of Oceanography, Hellenic Centre for Marine Research, 71003 Heraklion, Crete, Greece

Iordanis Magiopoulos – Institute of Oceanography, Hellenic Centre for Marine Research, 71003 Heraklion, Crete, Greece
Chiara Santinelli – CNR-IBF, Istituto di Biofisica, Area della Ricerca, I-56124 Pisa, Italy

Giovanni Checcucci – CNR-IBF, Istituto di Biofisica, Area della Ricerca, I-56124 Pisa, Italy

Silvia Valsecchi – Faculty of Life Sciences, University of Madeira, 9000-072 Funchal, Madeira Island, Portugal; Marine and Environmental Sciences Centre (MARE)/Aquatic Research Network (ARNET), Regional Agency for the Development of Research, Technology and Innovation (ARDITI), 9020-105 Funchal, Madeira Island, Portugal

Laura Borromeo – Department of Earth and Environmental Sciences, University of Milano-Bicocca, 20126 Milano, Italy

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acsestwater.6c00342>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

The authors would like to thank the staff at the HCMR Cretacosmos facility, part of the Greek National Research Infrastructure CMBR (Centre for the study and sustainable exploitation of Marine Biological Resources), for their valuable help with the mesocosm setup, installation, and runs, with special thanks to Anastasia Tsiola and Filomena Romano as well as the POSEIDON team for assistance with filling. The authors are grateful to Manolis Tsapakis and Eleni Dafnomili (HCMR) for their support in the inorganic nutrient analyses, and to Mirco Guerrazzi and Marco Carloni (CNR-IBF team of Pisa) for their support with sample processing. The authors thank Eleonora Barbaccia for kindly providing the photograph of the mesocosms in Figure 1. The research was funded by the Prince Albert II of Monaco Foundation in the framework of the OACIS Initiative (<https://www.fpa2.org/>), project no. 4000.1-1, and the EU H2020-INFRAIA project (No. 871081) AQUACOSM-plus: Network of Leading Ecosystem Scale Experimental AQUatic MesoCOSM Facilities Connecting

Rivers, Lakes, Estuaries and Oceans in Europe and beyond. The work was further supported by the Transnational Access program of the project AQUACOSM-plus that funded the participation of Selene Varliero, Giovanni Checcucci, Silvia Valsecchi, and Daniela Basso (TA proposal OLIGOS) in the experiment. Silvia Valsecchi is currently supported by a Doctoral Grant (024.04330.BD), financed by the Portuguese Foundation for Science and Technology (FCT).

REFERENCES

- (1) Trends in atmospheric concentrations of CO₂ (ppm), CH₄ (ppb) and N₂O (ppb), between 1800 and 2017. European Environmental Agency, 2024. <https://www.eea.europa.eu/en/analysis/maps-and-charts/atmospheric-concentration-of-carbon-dioxide-5?activeTab=6fbd444d-c422-4a78-8492-fd496bd61b7a> (accessed 2024-01-12).
- (2) Trends in CO₂. National Oceanic and Atmospheric Administration, 2024. <https://gml.noaa.gov/ccgg/trends/> (accessed 2024-01-12).
- (3) IPCC. Summary for Policymakers. *Climate Change 2022 - Mitigation of Climate Change: Working Group III Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*; Cambridge University Press, 2022.
- (4) IPCC. Summary for Policymakers. *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty*; Cambridge University Press, 2018; pp 3–24.
- (5) IPCC Chapter 12: Cross-sectoral perspectives. *Climate Change 2022 - Mitigation of Climate Change: Working Group III Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*; Cambridge University Press, 2022.
- (6) Middelburg, J. J.; Soetaert, K.; Hagens, M. Ocean alkalinity, buffering and biogeochemical processes. *Reviews of Geophysics* **2020**, 58, No. e2019RG000681.
- (7) Rohling, E. J. Marine methods for carbon dioxide removal: fundamentals and myth-busting for the wider community. *Oxford Open Climate Change* **2023**, 3, kgad004.
- (8) Ho, D. T.; Bopp, L. Marine carbon dioxide removal may be a future climate solution. *Dialogues on Climate Change* **2024**, 1, 56.
- (9) Zeebe, R. E.; Wolf-Gladrow, D. CO₂ in seawater: equilibrium, kinetics, isotopes; Gulf Professional Publishing, 2001.
- (10) Lee, C.-H.; Subhas, A. V.; Kim, J.-H.; Lee, K. Ocean Carbon Dioxide Removal and Storage. *Chem. Rev.* **2026**, 126, 1110–1144.
- (11) Decline in ocean pH measured at the Aloha station and yearly mean surface seawater pH reported on a global scale. European Environmental Agency, 2024. <https://www.eea.europa.eu/data-and-maps/figures/decline-in-ocean-ph-measured-4/> (accessed 2024-11-02).
- (12) Kheshgi, H. S. Sequestering atmospheric carbon dioxide by increasing ocean alkalinity. *Energy* **1995**, 20, 915–922.
- (13) Eisaman, M. D.; Geilert, S.; Renforth, P.; Bastianini, L.; Campbell, J.; Dale, A. W.; Foteinis, S.; Grasse, P.; Hawrot, O.; Löscher, C. R.; Rau, G. H.; Ronning, J. Assessing the technical aspects of ocean-alkalinity-enhancement approaches. *State of the Planet* **2023**, 2-0ae2023, 1.
- (14) Renforth, P.; Kruger, T. Coupling mineral carbonation and ocean liming. *Energy Fuels* **2013**, 27, 4199–4207.
- (15) Ringham, M. C.; Hirtle, N.; Shaw, C.; Lu, X.; Herndon, J.; Carter, B. R.; Eisaman, M. D. An assessment of ocean alkalinity enhancement using aqueous hydroxides: kinetics, efficiency, and precipitation thresholds. *Biogeosciences* **2024**, 21, 3551–3570.
- (16) Moras, C. A.; Bach, L. T.; Cyronak, T.; Joannes-Boyau, R.; Schulz, K. G. Ocean alkalinity enhancement - avoiding runaway CaCO₃ precipitation during quick and hydrated lime dissolution. *Biogeosciences* **2022**, 19, 3537–3557.

- (17) Hartmann, J.; Suitner, N.; Lim, C.; Schneider, J.; Marín-Samper, L.; Aristegui, J.; Renforth, P.; Taucher, J.; Riebesell, U. Stability of alkalinity in ocean alkalinity enhancement (OAE) approaches – consequences for durability of CO₂ storage. *Biogeosciences* **2023**, *20*, 781–802.
- (18) Schwinger, J.; Bourgeois, T.; Rickels, W. On the emission-path dependency of the efficiency of ocean alkalinity enhancement. *Environmental Research Letters* **2024**, *19*, 074067.
- (19) Jeltsch-Thömmes, A.; Tran, G.; Lienert, S.; Keller, D. P.; Oschlies, A.; Joos, F. Earth system responses to carbon dioxide removal as exemplified by ocean alkalinity enhancement: tradeoffs and lags. *Environmental Research Letters* **2024**, *19*, 054054.
- (20) Bach, L. T. The additionality problem of ocean alkalinity enhancement. *Biogeosciences* **2024**, *21*, 261–277.
- (21) Suitner, N.; Faucher, G.; Lim, C.; Schneider, J.; Moras, C. A.; Riebesell, U.; Hartmann, J. Ocean alkalinity enhancement approaches and the predictability of runaway precipitation processes: results of an experimental study to determine critical alkalinity ranges for safe and sustainable application scenarios. *Biogeosciences* **2024**, *21*, 4587–4604.
- (22) Varliero, S.; Buono, A.; Caserini, S.; Raos, G.; Macchi, P. Chemical Aspect of Ocean Liming for CO₂ Removal: Dissolution Kinetics of Calcium Hydroxide in Seawater. *ACS Engineering Au* **2024**, *4*, 422.
- (23) Varliero, S.; Jamali Alamooti, S.; Campo, F. P.; Cappello, G.; Cappello, S.; Caserini, S.; Comazzi, F.; Macchi, P.; Raos, G. Assessing the Limit of CO₂ Storage in Seawater as Bicarbonate-Enriched Solutions. *Molecules* **2024**, *29*, 4069.
- (24) Jamali Alamooti, S.; Comazzi, F.; Thaler, E. K.; Gropelli, S.; Calvi, D.; Raos, G.; Macchi, P. pH-equilibrated ocean alkalization: Mesoscale evaluation of long-term stability. *International Journal of Greenhouse Gas Control* **2026**, *151*, 104589.
- (25) Basso, D.; et al. Ocean liming in eutrophic water: a mesocosm scale approach. *EGU General Assembly 2023*, 2023. DOI: [10.5194/egusphere-egu23-10342](https://doi.org/10.5194/egusphere-egu23-10342).
- (26) Serret, P.; et al. The impact of ocean liming on phytoplankton size-structure and the balance of photosynthesis and respiration in two contrasting environments. *EGU General Assembly 2024*, 2024. DOI: [10.5194/egusphere-egu24-13093](https://doi.org/10.5194/egusphere-egu24-13093).
- (27) Tselepidis, A.; Zervakis, V.; Polychronaki, T.; Danovaro, R.; Chronis, G. Distribution of nutrients and particulate organic matter in relation to the prevailing hydrographic features of the Cretan Sea (NE Mediterranean). *Progress in Oceanography* **2000**, *46*, 113–142.
- (28) Nocera, A. C.; Traboni, C.; Courboulès, J.; Romano, F.; Chantzaras, C.; Magiopoulos, I.; Varliero, S.; Azzellino, A.; Bazzicalupo, P.; Macchi, P.; Raos, G.; Basso, D.; Pitta, P. The effect of ocean alkalinity enhancement on zooplankton standing stock and community composition in the Eastern Mediterranean Sea: a mesocosm study. *Marine Environmental Research* **2025**, *212*, 107489.
- (29) Caserini, S.; Pagano, D.; Campo, F.; Abbà, A.; De Marco, S.; Righi, D.; Renforth, P.; Grosso, M. Potential of maritime transport for ocean liming and atmospheric CO₂ removal. *Frontiers in Climate* **2021**, *3*, 575900.
- (30) Badocco, D.; Pedrini, F.; Pastore, A.; di Marco, V.; Marin, M. G.; Bogialli, S.; Roverso, M.; Pastore, P. Use of a simple empirical model for the accurate conversion of the seawater pH value measured with NIST calibration into seawater pH scales. *Talanta* **2021**, *225*, 122051.
- (31) Lewis, E.; Perkin, R. The practical salinity scale 1978: conversion of existing data. *Deep Sea Research Part A. Oceanographic Research Papers* **1981**, *28*, 307–328.
- (32) ASTM D7573-18ae1: Standard Test Method for Total Carbon and Organic Carbon in Water by High Temperature Catalytic Combustion and Infrared Detection. *Book of Standards*; ASTM, 2019; Vol. 11.02. DOI: [10.1520/D7573-18AE01](https://doi.org/10.1520/D7573-18AE01).
- (33) Montgomery, D. C. *Introduction to Statistical Quality Control*; Wiley, 2020.
- (34) Humphreys, M. P.; Lewis, E. R.; Sharp, J. D.; Pierrot, D. PyCO2SYS v1.8: marine carbonate system calculations in Python. *Geoscientific Model Development* **2022**, *15*, 15–43.
- (35) Lueker, T. J.; Dickson, A. G.; Keeling, C. D. Ocean pCO₂ calculated from dissolved inorganic carbon, alkalinity, and equations for K₁ and K₂: validation based on laboratory measurements of CO₂ in gas and seawater at equilibrium. *Marine chemistry* **2000**, *70*, 105–119.
- (36) Dickson, A. G. Standard potential of the reaction: AgCl(s) + 12H₂(g) = Ag(s) + HCl(aq), and the standard acidity constant of the ion in synthetic sea water from 273.15 to 318.15 K. *J. Chem. Thermodyn.* **1990**, *22*, 113–127.
- (37) Uppström, L. R. The boron/chlorinity ratio of deep-sea water from the Pacific Ocean. *Deep Sea Research and Oceanographic Abstracts* **1974**, *21*, 161–162.